

ON WEAKLY MAXIMAL REPRESENTATIONS OF SURFACE GROUPS

G. BEN SIMON, M. BURGER, T. HARTNICK, A. IOZZI, AND A. WIENHARD

ABSTRACT. We introduce and study a new class of representations of surface groups into Lie groups of Hermitian type, called *weakly maximal* representations. We prove that weakly maximal representations are discrete and injective and we describe the structure of the Zariski closure of their image. Furthermore we prove that the set of weakly maximal representations is a closed subset of the representation variety and describe its relation to other geometrically significant subsets of the representations variety.

1. INTRODUCTION

Given a compact oriented surface Σ of negative Euler characteristic, possibly with boundary, a general theme is to study the space of representations $\text{Hom}(\pi_1(\Sigma), G)$ of the fundamental group $\pi_1(\Sigma)$ of Σ into a semisimple Lie group G , and in particular to distinguish subsets of geometric significance.

In recent years these studies led to the discovery of new subsets of representation varieties: Hitchin components [38, 18, 39, 30, 31], positive representations [21, 22], maximal representations [26, 27, 42, 37, 15, 12, 10, 11, 44, 36, 28, 3, 2, 4, 23] and Anosov representations [33, 39, 32]. Even though these subsets are defined and investigated by very different methods, they exhibit several common properties, and their study is summarized under the terminology *higher Teichmüller theory*.

Date: December 25, 2014.

M. B. was partial supported by the Swiss National Science Foundation project 200020-144373; T. H. was partial supported by the Swiss National Science Foundation project 2000021-127016/2; A. I. was partial supported by the Swiss National Science Foundation projects 2000021-127016/2 and 200020-144373; A. W. was partially supported by the National Science Foundation under agreement No. DMS-1065919 and 0846408, by the Sloan Foundation, by the Deutsche Forschungsgemeinschaft, and by the ERCEA under ERC-Consolidator grant no. 614733. Support by the Institut Mittag-Leffler (Djursholm, Sweden) and by the Institute for Advanced Study (Princeton, NJ) is gratefully acknowledged.

Here we introduce a new class of representations of $\pi_1(\Sigma)$ into a Lie group of Hermitian type, the set of *weakly maximal representations*. We establish several results showing that weakly maximal representations are of geometric significance. We also construct various nontrivial examples of weakly maximal representations, establish a relation to orders on Lie groups, and discuss the relation of the set of weakly maximal representations with other subsets of the representation variety.

Weakly maximal representations with non-zero Toledo number also admit a geometric characterization. This geometric characterization is discussed in detail in the companion paper [1]. We show there that weakly maximal representations with non-zero Toledo number are representations that are order preserving in an appropriate sense with respect to an arbitrary bi-invariant continuous partial order on the cyclic covering on G . If the symmetric space associated to G is a Hermitian symmetric space of tube type, they are also order-preserving with respect to the order induced by the Kaneyuki causal ordering on the corresponding Shilov boundary. This leads to a particular simple geometric characterization in the tube type case.

1.1. Structure Theorems for Weakly Maximal Representations.

Recall that the Toledo invariant $T(\rho)$ of a representation $\rho : \pi_1(\Sigma) \rightarrow G$ is defined as the evaluation, in an appropriate sense, of the pullback $\rho^*(\kappa_G^b)$ of the bounded Kähler class $\kappa_G^b \in H_{cb}^2(G, \mathbb{R})$ on the relative fundamental class $[\Sigma, \partial\Sigma]$ (see § 2 for details). This definition in terms of bounded cohomology leads to a chain of inequalities

$$|T(\rho)| \leq 2\|\rho^*(\kappa_G^b)\| \cdot |\chi(\Sigma)| \leq 2\|\kappa_G^b\| \cdot |\chi(\Sigma)|,$$

where $\|\cdot\|$ denotes the canonical norm on the Banach spaces $H_{cb}^2(G, \mathbb{R})$ and $H_b^2(\pi_1(\Sigma), \mathbb{R})$, while $|\chi(\Sigma)|$ appears as the ℓ^1 -simplicial norm of $[\Sigma, \partial\Sigma]$. *Maximal representations* are those for which both inequalities are equalities, while representations for which the second inequality is an equality are *tight homomorphisms*; these constitute a very interesting class of homomorphisms which can be defined and studied in much greater generalities, see [14, 34, 35]. In this article we set ourselves the goal to analyse the structure of representations for which the first inequality is an equality.

DEFINITION 1.1 ([43, Chapter 8, Definition 2.1]). A representation $\rho : \pi_1(\Sigma) \rightarrow G$ is *weakly maximal* if it satisfies

$$(1.1) \quad T(\rho) = 2\|\rho^*(\kappa_G^b)\| \cdot |\chi(\Sigma)|.$$

REMARK 1.2. By definition a weakly maximal representation has non-negative Toledo number. Analogously one can define weakly minimal

representations. Since the composition of a representation with an orientation reversing outer automorphism changes the sign of the Toledo number the theory is completely analogous.

Whereas for maximal representations the Toledo invariant is a fixed number independent of ρ , for weakly maximal representations the right hand side of the equality is $2\|\rho^*(\kappa_G^b)\| \cdot |\chi(\Sigma)|$ and depends on ρ . In particular, there is (a priori) no restriction on the Toledo invariant of a weakly maximal representation.

In order to get a better understanding of weakly maximal representations it is therefore important to give alternative characterizations of weakly maximal representations. One such characterization is that a representation ρ is weakly maximal if and only if there exists $\lambda \geq 0$ such that

$$(1.2) \quad \rho^*(\kappa_G^b) = \lambda \kappa_\Sigma^b,$$

where κ_Σ^b is the bounded fundamental class of Σ (see (3.1)). This characterization was in effect first established in [43, Corollary 3.4] for compact surfaces and in [15, Cor. 4.15] in the general case; a different proof using Bavard's duality was later obtained in [17] (for a more thorough description of the relation with Calegari's work, see [11]).

The constant λ appearing in (1.2) is related to the Toledo invariant by

$$T(\rho) = \lambda |\chi(\Sigma)|.$$

When Σ is a surface without boundary the Toledo invariant is a characteristic number with values in $q_G^{-1}\mathbb{Z}$, where q_G is a natural number depending only on G (namely, q_G is the smallest integer such that $q_G \kappa_G^b$ is an integral class, see Remark 6.5). On the other hand, when Σ is a surface with boundary, the Toledo invariant ranges over the whole closed interval $[-2\|\kappa_G^b\| \cdot |\chi(\Sigma)|, 2\|\kappa_G^b\| \cdot |\chi(\Sigma)|]$.

It is remarkable that for weakly maximal representations we can restrict the possible values of the Toledo-invariant by the following

THEOREM 1.3 (Rationality Theorem). *There is a natural number ℓ_G depending only on G , such that for every weakly maximal representation $\rho : \pi_1(\Sigma) \rightarrow G$ we have $T(\rho) \in \frac{|\chi(\Sigma)|}{\ell_G} \mathbb{Z}$. In particular, T takes only finitely many values on the set of weakly maximal representations.*

REMARK 1.4. The integer ℓ_G depends in an explicit way on q_G and on the degree of non-integrality of the restriction $\kappa_G^b|_H$ to various connected semisimple subgroups H of G , as well as on the cardinality of their center (see the proof of Corollary 6.4).

The Rationality Theorem is an essential ingredient in order to prove

THEOREM 1.5. *Let $\rho : \pi_1(\Sigma) \rightarrow G$ be a weakly maximal representation and $T(\rho) \neq 0$. Then ρ is faithful with discrete image.*

The methods of proof for this theorem are very different from those used in the case of maximal representations [15] and give in particular a new and simpler proof of the injectivity and discreteness of maximal representations.

Whereas the Zariski closure for maximal representations is always reductive and subject to strong restrictions, the Zariski closure of the image of a weakly maximal representation can be quite wild. In particular, it need not be reductive (see § 9.6). However, we can establish a structure theorem describing the Zariski closure of weakly maximal representations as follows. Given a closed subgroup $L < G$, we show that there exists a unique maximal normal subgroup of L , that we call the *Kähler radical* $\text{Rad}_{\kappa_G^b}(L)$ of L , on which $\kappa_G^b|_L$ vanishes. Since $\text{Rad}_{\kappa_G^b}(L)$ contains the solvable radical of L , the quotient $L/\text{Rad}_{\kappa_G^b}(L)$ is semisimple. We then establish the following as part of Theorem 7.2:

THEOREM 1.6. *Let $\rho : \pi_1(\Sigma) \rightarrow G$ be a weakly maximal representation into a Lie group $G := \mathbf{G}(\mathbb{R})^\circ$ of Hermitian type, where \mathbf{G} is a connected algebraic group defined over \mathbb{R} . Let $L := \mathbf{L}(\mathbb{R})$, where \mathbf{L} is the Zariski closure of $\rho(\pi_1(\Sigma))$ in \mathbf{G} , and let $H = L/\text{Rad}_{\kappa_G^b}(L)$ be the quotient of L by its Kähler radical. Assume that $T(\rho) \neq 0$. Then:*

- (1) *H is adjoint of Hermitian type and all of its simple factors are of tube type.*
- (2) *The composition $\pi_1(\Sigma) \rightarrow L \rightarrow H$ is faithful with discrete image.*

Recall that a Hermitian Lie group is of *tube type* if the associated symmetric space is of tube type, that is biholomorphic to $\mathbb{R}^n + iC$, for some $n \in \mathbb{N}$, where $C \subset \mathbb{R}^n$ is an open convex cone.

REMARK 1.7. In the above Theorems 1.5 and 1.6 it is essential that the Toledo invariant is non-zero. However the class of weakly maximal representations with $T(\rho) = 0$ is also of interest. This is precisely the set where $\rho^*(\kappa_G^b) = 0$. In the case when $G = \text{PU}(1, n)$ such representations have been studied and characterized in [9] as representations that preserve a totally real subspace of complex hyperbolic space $\mathcal{H}_\mathbb{C}^n$.

1.2. Comparison with Other Classes of Representations. Consider the representation variety $\text{Hom}(\pi_1(\Sigma), G)$.

The subset of weakly maximal representations $\text{Hom}_{wm}(\pi_1(\Sigma), G)$ decomposes as a disjoint union

$$\text{Hom}_{wm}(\pi_1(\Sigma), G) = \text{Hom}_{wm}^*(\pi_1(\Sigma), G) \sqcup \text{Hom}_0(\pi_1(\Sigma), G)$$

of the set $\text{Hom}_{wm}^*(\pi_1(\Sigma), G)$ of weakly maximal representations with non-zero Toledo invariant and the set of representations with vanishing Toledo invariant

$$\text{Hom}_0(\pi_1(\Sigma), G) := \{\rho : \pi_1(\Sigma) \rightarrow G : \rho^*(\kappa_G^b) = 0\}.$$

The latter contains $\text{Hom}(\pi_1(\Sigma), L)$ for every closed subgroup $L < G$ for which $\kappa_G^b|_L = 0$. In particular, representations in $\text{Hom}_0(\pi_1(\Sigma), G)$ are not necessarily injective, and their images are not necessarily discrete (see Remark 1.7), while, according to Theorem 1.5, $\text{Hom}_{wm}^*(\pi_1(\Sigma), G)$ is contained in the set $\text{Hom}_{d,i}(\pi_1(\Sigma), G)$ of injective homomorphisms with discrete image. We prove (cf. Corollary 8.4):

THEOREM 1.8. *The following*

$$\begin{array}{ccc} & & \text{Hom}_{d,i}(\pi_1(\Sigma), G) \\ & \nearrow & \\ \text{Hom}_{max}(\pi_1(\Sigma), G) \hookrightarrow & \text{Hom}_{wm}^*(\pi_1(\Sigma), G) & \\ & \searrow & \\ & & \text{Hom}_{wm}(\pi_1(\Sigma), G) \end{array}$$

is a diagram of $\text{Aut}(\pi_1(\Sigma))$ -invariant **closed** subsets of $\text{Hom}(\pi_1(\Sigma), G)$.

If G is real algebraic, $\text{Hom}_{max}(\pi_1(\Sigma), G)$ is a real semi-algebraic subset of $\text{Hom}(\pi_1(\Sigma), G)$, [15, Corollary 14], but we do not have such precise information on the other sets appearing in Theorem 1.8.

A prominent role in higher Teichmüller theory is played by Anosov representations, a notion introduced by F. Labourie in his study of Hitchin representations [39], then studied for Hermitian Lie groups in [10, 12] and in greater generality in [33]. The property of a representation to be Anosov is defined with respect to a parabolic subgroup $P < G$. In the context of weakly maximal representations the interest lies on representations $\rho : \pi_1(\Sigma) \rightarrow G$ which are Anosov with respect to the parabolic group Q which is the stabilizer in G of a point in the Shilov boundary \check{S} of the bounded symmetric domain realization of \mathcal{X} [10, 12, 33]. We will call such an Anosov representation *Shilov-Anosov*.

In the case in which $\partial\Sigma = \emptyset$, if we denote by $\text{Hom}_{\check{S}-An}(\pi_1(\Sigma), G)$ the set of such representations and by $\text{Hom}_{\check{S}-An}^*(\pi_1(\Sigma), G)$ the subset of Shilov-Anosov representations with positive Toledo invariant, we have

$$\text{Hom}_{max}(\pi_1(\Sigma), G) \hookrightarrow \text{Hom}_{\check{S}-An}^*(\pi_1(\Sigma), G) \hookrightarrow \text{Hom}_{\check{S}-An}(\pi_1(\Sigma), G),$$

where $\text{Hom}_{max}(\pi_1(\Sigma), G)$ is a union of components and $\text{Hom}_{\check{S}-An}(\pi_1(\Sigma), G)$ is an open subset of $\text{Hom}(\pi_1(\Sigma), G)$, [10, 12, 33].

On the other hand we prove in Corollary 8.5 the following:

THEOREM 1.9. *If G is of tube type and $\partial\Sigma = \emptyset$, then*

$$\overline{\text{Hom}_{\check{S}-An}(\pi_1(\Sigma), G)} \hookrightarrow \text{Hom}_{wm}(\pi_1(\Sigma), G).$$

If $G = \text{Sp}(2n, \mathbb{R})$ with $n \geq 2$, we will see that $\text{Hom}_{\check{S}-An}(\pi_1(\Sigma), G)$ is open but not closed. The study of the closure of the set of Anosov representations is a completely open problem. Since the set of weakly maximal representations is closed (see Corollary 8.3), it provides a natural framework in which to study limits of Shilov-Anosov representations. We refer the reader to § 9.5 for examples of weakly maximal representations into $\text{Sp}(2n, \mathbb{R})$ for $n \geq 6$, that are not Shilov-Anosov, but are the limit of Shilov-Anosov representations.

For a closed surface Σ and a Hermitian group G of tube type we summarize the relation of various special subsets of the representation variety and results from [39, 14] in the following diagram. Here we denote by $\text{Hom}_{red}(\pi_1(\Sigma), G)$ the set of representations with reductive Zariski closure and by $\text{Hom}_{Hitchin}(\pi_1(\Sigma), G)$ the Hitchin components in case G is locally isomorphic to $\text{Sp}(2n, \mathbb{R})$ (and the empty set otherwise). Then we have the following inclusions:

$$\begin{array}{ccccccc}
 & & & \text{Hom}_{\check{S}-An} & \hookrightarrow & \text{Hom}_{wm} & \\
 & & & \uparrow & & \uparrow & \\
 & & & \text{Hom}_{\check{S}-An}^* & \hookrightarrow & \text{Hom}_{wm}^* & \hookrightarrow \text{Hom}_{d,i} \\
 \text{Hom}_{Hitchin} & \hookrightarrow & \text{Hom}_{max} & \hookrightarrow & \text{Hom}_{\check{S}-An}^* & \hookrightarrow & \text{Hom}_{wm}^* & \hookrightarrow & \text{Hom}_{d,i} \\
 & & \searrow & & & & & & \\
 & & & \text{Hom}_{tight} & \hookrightarrow & \text{Hom}_{red} & & &
 \end{array}$$

2. THE TOLEDO INVARIANT

We recall here the definition of the Toledo invariant in a general context and indicate how the Milnor–Wood inequality follows from known isometric isomorphisms in bounded cohomology.

Let Σ be a compact oriented surface with (possibly empty) boundary $\partial\Sigma$. Let G be a locally compact group, $\kappa \in H_{cb}^2(G, \mathbb{R})$ a fixed continuous bounded class and $\rho : \pi_1(\Sigma) \rightarrow G$ a homomorphism. Recall that there is an isometric isomorphism

$$g_\Sigma : H_b^2(\pi_1(\Sigma), \mathbb{R}) \longrightarrow H_b^2(\Sigma, \mathbb{R})$$

whose existence in general is due to Gromov [29]; in our case, the universal covering $\tilde{\Sigma}$ is contractible and the existence and isometric property of g_Σ are easily established. Next, the inclusion of pairs $(\Sigma, \emptyset) \hookrightarrow (\Sigma, \partial\Sigma)$ gives rise to a map

$$j_{\partial\Sigma} : H_b^2(\Sigma, \partial\Sigma, \mathbb{R}) \longrightarrow H_b^2(\Sigma, \mathbb{R})$$

where the left hand side refers to bounded relative cohomology. Since every connected component of $\partial\Sigma$ is a circle and hence has amenable fundamental group, the map $j_{\partial\Sigma}$ is an isometric isomorphism, [5, Theorem 1]. We hence define

$$T_\kappa(\rho) := \langle (j_{\partial\Sigma})^{-1} g_\Sigma(\rho^*(\kappa)), [\Sigma, \partial\Sigma] \rangle,$$

where $[\Sigma, \partial\Sigma]$ is the relative fundamental class. Since g_Σ and $j_{\partial\Sigma}$ are isometries, we deduce that

$$|T_\kappa(\rho)| \leq \|\rho^*(\kappa)\| \|[\Sigma, \partial\Sigma]\|_1,$$

where the second factor refers to the norm in relative ℓ^1 -homology. Since $\|[\Sigma, \partial\Sigma]\|_1 = 2|\chi(\Sigma)|$ and the pullback is norm decreasing, then

$$(2.1) \quad |T_\kappa(\rho)| \leq 2\|\rho^*(\kappa)\| |\chi(\Sigma)| \leq 2\|\kappa\| |\chi(\Sigma)|.$$

In view of (2.1), we give the following definition:

DEFINITION 2.1. A representation $\rho : \pi_1(\Sigma) \rightarrow G$ is *κ -weakly maximal* if $T_\kappa(\rho) = \|\rho^*(\kappa)\| |\chi(\Sigma)|$.

REMARK 2.2. If G is of Hermitian type and $\kappa = \kappa_G^b$ is the bounded Kähler class, then $\|\kappa_G^b\| = \text{rank}(G)/2$ (see [15, § 2.1]) and one obtains for the corresponding Toledo invariant $T(\rho)$ the familiar Milnor–Wood inequality

$$|T(\rho)| \leq \text{rank}(G) |\chi(\Sigma)|.$$

This will be used in the examples in § 9, but otherwise never in the paper the explicit computation of the norm is used.

3. A CHARACTERIZATION OF WEAKLY MAXIMAL REPRESENTATIONS

In this section we develop the general framework for the study of weakly maximal representations and establish some of their basic properties. In particular we show that weak maximality of a representation $\rho : \pi_1(\Sigma) \rightarrow G$ is reflected in a relation between its lift to appropriate central extensions of $\pi_1(\Sigma)$ and G and canonically defined quasimorphisms on those central extensions (cf. Proposition 3.2).

3.1. The Central Extension of $\pi_1(\Sigma)$. Let h be any complete hyperbolic structure on the interior Σ° of Σ compatible with the fixed orientation and let $\rho_h : \pi_1(\Sigma) \rightarrow \mathrm{PU}(1, 1)$ be the corresponding holonomy representation. Then

$$(3.1) \quad \kappa_\Sigma^b := \rho_h^*(\kappa_{\mathrm{PU}(1,1)}^b) \in H_b^2(\pi_1(\Sigma), \mathbb{R})$$

is independent of the choice of h and is called the *(real) bounded fundamental class* of Σ , [15, § 8.2]. If $\widehat{\Gamma}$ denotes the central extension

$$(3.2) \quad 0 \longrightarrow \mathbb{Z} \longrightarrow \widehat{\Gamma} \xrightarrow{p_\Sigma} \pi_1(\Sigma) \longrightarrow 0$$

corresponding to the positive generator of $H^2(\pi_1(\Sigma), \mathbb{Z})$ if $\partial\Sigma = \emptyset$ and $\widehat{\Gamma} = \widetilde{\pi_1(\Sigma)}$ otherwise, then ρ_h lifts to a homomorphism $\widetilde{\rho}_h : \widehat{\Gamma} \rightarrow \widetilde{\mathrm{PU}(1, 1)}$, where we think of the universal covering $\widetilde{\mathrm{PU}(1, 1)}$ as contained in the group $\mathrm{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$ of increasing homeomorphisms of the real line \mathbb{R} commuting with integer translations. Denote by $\tau : \widetilde{\mathrm{PU}(1, 1)} \rightarrow \mathbb{R}$ the Poincaré translation quasimorphism defined by $\tau(f) := \lim_{n \rightarrow \infty} \frac{f^n(x) - x}{n}$ for $x \in \mathbb{R}$. Then the composition $\tau \circ \widetilde{\rho}_h$ is a homogeneous quasimorphism which is \mathbb{Z} -valued

$$(3.3) \quad \tau \circ \widetilde{\rho}_h : \widehat{\Gamma} \rightarrow \mathbb{Z},$$

since every element in the image of ρ_h has a fixed point. Moreover, the homogeneous quasimorphism τ corresponds in real bounded cohomology to the pullback, via the projection $p : \widetilde{\mathrm{PU}(1, 1)} \rightarrow \mathrm{PU}(1, 1)$, of the real bounded cohomology class $\kappa_{\mathrm{PU}(1,1)}^b$

$$(3.4) \quad [d\tau] = p^*(\kappa_{\mathrm{PU}(1,1)}^b),$$

and hence, again in real bounded cohomology,

$$(3.5) \quad [d(\tau \circ \widetilde{\rho}_h)] = p_\Sigma^*(\kappa_\Sigma^b).$$

While $\tau \circ \widetilde{\rho}_h$ depends on the hyperbolization ρ_h and on its lift $\widetilde{\rho}_h$, its restriction $(\tau \circ \widetilde{\rho}_h)|_\Lambda$ to $\Lambda := [\widehat{\Gamma}, \widehat{\Gamma}]$ is independent of the above choices, (as the equivalence in the Proposition 3.2 below shows).

3.2. The Central Extension for a Locally Compact Group G .

We now turn to the construction of the central extension (depending on a “rational” class) and of the associated quasimorphism for a general locally compact group. The relation with § 3.1 is outlined in § 3.3.

DEFINITION 3.1. We say that a bounded cohomology class κ is *rational* if there is an integer $n \geq 1$ such that $n\kappa$ is representable by a bounded \mathbb{Z} -valued Borel cocycle.

Let κ be rational and let $c : G^2 \rightarrow \mathbb{Z}$ be a normalized bounded Borel cocycle representing $n\kappa$. Endow $G \times \mathbb{Z}$ with the group structure defined by the Borel map

$$(g_1, n_1)(g_2, n_2) := (g_1g_2, n_1 + n_2 + c(g_1, g_2))$$

and let $G_{n\kappa}$ denote the Borel group $G \times \mathbb{Z}$ endowed with the unique compatible locally compact group topology [40]. This gives the topological central extension

$$(3.6) \quad 0 \longrightarrow \mathbb{Z} \xrightarrow{i} G_{n\kappa} \xrightarrow{p_{n\kappa}} G \longrightarrow e.$$

Then $f'_{n\kappa}(g, m) = \frac{1}{n}m$ is a Borel quasimorphism such that $df'_{n\kappa}$ represents $p_{n\kappa}^*(\kappa)$. Its homogenization $f_{n\kappa} : G_{n\kappa} \rightarrow \mathbb{R}$ is a continuous homogeneous quasimorphism [15, Lemma 7.4] such that

$$(3.7) \quad [df_{n\kappa}] = p_{n\kappa}^*(\kappa),$$

and

$$(3.8) \quad f_{n\kappa}(i(m)) = \frac{1}{n}m.$$

The homogeneous quasimorphisms $f_{n\kappa}$ are well defined in the following sense. Let $\zeta = n\kappa$ be the smallest integer multiple of κ representable by a \mathbb{Z} -valued bounded Borel cocycle and let $\ell \in \mathbb{Z}$, $\ell \neq 0$. Then the map $G \times \mathbb{Z} \rightarrow G \times \mathbb{Z}$ defined by $(g, m) \mapsto (g, \ell m)$ induces a continuous homomorphism that identifies G_ζ with a closed subgroup of finite index in $G_{\ell\zeta}$. Via this identification, the quasimorphism f_ζ is the restriction to G_ζ of $f_{\ell\zeta}$.

The following result describes the property of being weakly maximal in terms of quasimorphisms and will be essential in the sequel.

PROPOSITION 3.2. *Let G be a locally compact group and $\kappa \in H_b^2(G, \mathbb{R})$ a rational class with $n\kappa$ integral. Let $\rho : \pi_1(\Sigma) \rightarrow G$ be a homomorphism and $\tilde{\rho} : \widehat{\Gamma} \rightarrow G_{n\kappa}$ some lift, where $\widehat{\Gamma}$ and $G_{n\kappa}$ are defined respectively in (3.2) and (3.6). The following are equivalent:*

- (1) ρ is κ -weakly maximal;
- (2) there exists $\lambda \geq 0$ such that $\rho^*(\kappa) = \lambda\kappa_\Sigma^b$;
- (3) there exists $\lambda \geq 0$ and $\psi \in \text{Hom}(\widehat{\Gamma}, \mathbb{R})$ such that

$$f_{n\kappa} \circ \tilde{\rho} = \lambda(\tau \circ \tilde{\rho}_h) + \psi;$$

- (4) there exists $\lambda \geq 0$ such that

$$(f_{n\kappa} \circ \tilde{\rho})|_\Lambda = \lambda(\tau \circ \tilde{\rho}_h)|_\Lambda.$$

REMARK 3.3. The constant $\lambda \geq 0$ that appears in Proposition 3.2 is nothing but $\lambda = \frac{T_\kappa(\rho)}{|\chi(\Sigma)|}$, as it can be easily seen from the definition of the Toledo invariant in § 2.

Proof of Proposition 3.2. The equivalence of (1) and (2) is [11, Corollary 4.15].

Observe that, modulo modifying the lift $\tilde{\rho}$ by a homomorphism with values in $\mathbb{Z} \hookrightarrow G_{n\kappa}$, the diagram

$$(3.9) \quad \begin{array}{ccc} \widehat{\Gamma} & \xrightarrow{\tilde{\rho}} & G_{n\kappa} \\ p_\Sigma \downarrow & & \downarrow p_{n\kappa} \\ \pi_1(\Sigma) & \xrightarrow{\rho} & G \end{array}$$

commutes. This, together with the relation (3.5) and the fact that p_Σ^* induces an isomorphism in real bounded cohomology, proves the equivalence of (2) and (3).

Finally, by taking coboundaries on both sides of the equality in (4), one obtains that the bounded classes $[d(f_{n\kappa} \circ \tilde{\rho})]$ and $\lambda[d(\tau \circ \tilde{\rho}_h)]$ coincide on Λ . Since $\widehat{\Gamma}/\Lambda$ is amenable, we deduce the equality in $\widehat{\Gamma}$, which in turns implies (3). \square

3.3. The Hermitian Case. Assume that G is of Hermitian type and almost simple. We apply the above discussion to the bounded Kähler class κ_G^b , which is rational, [7, § 5]. If $\kappa = n\kappa_G^b$ is any integer multiple represented by an integral cocycle, the topological central extension

$$0 \longrightarrow \mathbb{Z} \xrightarrow{i} G_\kappa \xrightarrow{p_\kappa} G \longrightarrow e$$

is not trivial. Since $\pi_1(G)$ is isomorphic to \mathbb{Z} modulo torsion, there is a unique connected central \mathbb{Z} -extension \widehat{G} and, as a result, the connected component of the identity $(G_\kappa)^\circ$ is isomorphic to \widehat{G} . We denote by $f_{\widehat{G}} : \widehat{G} \rightarrow \mathbb{R}$ the continuous homogeneous quasimorphism corresponding to f_κ under this isomorphism and hence obtain that

$$(3.10) \quad [df_{\widehat{G}}] = p^*(\kappa_G^b),$$

where $p = p|_{(G_\kappa)^\circ} : \widehat{G} \rightarrow G$ is the projection. For example, if $G = \mathrm{PU}(1, 1)$, the bounded Kähler class $\kappa_{\mathrm{PU}(1,1)}^b$ is already integral as it is the image of the bounded Euler class e^b under the change of coefficients $H_{\mathrm{cb}}^2(\mathrm{PU}(1, 1), \mathbb{Z}) \rightarrow H_{\mathrm{cb}}^2(\mathrm{PU}(1, 1), \mathbb{R})$. Then \widehat{G} is the universal covering $\mathrm{PU}(1, 1)$ and $f_{\widehat{G}}$ is the Poincaré translation quasimorphism.

4. A GENERAL FRAMEWORK FOR INJECTIVITY AND DISCRETENESS

In this section we give two results on weakly maximal representations: one concerns injectivity with general locally compact targets, and the other discreteness of the image in the case of Hermitian targets. In both cases we use the characterization of weakly maximal representations in § 3. However, while the characterization holds also if the Toledo invariant vanishes, both results in this section are obtained under the assumption that $T_\kappa(\rho) \neq 0$. Discreteness requires in addition that $T_\kappa(\rho)$ is rational. The rationality question will be further addressed in Section 6.

PROPOSITION 4.1. *Let $\rho : \pi_1(\Sigma) \rightarrow G$ be κ -weakly maximal and assume that $T_\kappa(\rho) \neq 0$. Then ρ is injective.*

Proof. Applying Proposition 3.2, we have that

$$f_\kappa \circ \tilde{\rho} = \lambda(\tau \circ \tilde{\rho}_h) + \psi$$

with $\lambda = \frac{T_\kappa(\rho)}{|\chi(\Sigma)|} \neq 0$. Since $p_\Sigma^{-1}(\ker \rho) = \tilde{\rho}^{-1}(i(\mathbb{Z}))$, we deduce from $\lambda \neq 0$ that $\tau \circ \tilde{\rho}_h$ is a homomorphism on $p_\Sigma^{-1}(\ker \rho)$. A formal argument using the central extension (3.2) restricted to $\ker \rho < \pi_1(\Sigma)$, the commutativity of the diagram

$$\begin{array}{ccc} \widehat{\Gamma} & \xrightarrow{\tilde{\rho}_h} & \widetilde{\text{PU}(1,1)} \\ p_\Sigma \downarrow & & \downarrow p \\ \pi_1(\Sigma) & \xrightarrow{\rho_h} & \text{PU}(1,1), \end{array}$$

and (3.4) implies that

$$(\rho_h|_{\ker \rho})^*(\kappa_{\text{PU}(1,1)}^b) = 0.$$

Since this implies that $\rho_h(\ker \rho)$ is elementary (see next lemma), we conclude that $\ker \rho$ is trivial. \square

We provide a proof of the following easy lemma for ease of reference.

LEMMA 4.2. *Let $\pi : \Delta \rightarrow \text{PU}(1,1)$ be a homomorphism such that $\pi^*(\kappa_{\text{PU}(1,1)}^b) = 0$. Then $\pi(\Delta)$ is elementary, that is $\pi(\Delta)$ has a finite orbit in $\overline{\mathbb{D}}$.*

Proof. Since $\pi^*(\kappa_{\text{PU}(1,1)}^b) = 0$, we conclude from the exactness of the sequence

$$\text{Hom}(\Delta, \mathbb{R}/\mathbb{Z}) \xrightarrow{\delta} H_b^2(\Delta, \mathbb{Z}) \longrightarrow H_b^2(\Delta, \mathbb{R})$$

that $\pi^*(e^b) = \delta(\chi)$ for some homomorphism $\chi : \Delta \rightarrow \mathbb{R}/\mathbb{Z}$. In other words, if we consider χ as a homomorphism into the group of rotations in $\mathrm{PU}(1, 1)$, we have that $\chi^*(e^b) = \pi^*(e^b)$. In particular $(\pi|_{[\Delta, \Delta]})^*(e^b) = 0$. Hence, by Ghys' theorem [24, Theorem 6.6], $[\Delta, \Delta]$ has a fixed point in $\partial\mathbb{D}$. There are then two cases, depending on whether or not $[\Delta, \Delta]$ has infinitely many fixed points. In the first case $[\pi(\Delta), \pi(\Delta)]$ is trivial, that is $\pi(\Delta)$ is abelian. In the second case $\pi(\Delta)$ has a finite orbit in $\partial\mathbb{D}$. In either cases $\pi(\Delta)$ is elementary. \square

The following proposition gives a criterion to insure the discreteness of the image of a κ -weakly maximal representation. In Theorem 7.2 we are going to show that such criterion is always satisfied if κ is the bounded Kähler class.

PROPOSITION 4.3. *Let G be a Lie group of Hermitian type and let $\kappa \in H_{\mathrm{cb}}^2(G, \mathbb{R})$ be a rational class. Assume that $\rho : \pi_1(\Sigma) \rightarrow G$ is κ -weakly maximal and that $T_\kappa(\rho) \in \mathbb{Q}^\times$. Then ρ has discrete image.*

REMARK 4.4. In the case in which $\partial\Sigma = \emptyset$, we have automatically that $T_\kappa(\rho) \in \mathbb{Q}$, so that in this case the condition just reads $T_\kappa(\rho) \neq 0$.

Proof. Let $\Gamma := \pi_1(\Sigma)$. It will be enough to show that $\rho([\Gamma, \Gamma])$ is discrete. In fact $\rho([\Gamma, \Gamma])$ is a normal subgroup of $\rho(\Gamma)$ and, if discrete, also of $\overline{\rho(\Gamma)}$; thus $\rho([\Gamma, \Gamma])$ centralizes the connected component of the identity $\overline{\rho(\Gamma)}^\circ$ of $\overline{\rho(\Gamma)}$. If $\overline{\rho(\Gamma)}^\circ$ were not trivial, we could find $\rho(\gamma) \in \rho(\Gamma) \cap \overline{\rho(\Gamma)}^\circ$ with $\rho(\gamma) \neq e$: indeed $\overline{\rho(\Gamma)}^\circ$ is an open subgroup of $\rho(\Gamma)$ and $\rho(\Gamma)$ is dense in $\overline{\rho(\Gamma)}$. But since, by Proposition 4.1, ρ is injective, this would imply that γ centralizes $[\Gamma, \Gamma]$. This is a contradiction, which shows that $\rho(\Gamma)$ is discrete.

To show the discreteness of $\rho([\Gamma, \Gamma])$, we retain the notation of § 3.3 and we define $L := \overline{\rho([\Gamma, \Gamma])}$ and $L' := p^{-1}(L)$. Observe that (3.6) and (3.9) imply that $p^{-1}(\rho([\Gamma, \Gamma])) = \tilde{\rho}(\widehat{[\Gamma, \Gamma]})\ker p$. Since p is a covering, then $p^{-1}(\overline{\rho([\Gamma, \Gamma])}) = \overline{p^{-1}(\rho([\Gamma, \Gamma]))}$, so that

$$(4.1) \quad L' = \overline{\tilde{\rho}(\widehat{[\Gamma, \Gamma]})\ker p}.$$

We apply now the implication (1) \Rightarrow (3) in Proposition 3.2 to obtain that

$$f_{n\kappa} \circ \tilde{\rho} = \lambda(\tau \circ \tilde{\rho}_h) + \psi,$$

where $\lambda = \frac{T_\kappa(\rho)}{|\chi(\Sigma)|}$ and $\psi \in \mathrm{Hom}(\widehat{[\Gamma, \Gamma]}, \mathbb{R})$. Since $\tau \circ \tilde{\rho}$ is \mathbb{Z} -valued, we deduce that

$$f_{n\kappa}(\tilde{\rho}(\widehat{[\Gamma, \Gamma]})) = \lambda\tau(\tilde{\rho}_h(\widehat{[\Gamma, \Gamma]})) \subset \lambda\tau(\tilde{\rho}_h(\widehat{[\Gamma, \Gamma]})) \subset \lambda\mathbb{Z},$$

which, together with (3.8) implies that

$$f_{n\kappa}(\tilde{\rho}(\widehat{\Gamma}, \widehat{\Gamma}))\ker p \subset \frac{1}{n}\mathbb{Z} + \lambda\mathbb{Z}.$$

Since $\lambda \in \mathbb{Q}$, then $\frac{1}{n}\mathbb{Z} + \lambda\mathbb{Z}$ is discrete in \mathbb{R} . Taking into account (4.1), this implies that $f_{n\kappa}(L') \subset \mathbb{R}$ is a discrete subset. Hence, if we denote by $(L')^\circ$ the identity component of L' , then

$$(4.2) \quad f_{n\kappa}|_{(L')^\circ} = 0.$$

This implies easily that $\kappa|_{L^\circ}$ vanishes as a bounded real class: in fact, because of (4.2) and (3.7),

$$0 = df_{n\kappa}|_{(L')^\circ} = p^*(\kappa)|_{(L')^\circ} = (p|_{(L')^\circ})^*(\kappa|_{L^\circ}),$$

where we used that, since $p : G_{n\kappa} \rightarrow G$ is an open map, $p((L')^\circ) = L^\circ$.

Consider now the subgroup $\Delta := \rho^{-1}(\rho([\Gamma, \Gamma]) \cap L^\circ)$ of Γ . Then $(\rho|_\Delta)^*(\kappa|_{L^\circ}) = 0$ and hence, by hypothesis and definition of κ -weak maximality, $\lambda\kappa_\Sigma^b|_\Delta = 0$. Since by hypothesis $\lambda \neq 0$, then it must be that $\kappa_\Sigma^b|_\Delta = 0$. Thus $\rho_h|_\Delta : \Delta \rightarrow \text{PU}(1, 1)$ satisfies the hypotheses of Lemma 4.2 and hence is elementary. Since $\Delta \triangleleft \pi_1(\Sigma)$, this implies Δ is trivial and hence $\rho([\Gamma, \Gamma]) \cap L^\circ$ is trivial as well.

But since L is a Lie group (as a Lie subgroup of G), L° is open in L and $\rho([\Gamma, \Gamma]) \cap L^\circ$ is dense in L° . Hence L° is trivial, and thus L , and consequently $\rho([\Gamma, \Gamma])$, is discrete. \square

5. ON THE RADICAL DEFINED BY A BOUNDED CLASS

In this section, given a locally compact group L and a bounded rational class $\kappa \in \mathbb{H}_{\text{cb}}^2(L, \mathbb{R})$, we show the existence of a largest normal closed subgroup $\text{Rad}_\kappa(L)$ on which the restriction of the class vanishes. We show moreover that the class κ comes from a bounded real class on the quotient $L/\text{Rad}_\kappa(L)$, the radical of which is trivial. If L is a connected Lie group, the quotient $L/\text{Rad}_\kappa(L)$ is adjoint semisimple without compact factors.

PROPOSITION 5.1. *Let L be a locally compact second countable group and let $\kappa \in \mathbb{H}_{\text{cb}}^2(L, \mathbb{R})$ be a rational class. There is a unique largest normal subgroup $N \triangleleft L$ with $\kappa|_N = 0$ which, in addition, is closed.*

This relies on the following:

LEMMA 5.2. *Let $f : L \rightarrow \mathbb{R}$ be a continuous homogeneous quasimorphism.*

- (1) *There is a unique largest normal subgroup $N_1 \triangleleft L$ with $f|_{N_1} = 0$.*

- (2) *There is a unique largest normal subgroup $N_2 \triangleleft L$ on which f is a homomorphism.*

Both N_1 and N_2 are closed.

Proof. Clearly (2) implies (1) with $N_1 = \ker(f|_{N_2})$.

Let now M_1, M_2 be normal subgroups of L such that $f|_{M_i} : M_i \rightarrow \mathbb{R}$ is a homomorphism. For $m_1 \in M_1$ and $m_2 \in M_2$, let $\chi(m_1 m_2) := f(m_1) + f(m_2)$. We claim that χ is well defined. Indeed, if $m_1 m_2 = m'_1 m'_2$ with $m_i, m'_i \in M_i$, then $(m'_1)^{-1} m_1 = m'_2 m_2^{-1} \in M_1 \cap M_2$. Thus $f((m'_1)^{-1} m_1) = f(m'_2 m_2^{-1})$, which implies, taking into account that f is a homomorphism on M_1 and M_2 , that

$$-f(m'_1) + f(m_1) = f(m'_2) - f(m_2).$$

This shows that χ is well defined. Next we claim that χ is a homomorphism. If $m_i, m'_i \in M_i$, and since M_1 is normal in L , we have

$$\begin{aligned} \chi((m_1 m_2)(m'_1 m'_2)) &= \chi(m_1(m_2 m'_1 m_2^{-1})m_2 m'_2) \\ &= f(m_1(m_2 m'_1 m_2^{-1})) + f(m_2 m'_2) \\ &= f(m_1) + f(m_2 m'_1 m_2^{-1}) + f(m_2) + f(m'_2). \end{aligned}$$

Since f is a homogeneous quasimorphism, we have $f(m_2 m'_1 m_2^{-1}) = f(m'_1)$, which implies that $\chi : M_1 M_2 \rightarrow \mathbb{R}$ is a homomorphism. Since f is a quasimorphism, we have in particular that for all $m_i \in M_i$

$$|f(m_1 m_2) - \chi(m_1 m_2)| = |f(m_1 m_2) - f(m_1) - f(m_2)| \leq C,$$

for some constant C . Thus the homogeneous quasimorphism $f|_{M_1 M_2}$ is at finite distance from the homomorphism χ and hence $f|_{M_1 M_2} = \chi$.

This shows the existence of a unique largest normal subgroup $N_2 \triangleleft L$ on which f is a homomorphism. By continuity of f , the subgroup N_2 is closed. \square

Proof of Proposition 5.1. Let

$$0 \longrightarrow \mathbb{Z} \xrightarrow{i} L_{n\kappa} \xrightarrow{p} L \longrightarrow e$$

be the topological central extension in (3.6) and $f_{n\kappa} : L_{n\kappa} \rightarrow \mathbb{R}$ the continuous homogeneous quasimorphism such that $[df_{n\kappa}] = p^*(\kappa) \in H_{\text{cb}}^2(L_{n\kappa}, \mathbb{R})$. It is an easy verification that, given a subgroup $N < L$, the property $\kappa|_N = 0$ is equivalent to the property that $f_{n\kappa}|_{p^{-1}(N)}$ is a homomorphism. Together with Lemma 5.2, this concludes the proof of the proposition. \square

DEFINITION 5.3. Let $\kappa \in H_{\text{cb}}^2(L, \mathbb{R})$ be a rational class. Denote by $\text{Rad}_\kappa(L)$ the normal closed subgroup of L given by Proposition 5.1 and call it the κ -radical of L .

COROLLARY 5.4. *Let L be a locally compact second countable group and let $\kappa \in H_{\text{cb}}^2(L, \mathbb{R})$ be a rational class.*

- (1) $\text{Rad}_{\kappa}(L) \supset \text{Rad}_a(L)$, where $\text{Rad}_a(L)$ denotes the amenable radical of L .
- (2) *Let*

$$(5.1) \quad \pi : L \rightarrow L/\text{Rad}_{\kappa}(L)$$

denote the canonical projection. Then there is a unique class $u \in H_{\text{cb}}^2(L/\text{Rad}_{\kappa}(L), \mathbb{R})$ with $\pi^(u) = \kappa$. The restriction of u to any non-trivial closed normal subgroup of $L/\text{Rad}_{\kappa}(L)$ does not vanish.*

Proof. (1) follows from the fact that the bounded cohomology of an amenable group vanishes.

(2) The first assertion follows from the exactness of the short sequence

$$0 \longrightarrow H_{\text{cb}}^2(L/N, \mathbb{R}) \longrightarrow H_{\text{cb}}^2(L, \mathbb{R}) \longrightarrow H_{\text{cb}}^2(N, \mathbb{R}),$$

where $N \triangleleft L$ is any closed normal subgroup [16, Theorem 4.1.1]. The second assertion follows from the first and the maximality of $\text{Rad}_{\kappa}(L)$. \square

We denote by $\hat{H}_{\text{cb}}^2(L, \mathbb{Z})$ the cohomology of the complex of integer valued bounded Borel cochains on L . If $\varkappa \in \hat{H}_{\text{cb}}^2(L, \mathbb{Z})$, we denote by $\varkappa^{\mathbb{R}}$ the image of \varkappa in $H_{\text{cb}}^2(L, \mathbb{R})$.

The above discussion applied to a general connected Lie group L has the following nice consequences.

COROLLARY 5.5. *Let L be a connected Lie group and $\varkappa \in \hat{H}_{\text{cb}}^2(L, \mathbb{Z})$. Then $H := L/\text{Rad}_{\varkappa^{\mathbb{R}}}(L)$ is connected adjoint of Hermitian type and a direct product $H = H_1 \times \cdots \times H_n$ of simple non-compact factors.*

Proof. Since the quotient $L/\text{Rad}_a(L)$ of L by its amenable radical is adjoint semisimple without compact factors, so is H (Corollary 5.4(1)). Let $H = H_1 \times \cdots \times H_n$ be the direct product decomposition into simple factors. Let $u \in H_{\text{cb}}^2(H, \mathbb{R})$ be such that $\pi^*(u) = \varkappa^{\mathbb{R}}$, where $\pi : L \rightarrow H$ is the projection in (5.1). According to Corollary 5.4(2), $u|_{H_j} \neq 0$ and hence $H_{\text{cb}}^2(H_j, \mathbb{R}) \neq 0$: thus H_j is of Hermitian type for every $1 \leq j \leq n$ and hence so is H . \square

The following definition will be needed in the next section.

DEFINITION 5.6. Let L be a connected Lie group which admits a closed Levi factor and $\varkappa \in \hat{H}_{\text{cb}}^2(L, \mathbb{Z})$. A \varkappa -Levi factor of L is a connected

semisimple subgroup S with finite center such that $\pi(S) = L/\text{Rad}_\kappa(L)$, where π is as in (5.1), and $\ker(\pi|_S)$ is the center $Z(S)$ of S .

If L_0 is a Levi factor of L , a \varkappa -Levi factor of L is nothing but the product of the almost simple factors in L whose image via π is non-trivial.

6. RATIONALITY QUESTIONS

The class u in Corollary 5.4(2) is *a priori* only a real class. We show in this section that if L is a closed subgroup of a group G of Hermitian type and κ is the restriction of a bounded rational class on G , then u has nice integrality properties.

If in particular $\kappa = \kappa_G^b|_L$ is the restriction of the bounded Kähler class, then the class u is the linear combination of the bounded Kähler classes of the individual simple factors of $L/\text{Rad}_{\kappa_G^b}(L)$ with rational coefficients whose denominators are bounded by an integer depending only on G .

If G is a Lie group of Hermitian type, $L < G$ a closed subgroup and $\varkappa \in \hat{H}_{\text{cb}}^2(G, \mathbb{Z})$, we denote by $\varkappa_L^{\mathbb{R}}$ the restriction of $\varkappa^{\mathbb{R}}$ to L .

PROPOSITION 6.1. *Let G be a Lie group of Hermitian type and let $L < G$ be a closed connected subgroup that admits a closed Levi factor S . Let $\varkappa \in \hat{H}_{\text{cb}}^2(G, \mathbb{Z})$ and let $\pi : L \rightarrow H := L/\text{Rad}_{\varkappa_L^{\mathbb{R}}}(L)$ denote the canonical projection. Let $p_j : H \rightarrow H_j$ be the projection onto the simple factors of H (see Corollary 5.5) and let \varkappa_{H_j} be a generator of $\hat{H}_{\text{cb}}^2(H_j, \mathbb{Z})$, for $j = 1, \dots, n$. If $u \in H_{\text{cb}}^2(H, \mathbb{R})$ is such that $\pi^*(u) = \varkappa_L^{\mathbb{R}}$ (see Corollary 5.4(2)), then*

$$(6.1) \quad u = \sum_{j=1}^n \lambda_j p_j^*(\varkappa_{H_j}^{\mathbb{R}})$$

with

$$(6.2) \quad \lambda_j \in \frac{1}{|Z(S)|} \mathbb{Z},$$

where $Z(S)$ denotes the center of the \varkappa -Levi factor of L .

Proof. According to [15, Proposition 7.7 (3)], the set $\{p_j^*(\varkappa_{H_j}) : 1 \leq j \leq n\}$ is a basis of $\hat{H}_{\text{cb}}^2(H, \mathbb{Z})$ corresponding to the basis $\{p_j^*(\varkappa_{H_j}^{\mathbb{R}}) : 1 \leq j \leq n\}$ of $H_{\text{cb}}^2(H, \mathbb{R})$. It follows that

$$u = \sum_{j=1}^n \lambda_j p_j^*(\varkappa_{H_j}^{\mathbb{R}}),$$

with $\lambda_j \in \mathbb{R}$.

In order to show that the λ_j are rational with an universal bound on the denominator, we consider the following diagram

$$(6.3) \quad \begin{array}{ccccc} & & \xrightarrow{\pi^*} & & \\ & & \text{---} & & \\ \mathrm{H}_{\mathrm{cb}}^2(H, \mathbb{R}) & \xrightarrow{(\pi|_S)_{\mathbb{R}}^*} & \mathrm{H}_{\mathrm{cb}}^2(S, \mathbb{R}) & \xleftarrow{\mathrm{Res}} & \mathrm{H}_{\mathrm{cb}}^2(L, \mathbb{R}) \\ \uparrow & & \uparrow & & \uparrow \\ \hat{\mathrm{H}}_{\mathrm{cb}}^2(H, \mathbb{Z}) & \xrightarrow{(\pi|_S)_{\mathbb{Z}}^*} & \hat{\mathrm{H}}_{\mathrm{cb}}^2(S, \mathbb{Z}) & \xleftarrow{\mathrm{Res}} & \hat{\mathrm{H}}_{\mathrm{cb}}^2(L, \mathbb{Z}), \end{array}$$

where Res is the restriction map in cohomology and the vertical arrows are the change of coefficients.

If $(\pi|_S)_{\mathbb{Z}}^*$ were surjective, again [15, Proposition 7.7 (3)], the commutativity of (6.3) and the fact that $(\pi|_S)_{\mathbb{R}}^*$ is an isomorphism would readily imply the integrality of the λ_j . This is however not necessarily true and the following lemma identifies explicitly the nature of the map $(\pi|_S)_{\mathbb{Z}}^*$.

LEMMA 6.2. *Let $\omega : S \rightarrow H$ be a surjective homomorphism between connected semisimple Lie groups with finite center. Then the map*

$$\omega_{\mathbb{Z}}^* : \hat{\mathrm{H}}_{\mathrm{cb}}^2(H, \mathbb{Z}) \longrightarrow \hat{\mathrm{H}}_{\mathrm{cb}}^2(S, \mathbb{Z})$$

is injective and

$$\mathrm{Image}(\omega_{\mathbb{Z}}^*) \supset |\ker \omega| \hat{\mathrm{H}}_{\mathrm{cb}}^2(S, \mathbb{Z}),$$

where $|\ker \omega|$ denotes the cardinality of $\ker \omega$ and $\hat{\mathrm{H}}_{\mathrm{cb}}^2(S, \mathbb{Z})$ is considered as a \mathbb{Z} -module.

We postpone the proof of the lemma and use its conclusion with $\omega = \pi|_S$. If $Z(S) = \ker(\pi|_S)$ denotes the center of S , the same argument as above, applied to $|Z(S)| \mathcal{X}_S^{\mathbb{R}} \in (\pi|_S)_{\mathbb{Z}}^*(\hat{\mathrm{H}}_{\mathrm{cb}}^2(H, \mathbb{Z}))$ shows that $|Z(S)| u$ is, in fact, in $\hat{\mathrm{H}}_{\mathrm{cb}}^2(H, \mathbb{Z})$ and hence its coordinates $|Z(S)| \lambda_j$ are integers, $j = 1, \dots, n$. \square

Proof of Lemma 6.2. If M is any connected semisimple Lie group with finite center and maximal compact K_M , Wigner's theorem asserts that the restriction map

$$\hat{\mathrm{H}}_{\mathrm{c}}^2(M, \mathbb{Z}) \longrightarrow \hat{\mathrm{H}}_{\mathrm{c}}^2(K_M, \mathbb{Z})$$

is an isomorphism [45]. This, together with the fact that the comparison map

$$\hat{\mathrm{H}}_{\mathrm{cb}}^2(M, \mathbb{Z}) \longrightarrow \hat{\mathrm{H}}_{\mathrm{c}}^2(M, \mathbb{Z})$$

is an isomorphism [15, Proposition 7.7], implies the isomorphism

$$j : \hat{H}_{\text{cb}}^2(M, \mathbb{Z}) \longrightarrow \hat{H}_c^2(K_M, \mathbb{Z}).$$

Using now the long exact sequence in cohomology associated to the short exact sequence

$$0 \longrightarrow \mathbb{Z} \longrightarrow \mathbb{R} \longrightarrow \mathbb{R}/\mathbb{Z} \longrightarrow 0,$$

and the fact that the terms of positive degree and real coefficients vanish, we obtain that the connecting homomorphism

$$\text{Hom}_c(K_M, \mathbb{R}/\mathbb{Z}) \xrightarrow{\delta_{K_M}} \hat{H}_c^2(K_M, \mathbb{Z})$$

is an isomorphism.

Given $\omega : S \rightarrow H$ as in the statement of the lemma, let $K_H < H$ be a maximal compact subgroup. Then $K_S := \omega^{-1}(K_H)$ is a maximal compact in S . One verifies then that the diagram

$$\begin{array}{ccc} \hat{H}_{\text{cb}}^2(H, \mathbb{Z}) & \xrightarrow{\omega_{\mathbb{Z}}^*} & \hat{H}_{\text{cb}}^2(S, \mathbb{Z}) \\ \delta_{K_H}^{-1} \circ j \downarrow & & \downarrow \delta_{K_S}^{-1} \circ j \\ \text{Hom}_c(K_H, \mathbb{R}/\mathbb{Z}) & \xrightarrow{\omega^*} & \text{Hom}_c(K_S, \mathbb{R}/\mathbb{Z}), \end{array}$$

where the vertical arrows are isomorphisms and the bottom one is the precomposition with $\omega|_{K_S}$, is commutative. Then the assertion of the lemma follows from the fact that $\ker \omega \subset K_S$ and that $\omega|_{K_S} : K_S \rightarrow K_H$ is surjective. \square

Recall that the bounded Kähler class $\kappa_G^{\text{b}} \in \text{H}_{\text{cb}}^2(G, \mathbb{R})$ of a Lie group G of Hermitian type is rational [7, § 5].

DEFINITION 6.3. Let G be a Lie group of Hermitian type and $\kappa_G^{\text{b}} \in \text{H}_{\text{cb}}^2(G, \mathbb{R})$ its bounded Kähler class. The *Kähler radical* $\text{Rad}_{\kappa_G^{\text{b}}}(L)$ of a closed subgroup $L < G$ is the $\kappa_G^{\text{b}}|_L$ -radical of L .

If L admits a closed Levi factor, a *Kähler-Levi factor* of L is a $\kappa_G^{\text{b}}|_L$ -Levi factor.

COROLLARY 6.4. *Let G be a Lie group of Hermitian type and let $L < G$ be a closed connected subgroup that admits a closed Levi factor. Then the group $H := L/\text{Rad}_{\kappa_G^{\text{b}}}(L)$ is connected adjoint of Hermitian type, and admits a direct product decomposition $H = H_1 \times \cdots \times H_n$ of simple non-compact factors.*

Moreover if $\pi : L \rightarrow H$ and $p_j : H \rightarrow H_j$ are the canonical projections, then there exists an integer $\ell_G \geq 1$ depending only on G such

that

$$\kappa_G^b|_L = \pi^* \left(\sum_{j=1}^n \nu_j \rho_j^*(\kappa_{H_j}^b) \right),$$

where $\nu_j \in \frac{1}{\ell_G} \mathbb{Z}$.

REMARK 6.5. Let H be a Lie group of Hermitian type. We denote by q_H the smallest integer such that there exists $\varkappa \in H_{\text{cb}}^2(H, \mathbb{Z})$ with $q_H \kappa_H^b = \varkappa^{\mathbb{R}}$. If in addition H is simple and \varkappa_H is a generator of $H_{\text{cb}}^2(H, \mathbb{Z})$, then there exists an integer m_H such that $\varkappa = m_H \varkappa_H$. It follows that

$$(6.4) \quad \varkappa_H^{\mathbb{R}} = \frac{q_H}{m_H} \kappa_H^b.$$

Proof of Corollary 6.4. We apply Proposition 6.1 to $\varkappa \in H_{\text{cb}}^2(G, \mathbb{Z})$ such that $\varkappa^{\mathbb{R}} = q_G \kappa_G^b$. Then (6.1) and (6.2), together with (6.4), show that

$$\kappa_G^b|_L = \pi^* \left(\sum_{j=1}^n \nu_j \rho_j^*(\kappa_{H_j}^b) \right),$$

with

$$\nu_j \in \frac{q_{H_j}}{q_G m_{H_j} |Z(S)|} \mathbb{Z},$$

where $|Z(S)|$ is the cardinality of the center of a Kähler-Levi factor S .

Since there are only finitely many possible conjugacy classes of connected semisimple subgroups of G , we obtain the result. \square

7. STRUCTURE OF WEAKLY MAXIMAL REPRESENTATIONS

In § 4 the discreteness of a κ -weakly maximal representation was proven under the assumption that the Toledo invariant $T_\kappa(\rho)$ is rational. In this section we prove that if κ is the bounded Kähler class, this is always the case and that the representation into the quotient of its Zariski closure by the Kähler radical is also discrete and injective. The definition and properties of the Kähler radical will be essential to show that the Toledo invariant of the representation into the quotient is also non-zero.

An interesting feature of the proof of the rationality of the Toledo invariant is that it depends upon showing first that the single factors of the quotient by the Kähler radical are of tube type.

Let G be a Lie group of Hermitian type and $\kappa_G^b \in H_{\text{cb}}^2(G, \mathbb{R})$ its bounded Kähler class. For ease of notation, we denote the Toledo invariant $T_{\kappa_G^b}$ by

$$T : \text{Hom}(\pi_1(\Sigma), G) \rightarrow \mathbb{R}.$$

DEFINITION 7.1. A homomorphism $\rho : \pi_1(\Sigma) \rightarrow G$ is weakly maximal if

$$T(\rho) = \|\rho^*(\kappa_G^b)\| |\chi(\Sigma)|,$$

that is if ρ is κ_G^b -weakly maximal in the sense of Definition 2.1.

THEOREM 7.2. *Let $G = \mathbf{G}(\mathbb{R})^\circ$ be of Hermitian type, where \mathbf{G} is a connected semisimple algebraic group defined over \mathbb{R} , and let $\rho : \pi_1(\Sigma) \rightarrow G$ be a weakly maximal homomorphism. Let L be the connected component of the real points of the Zariski closure of the image of ρ and let $\Gamma := \rho^{-1}(L)$. Assume that $T(\rho) \neq 0$. Then:*

- (1) *the group $L/\text{Rad}_{\kappa_G^b}(L)$ is Hermitian of tube type;*
- (2) *there is an integer $\ell_G \geq 1$ depending only on G (see Corollary 6.4), such that*

$$T(\rho) \in \frac{|\chi(\Sigma)|}{\ell_G} \mathbb{Z};$$

- (3) *the composition*

$$\Gamma \xrightarrow{\rho|_\Gamma} L \xrightarrow{\pi} L/\text{Rad}_{\kappa_G^b}(L)$$

is injective with discrete image.

Using that in a surface group there are no finite normal subgroups, one obtains immediately:

COROLLARY 7.3. *Let $\rho : \pi_1(\Sigma) \rightarrow G$ be a weakly maximal homomorphism with $T(\rho) \neq 0$. Then ρ is injective with discrete image.*

An important role in the proof is played by the generalized Maslov cocycle. Recall that if H is a connected Lie group of Hermitian type and \check{S} the Shilov boundary of the associated bounded symmetric domain, the *generalized Maslov cocycle* $\beta_{\check{S}} : \check{S}^3 \rightarrow \mathbb{R}$ is a bounded alternating H -invariant cocycle constructed by J.-L. Clerc in [19]. We will use it in two ways: on the one hand it represents the bounded Kähler class in a particular resolution useful to implement pullbacks in bounded cohomology (see (7.2)); on the other, through the relation (7.4) with the Hermitian triple product

$$\langle\langle \cdot, \cdot, \cdot \rangle\rangle : \check{S}^{(3)} \rightarrow \mathbb{R}^\times \setminus \mathbb{C}^\times,$$

it gives a criterion to detect when H is of tube type, [7, 13]

We refer the reader to [6, 7] and [13, § 4.2] for details.

Proof of Theorem 7.2. We use heavily in this proof techniques developed in [16, 6, 7, 13], to which we refer the reader for details.

According to Corollary 6.4, the group $H := L/\text{Rad}_{\kappa_G^b}(L) = H_1 \times \cdots \times H_n$ is of Hermitian type and $\kappa_G^b|_L = \pi^*(u)$, where

$$u = \sum_{j=1}^n \nu_j p_j^*(k_{H_j}^b),$$

and $p_j : H \rightarrow H_j$ and $\pi : L \rightarrow H$ are the canonical projections. We set aside for the moment that the ν_j are rational, and will pick it up towards the end of the proof.

Let $\xi := \pi \circ \rho|_\Gamma : \Gamma \rightarrow L \rightarrow H$ be the composition of π with $\rho|_\Gamma$. Observe that Γ is of finite index in $\pi_1(\Sigma)$. It follows that, since ρ is weakly maximal, that is $\rho^*(\kappa_G^b) = \lambda \kappa_\Sigma^b$ for $\lambda = \frac{\text{T}(\rho)}{|\chi(\Sigma)|}$, then

$$(7.1) \quad \xi^*(u) = \lambda \kappa_{\Sigma'}^b,$$

where $\Sigma' \rightarrow \Sigma$ is the finite covering corresponding to $\Gamma < \pi_1(\Sigma)$.

As usual, to realize the pullback ξ^* in bounded cohomology, we use boundary maps. According to [6], this is possible since if $(\mathcal{B}_{\text{alt}}^\infty(\check{S}^\bullet))$ denotes the complex of bounded alternating Borel cocycles on \check{S} , then the class $[\beta_{\check{S}}]$ defined by the generalized Maslov cocycle corresponds to the bounded Kähler class κ_H^b under the canonical map

$$(7.2) \quad \text{H}^\bullet(\mathcal{B}_{\text{alt}}^\infty(\check{S}^\bullet)^H) \longrightarrow \text{H}_{\text{cb}}^\bullet(H, \mathbb{R}).$$

Likewise, if $\check{S} = \check{S}_1 \times \cdots \times \check{S}_n$ is the decomposition into a product, where \check{S}_j is the Shilov boundary of H_j and $\check{p}_j : \check{S} \rightarrow \check{S}_j$ is the projection, a standard cohomological argument shows that the diagram

$$\begin{array}{ccc} \text{H}_{\text{cb}}^\bullet(H_j, \mathbb{R}) & \xrightarrow{p_j^*} & \text{H}_{\text{cb}}^\bullet(H, \mathbb{R}) \\ \uparrow & & \uparrow \\ \text{H}^\bullet(\mathcal{B}_{\text{alt}}^\infty(\check{S}_j^\bullet)^{H_j}) & \xrightarrow{\check{p}_j^*} & \text{H}_{\text{cb}}^\bullet(\mathcal{B}_{\text{alt}}^\infty(\check{S}^\bullet)^H) \end{array}$$

commutes. It follows that u is represented, again via (7.2), by the bounded Borel cocycle on \check{S}^3 defined by

$$(x, y, z) \longmapsto \sum_{j=1}^n \nu_j \beta_{\check{S}_j}(x_j, y_j, z_j).$$

To recall the existence of the boundary map, endow the interior of Σ' with a complete hyperbolic metric of finite area so that Γ is identified with a lattice in $\text{PU}(1, 1)$. Since $\xi(\Gamma)$ is Zariski dense in H , there is a

Γ -equivariant measurable map $\varphi : \partial\mathbb{D} \rightarrow \check{S}$. Because of the product structure of \check{S} , the map φ has the form

$$\varphi_1 \times \cdots \times \varphi_n : \partial\mathbb{D} \longrightarrow \check{S}_1 \times \cdots \times \check{S}_n,$$

where $\varphi_j : \partial\mathbb{D} \rightarrow \check{S}_j$ is measurable and equivariant with respect to $p_j \circ \xi$.

Consequently, according to [6] the pullback $\xi^*(u)$ is represented by the following bounded measurable Γ -invariant cocycle

$$(x, y, z) \longmapsto \sum_{j=1}^n \nu_j \beta_{\check{S}_j}(\varphi_j(x), \varphi_j(y), \varphi_j(z)),$$

for almost all $(x, y, z) \in (\partial\mathbb{D})^3$.

Using now again that the bounded fundamental class of Σ is represented by $\beta_{\partial\mathbb{D}}$, it follows from (7.1) that

$$(7.3) \quad \lambda \beta_{\partial\mathbb{D}}(x, y, z) = \sum_{j=1}^n \nu_j \beta_{\check{S}_j}(\varphi_j(x), \varphi_j(y), \varphi_j(z))$$

for almost all $(x, y, z) \in (\partial\mathbb{D})^3$.

Recall now that the Hermitian triple product is an H -invariant real algebraic map on $(\check{S})^3$ whose relation with the Maslov cocycle is given by

$$(7.4) \quad \langle\langle x, y, z \rangle\rangle_{\check{S}} \equiv e^{i\pi d_{\check{S}} \beta_{\check{S}}(x, y, z)} \pmod{\mathbb{R}^\times},$$

where $d_{\check{S}}$ is an integer given in terms of the root system of H , [13]. We will use this relation for the Hermitian triple product on each of the H_j . To uniformize the exponents, since $\nu_j \in \mathbb{Q}^\times$, we can pick an integer $m \geq 1$ such that $m\nu_j = n_j d_j$ for some $n_j \in \mathbb{Z}$, $1 \leq j \leq n$. Multiplying (7.3) by m , exponentiating and using that $\beta_{\partial\mathbb{D}}$ takes only $\pm 1/2$ as values, we obtain

$$(7.5) \quad e^{i\pi m \lambda (\pm 1/2)} = \prod_{j=1}^n \langle\langle \varphi_j(x), \varphi_j(y), \varphi_j(z) \rangle\rangle_{\check{S}_j}^{n_j},$$

for almost every $(x, y, z) \in (\partial\mathbb{D})^3$. If $x_j, y_j, z_j \in \check{S}$, we define now

$$R((x_j), (y_j), (z_j)) := \prod_{j=1}^n \langle\langle x_j, y_j, z_j \rangle\rangle_{\check{S}_j}^{n_j}.$$

Because of (7.5) the function R takes at most two values on $\varphi(\partial\mathbb{D})^3$; since the latter set is Zariski dense in \check{S}^3 and R is rational, it takes on at most two values on \check{S}^3 . The factors $\langle\langle x_j, y_j, z_j \rangle\rangle_{\check{S}_j}^{n_j}$ are pairwise independent rational functions, therefore each of these factors itself

can take only finitely many values. As previously recalled, it follows from [13] that the corresponding groups are of tube type, thus showing assertion (1).

Furthermore, since $\beta_{\mathfrak{S}_j}$ takes only values in $\frac{1}{2}\mathbb{Z}$ [20, Theorem 4.3], it follows from (7.3) and Corollary 6.4 that $\lambda \in \frac{1}{\ell_G}\mathbb{Z}$. This shows assertion (2).

The third assertion follows from Proposition 4.1 (2) and Proposition 4.3. \square

8. WEAKLY MAXIMAL REPRESENTATIONS AND RELATIONS WITH OTHER REPRESENTATION VARIETIES

In this section we first show that the set of weakly maximal representations and the set of weakly maximal representations with non-zero Toledo number are closed. Then we examine the relationship of weakly maximal representations with Shilov-Anosov representations.

Returning to our general framework in § 3, let G be locally compact and second countable, Γ a discrete group and $\kappa \in H_{\text{cb}}^2(G, \mathbb{R})$ a fixed class. We define now a (Hausdorff) topology on $H_{\text{b}}^2(\Gamma, \mathbb{R})$ with respect to which the map

$$(8.1) \quad \begin{aligned} \text{Hom}(\Gamma, G) &\rightarrow H_{\text{b}}^2(\Gamma, \mathbb{R}) \\ \rho &\longmapsto \rho^*(\kappa) \end{aligned}$$

will be continuous. To this purpose recall that $H_{\text{b}}^2(\Gamma, \mathbb{R}) := \ker \delta^2 / \text{im } \delta^1$, where

$$0 \longrightarrow \mathbb{R} \longrightarrow \ell^\infty(\Gamma) \xrightarrow{\delta^1} \ell^\infty(\Gamma^2) \xrightarrow{\delta^2} \ell^\infty(\Gamma^3) \xrightarrow{\delta^3} \dots$$

denotes the inhomogeneous bar resolution. If we endow each $\ell^\infty(\Gamma^n)$ with the weak-* topology as the dual of $\ell^1(\Gamma^n)$, then:

LEMMA 8.1. *$H_{\text{b}}^2(\Gamma, \mathbb{R})$ is a Hausdorff topological vector space with the quotient weak-* topology.*

Proof. It is clear that $\ker \delta^2$ is weak-* closed. Since $H_{\text{b}}^2(\Gamma, \mathbb{R})$ is a Banach space, $\text{im } \delta^1$ is closed and hence, by [41, Theorem 4.14], $\text{im } \delta^1$ is weak-* closed as well. \square

LEMMA 8.2. *The function in (8.1) is continuous with respect to the weak-* topology on $H_{\text{b}}^2(\Gamma, \mathbb{R})$.*

Proof. Let $c_\kappa : G^2 \rightarrow \mathbb{R}$ be a bounded continuous inhomogeneous cocycle representing κ . Let $(\rho_n)_{n \geq 1}$ be a sequence of elements in $\text{Hom}(\Gamma, G)$

converging to ρ . Setting

$$\begin{aligned} c_n(x, y) &:= c_\kappa(\rho_n(x), \rho_n(y)) \\ c(x, y) &:= c_\kappa(\rho(x), \rho(y)), \end{aligned}$$

we have that $c_n \in \ell^\infty(\Gamma^2)$ (respectively $c \in \ell^\infty(\Gamma^2)$) represent $\rho_n^*(\kappa)$ (respectively $\rho^*(\kappa)$). In addition

- (1) $c_n \rightarrow c$ pointwise on Γ^2 , and
- (2) $\|c_n\|_\infty$ and $\|c\|_\infty$ are bounded by $\|c_\kappa\|_\infty$.

Then if $f \in \ell^1(\Gamma^2)$, we have that

- (1) $f c_n \rightarrow f c$ pointwise, and
- (2) $|f(x, y) c_n(x, y)| \leq |f(x, y)| \|c_\kappa\|_\infty$,

then the Dominated Convergence Theorem implies that

$$\int_{\Gamma^2} f c_n \rightarrow \int_{\Gamma^2} f c$$

and shows therefore that $\rho_n^*(\kappa) \rightarrow \rho^*(\kappa)$ for the quotient topology in $H_b^2(\Gamma, \mathbb{R})$. \square

If now $\Gamma = \pi_1(\Sigma)$, then

COROLLARY 8.3. *The set $\text{Hom}_{wm}(\pi_1(\Sigma), G)$ of weakly maximal representations is closed in $\text{Hom}(\pi_1(\Sigma), G)$.*

Proof. It follows from Proposition 3.2 that

$$\text{Hom}_{wm}(\pi_1(\Sigma), G) = \{\rho \in \text{Hom}(\pi_1(\Sigma), G) : \rho^*(\kappa) = t\kappa_\Sigma^b \text{ for some } t \in [0, \infty)\}.$$

Since $H_{cb}^2(\pi_1(\Sigma), \mathbb{R})$ with the weak-* topology is Hausdorff, the subset $\{t\kappa_\Sigma^b : t \in [0, \infty)\}$ is closed. The assertion follows then from the continuity of the map $\rho \mapsto \rho^*(\kappa)$. \square

Let now set

$$\text{Hom}_{wm}^*(\pi_1(\Sigma), G) := \{\rho : \pi_1(\Sigma) \rightarrow G : \rho \text{ is weakly maximal and } T(\rho) \neq 0\}.$$

COROLLARY 8.4. *Let G be a Lie group of Hermitian type. Then the set of weakly maximal representations with non-zero Toledo invariant is closed in $\text{Hom}(\pi_1(\Sigma), G)$.*

Proof. By Theorem 7.2(2) we have

$$\text{Hom}_{wm}^*(\pi_1(\Sigma), G) = \left\{ \rho \in \text{Hom}_{wm}(\pi_1(\Sigma), G) : T(\rho) \geq \frac{\|\chi(\Sigma)\|}{l_G} \right\},$$

which implies the claim since $\rho \mapsto T(\rho)$ is continuous, [15, Proposition 3.10], and $\text{Hom}_{wm}(\pi_1(\Sigma), G)$ is closed. \square

Finally, we turn to the relation with Anosov representations. Let $\partial\Sigma = \emptyset$ and realize $\pi_1(\Sigma)$ as cocompact lattice Γ in $\mathrm{PU}(1, 1)$ via a hyperbolization. If G is, say, simple of tube type with Shilov boundary \check{S} , then a Shilov-Anosov representation $\rho : \Gamma \rightarrow G$ implies the existence of a (unique) continuous equivariant map $\varphi : \partial\mathbb{D} \rightarrow \check{S}$ with the additional property that for every $x, y \in \partial\mathbb{D}$ with $x \neq y$, $\varphi(x)$ and $\varphi(y)$ are transverse. Let $(\partial\mathbb{D})^{3,+}$ be the connected set of distinct, positively oriented triples in $(\partial\mathbb{D})^3$ and $\check{S}^{(3)}$ the set of triples of pairwise transverse points in \check{S}^3 . Then

$$\varphi \times \varphi \times \varphi : (\partial\mathbb{D})^{3,+} \rightarrow \check{S}^{(3)}$$

must send $(\partial\mathbb{D})^{3,+}$ into a connected component of $\check{S}^{(3)}$. Thus if $\beta_{\check{S}} : \check{S}^3 \rightarrow \mathbb{R}$ denotes the generalized Maslov cocycle, $\beta_{\check{S}} \circ \varphi^3$ is a multiple of the orientation cocycle. This and Proposition 3.2 imply that the set

$$\mathrm{Hom}_{\check{S}\text{-An}}^+(\pi_1(\Sigma), G)^\circ := \{\rho : \pi_1(\Sigma) \rightarrow G : \rho \text{ is Shilov-Anosov with } T(\rho) \geq 0\}$$

is contained in $\mathrm{Hom}_{wm}(\pi_1(\Sigma), G)$. Taking into account that the latter is closed, we get

COROLLARY 8.5.

$$(8.2) \quad \overline{\mathrm{Hom}_{\check{S}\text{-An}}^+(\pi_1(\Sigma), G)} \subset \mathrm{Hom}_{wm}(\pi_1(\Sigma), G).$$

9. EXAMPLES

In this section we describe some examples of weakly maximal representations.

9.1. Maximal Representations. Any maximal representation is a weakly maximal representation. Concrete examples of maximal representations are described in [15, 10, 32].

9.2. Embeddings of $\mathrm{SL}(2, \mathbb{R})$. Special examples of weakly maximal representations arise from embeddings of $\mathrm{SL}(2, \mathbb{R})$. Consider a faithful representation $\rho_0 : \pi_1(\Sigma) \rightarrow \mathrm{SL}(2, \mathbb{R})$ with discrete image, and let $\tau : \mathrm{SL}(2, \mathbb{R}) \rightarrow G$ be an injective homomorphism into a Lie group of Hermitian type G . Then the composition $\tau \circ \rho_0 : \pi_1(\Sigma) \rightarrow G$ is a weakly maximal representation.

More generally if H, G are Lie groups of Hermitian type, and $\tau : H \rightarrow G$ is a homomorphism such that $\tau^*(\kappa_G^b)$ is a multiple of κ_H^b . Then the composition of a weakly maximal representation $\rho : \pi_1(\Sigma) \rightarrow H$ with τ is a weakly maximal representation into G .

9.3. Shilov-Anosov Representations. We explained in § 8 that any Shilov-Anosov representation into a Hermitian Lie group G of tube type with non-negative Toledo number is weakly maximal. Here we just give an example of such a representation into $\mathrm{Sp}(2n, \mathbb{R})$. Let again $\rho_i : \pi_1(\Sigma) \rightarrow \mathrm{SL}(2, \mathbb{R})$ $i = 1, \dots, n$ be faithful representations with discrete image. Consider the embedding $\tau : \mathrm{SL}(2, \mathbb{R})^n \rightarrow \mathrm{Sp}(2n, \mathbb{R})$ corresponding to a maximal polydisk in the bounded symmetric domain realization of $\mathrm{Sp}(2n, \mathbb{R})$. Then $\rho := \tau \circ (\rho_1, \dots, \rho_n) : \pi_1(\Sigma) \rightarrow \mathrm{Sp}(2n, \mathbb{R})$ is a Shilov-Anosov representation, and hence a weakly maximal representations. Since the set of Shilov-Anosov representations is open, any small deformation of such a representation is also weakly maximal. Similarly to the construction in [10] one can explicitly construct bending deformations of the representation ρ with Zariski dense image.

Note that ρ is maximal if and only if the representations ρ_i are orientation preserving for all $i = 1, \dots, n$.

9.4. Cancelling Contributions. Let $G = G_1 \times \dots \times G_n$ be a semisimple Lie group of Hermitian type. If $\rho : \pi_1(\Sigma) \rightarrow G$ is a maximal representation, then $\rho_i = p_i \circ \rho : \pi_1(\Sigma) \rightarrow G_i$, $i = 1, \dots, n$ are maximal representations, where p_i denotes the projection onto the i -th factor.

In order to illustrate that this does not hold true for weakly maximal representations consider an arbitrary representation $\rho_a : \pi_1(\Sigma) \rightarrow \mathrm{SL}(2, \mathbb{R})$ and denote by $\bar{\rho}_a : \pi_1(\Sigma) \rightarrow \mathrm{SL}(2, \mathbb{R})$ the composition of ρ_a with the outer automorphism of $\mathrm{SL}(2, \mathbb{R})$ reversing the orientation. Let $\rho_0 : \pi_1(\Sigma) \rightarrow \mathrm{SL}(2, \mathbb{R})$ be a faithful representation with discrete image, as before. Then the representation $\rho = (\rho_0, \rho_a, \bar{\rho}_a) : \pi_1(\Sigma) \rightarrow \mathrm{SL}(2, \mathbb{R}) \times \mathrm{SL}(2, \mathbb{R}) \times \mathrm{SL}(2, \mathbb{R})$ is weakly maximal.

9.5. Limit of Shilov-Anosov Representations. Let $\partial\Sigma = \emptyset$. We have seen in Corollary 8.5 that any representation that is the limit of Shilov-Anosov representations is weakly maximal. By choosing appropriately ρ_a in § 9.4 it is easy to construct representations that are weakly maximal but not Shilov-Anosov. While we do not know whether the containment in (8.2) is strict, we give here an example to show that the containment

$$\mathrm{Hom}_{\mathcal{S}\text{-An}}^*(\pi_1(\Sigma), G) \subset \overline{\mathrm{Hom}_{\mathcal{S}\text{-An}}^*(\pi_1(\Sigma), G)}$$

is strict, that is an example of a representation that is the limit of Shilov-Anosov representations but is not Shilov-Anosov.

To this purpose we consider the group $\mathrm{Sp}(2n, \mathbb{R})$ and we denote for simplicity by $\mathcal{R}_d(n)$ the set of representations $\rho : \pi_1(\Sigma) \rightarrow \mathrm{Sp}(2n, \mathbb{R})$ with Toledo invariant equal to d and with $(\mathcal{R}_d(n))_{\mathcal{S}\text{-An}} \subset \mathcal{R}_d(n)$ the subset of Shilov-Anosov representations.

If $d = (g - 1)n$, that is if $\mathcal{R}_{(g-1)n}(n)$ consists of maximal representations, then by [10] there is the equality $(\mathcal{R}_{(g-1)n}(n))_{\check{S}-An} \equiv \mathcal{R}_{(g-1)n}(n)$. On the other hand, it is easy to construct a representation that has zero Toledo invariant but is not Shilov-Anosov (by taking for example $\rho_a \oplus \overline{\rho_a} : \pi_1(\Sigma) \rightarrow \mathrm{SL}(2, \mathbb{R}) \times \mathrm{SL}(2, \mathbb{R}) \hookrightarrow \mathrm{Sp}(4, \mathbb{R})$, where ρ_a is any representation with dense image in $\mathrm{SL}(2, \mathbb{R})$). Since for $n \geq 3$ the set $\mathcal{R}_0(n)$ is connected [23, Theorem 1.1 (1)] and $(\mathcal{R}_0(n))_{\check{S}-An}$ is open, this shows that

$$(\mathcal{R}_0(n))_{\check{S}-An} \subsetneq \overline{(\mathcal{R}_0(n))_{\check{S}-An}}.$$

We can use this fact as follows. Recall that the direct sum of symplectic vector spaces leads to a canonical homomorphism

$$\mathrm{Sp}(2n_1, \mathbb{R}) \times \mathrm{Sp}(2n_2, \mathbb{R}) \longrightarrow \mathrm{Sp}(2(n_1 + n_2), \mathbb{R})$$

and consequently to a continuous map

$$\begin{aligned} \mathcal{R}_{d_1}(n_1) \times \mathcal{R}_{d_2}(n_2) &\rightarrow \mathcal{R}_{d_1+d_2}(n_1 + n_2) \\ (\rho_1, \rho_2) &\mapsto \rho_1 \oplus \rho_2. \end{aligned}$$

Observe moreover that $\rho_1 \oplus \rho_2$ is Shilov-Anosov precisely if both ρ_1 and ρ_2 are. In particular if $n_1 \geq 3$ we can consider a sequence $\rho_1^{(k)} \in (\mathcal{R}_0(n_1))_{\check{S}-An}$ whose limit ρ_1 is not Shilov-Anosov. Then for any $\rho_2 \in (\mathcal{R}_d(n_2))_{\check{S}-An}$, the sequence $\rho_1^{(k)} \oplus \rho_2$ converges to the representation $\rho_1 \oplus \rho_2$ that is not Shilov-Anosov and has Toledo invariant equal to d .

9.6. Weakly maximal representations with non-reductive Zariski

closure. Let J_n be the $2n \times 2n$ matrix with block entries $J_1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

We consider the subgroup of $\mathrm{Sp}(2(n+1), \mathbb{R})$ given by

$$Q = \left\{ \begin{pmatrix} A & b & 0 \\ 0 & 1 & 0 \\ c & d & 1 \end{pmatrix} : A \in \mathrm{Sp}(2n, \mathbb{R}) \text{ and } c = {}^t b J_n A \right\},$$

which is the semidirect product of $\mathrm{Sp}(2n, \mathbb{R})$ and the n -dimensional Heisenberg group H_n .

A map $\rho : \Gamma \rightarrow Q$ with entries $\pi(\gamma), b(\gamma), c(\gamma)$ and $d(\gamma)$ is a homomorphism if and only if $\pi : \Gamma \rightarrow \mathrm{Sp}(2n, \mathbb{R})$ is a homomorphism, $b : \Gamma \rightarrow \mathbb{R}^{2n}$ is a 1-cocycle (with the Γ -module structure on \mathbb{R}^{2n} is given by π) and

$$(9.1) \quad d(\gamma_1 \gamma_2) = {}^t b(\gamma_1) J_n \pi(\gamma_1) b(\gamma_2) + d(\gamma_1) + d(\gamma_2).$$

To reinterpret (9.1), we observe that there is a bilinear symmetric form on $H^1(\Gamma, \pi)$ with values in $H^2(\Gamma, \mathbb{R})$ obtained by composing the cup product $H^1(\Gamma, \pi) \times H^1(\Gamma, \pi) \rightarrow H^2(\Gamma, \pi \otimes \pi)$ with the projection on

trivial coefficients given by the invariant symplectic form; denoting by Q_π the corresponding quadratic form, we have that

$$(\gamma_1, \gamma_2) \longmapsto {}^t b(\gamma_1) J_n \pi(\gamma_1) b(\gamma_2)$$

is a representative of $Q_\pi([b])$ and the existence of a function d satisfying (9.1) amounts to $Q_\pi([b]) = 0$.

Let now $\Gamma = \pi_1(S)$, where S is an oriented surface of genus g , and let $\rho_i : \Gamma \rightarrow \mathrm{Sp}(2n_i, \mathbb{R})$, where ρ_1 is in the Hitchin component of $\mathrm{Sp}(2n_1, \mathbb{R})$, while ρ_2 is the precomposition of a Hitchin representation into $\mathrm{Sp}(2n_2, \mathbb{R})$ with an orientation reversing automorphism of Γ . Identifying $H^2(\Gamma, \mathbb{R})$ with \mathbb{R} by means of the chosen orientation, we have [25] that Q_{ρ_1} is positive definite while Q_{ρ_2} is negative definite. Choose $[b_i] \in H^1(\Gamma, \rho_i)$ both non-zero such that

$$(9.2) \quad Q_{\rho_1}([b_1]) + Q_{\rho_2}([b_2]) = 0.$$

Set $\pi := \rho_1 \oplus \rho_2$, $b := b_1 + b_2$ and let d be a solution of (9.1), which exists by (9.2). Then $\rho : \pi_1(S) \rightarrow \mathrm{Sp}(2(n+1), \mathbb{R})$ is a weakly maximal representation with $T(\rho) = n_1 - n_2$, and the real Zariski closure of its image is the semi direct product of $G_1 \times G_2 < \mathrm{Sp}(2n, \mathbb{R})$ with H_n ; here G_i is the real Zariski closure of the image of ρ_i .

REFERENCES

1. G. Ben Simon, M. Burger, T. Hartnick, A. Iozzi and A. Wienhard, *Order preserving and causal representations*.
2. S. B. Bradlow, O. García-Prada, and P. B. Gothen, *Surface group representations and $U(p, q)$ -Higgs bundles*, J. Differential Geom. **64** (2003), no. 1, 111–170. MR 2015045 (2004k:53142)
3. ———, *Maximal surface group representations in isometry groups of classical Hermitian symmetric spaces*, Geom. Dedicata **122** (2006), 185–213. MR 2295550 (2008e:14013)
4. ———, *Deformations of maximal representations in $\mathrm{Sp}(4, \mathbb{R})$* , Q. J. Math. **63** (2012), 795–843, <http://arxiv.org/abs/0903.5496>.
5. M. Bucher, M. Burger, R. Frigerio, A. Iozzi, C. Pagliantini, and B. Pozzetti, *Isometric properties of relative bounded cohomology*, <http://arxiv.org/abs/1205.1022>.
6. M. Burger and A. Iozzi, *Boundary maps in bounded cohomology. Appendix to: “Continuous bounded cohomology and applications to rigidity theory” [Geom. Funct. Anal. **12** (2002), no. 2, 219–280; MR1911660 (2003d:53065a)] by Burger and N. Monod*, Geom. Funct. Anal. **12** (2002), no. 2, 281–292. MR 1911668 (2003d:53065b)
7. M. Burger and A. Iozzi, *Bounded Kähler class rigidity of actions on Hermitian symmetric spaces*, Ann. Sci. École Norm. Sup. (4) **37** (2004), no. 1, 77–103. MR 2050206 (2005b:32048)

8. M. Burger and A. Iozzi, *Bounded differential forms, generalized Milnor-Wood inequality and an application to deformation rigidity*, *Geom. Dedicata* **125** (2007), 1–23. MR 2322535 (2009c:53053)
9. M. Burger and A. Iozzi, *Bounded cohomology and totally real subspaces in complex hyperbolic geometry*, *Ergodic Theory Dynam. Systems* **32** (2012), no. 2, 467–478. MR 2901355
10. M. Burger, A. Iozzi, F. Labourie, and A. Wienhard, *Maximal representations of surface groups: symplectic Anosov structures*, *Pure Appl. Math. Q.* **1** (2005), no. 3, Special Issue: In memory of Armand Borel. Part 2, 543–590. MR 2201327 (2007d:53064)
11. M. Burger, A. Iozzi, and A. Wienhard, *Higher Teichmüller Spaces: from $SL(2, R)$ to other Lie groups*, To appear in Handbook of Teichmüller theory.
12. ———, *Maximal representations and Anosov structures*, In preparation.
13. M. Burger, A. Iozzi, and A. Wienhard, *Hermitian symmetric spaces and Kähler rigidity*, *Transform. Groups* **12** (2007), no. 1, 5–32. MR 2308026 (2009c:32044)
14. M. Burger, A. Iozzi, and A. Wienhard, *Tight homomorphisms and Hermitian symmetric spaces*, *Geom. Funct. Anal.* **19** (2009), no. 3, 678–721. MR 2563767 (2011d:53080)
15. ———, *Surface group representations with maximal Toledo invariant*, *Ann. of Math. (2)* **172** (2010), no. 1, 517–566. MR 2680425 (2012j:22014)
16. M. Burger and N. Monod, *Continuous bounded cohomology and applications to rigidity theory*, *Geom. Funct. Anal.* **12** (2002), no. 2, 219–280. MR 1911660 (2003d:53065a)
17. D. Calegari, *Faces of the scl norm ball*, *Geom. Topol.* **13** (2009), no. 3, 1313–1336. MR 2496047 (2010e:20062)
18. S. Choi and W. M. Goldman, *Convex real projective structures on closed surfaces are closed*, *Proc. Amer. Math. Soc.* **118** (1993), no. 2, 657–661. MR 1145415 (93g:57017)
19. J.-L. Clerc, *L'indice de Maslov généralisé*, *J. Math. Pures Appl. (9)* **83** (2004), no. 1, 99–114. MR 2032583 (2004k:53127)
20. J.-L. Clerc and B. Orsted, *The Maslov index revisited*, *Transform. Groups* **6** (2001), no. 4, 303–320. MR 1870049 (2002j:53103)
21. V. V. Fock and A. B. Goncharov, *Moduli spaces of local systems and higher Teichmüller theory*, *Publ. Math. Inst. Hautes Études Sci.* (2006), no. 103, 1–211. MR 2233852 (2009k:32011)
22. ———, *Moduli spaces of convex projective structures on surfaces*, *Adv. Math.* **208** (2007), no. 1, 249–273.
23. O. García-Prada, P. B. Gothen, and I. Mundet i Riera, *Higgs bundles and surface group representations in real symplectic groups*, <http://arxiv.org/abs/0809.0576>, to appear in *Journal of Topology*.
24. É. Ghys, *Groups acting on the circle*, *Enseign. Math. (2)* **47** (2001), no. 3–4, 329–407. MR 1876932 (2003a:37032)
25. W. M. Goldman, personal communication.
26. ———, *Discontinuous groups and the Euler class*, Thesis, University of California at Berkeley, 1980.
27. ———, *Characteristic classes and representations of discrete subgroups of Lie groups*, *Bull. Amer. Math. Soc. (N.S.)* **6** (1982), no. 1, 91–94. MR 634439 (83b:22012)

28. P. B. Gothen, *Components of spaces of representations and stable triples*, *Topology* **40** (2001), no. 4, 823–850. MR 1851565 (2002k:14017)
29. M. Gromov, *Volume and bounded cohomology*, *Inst. Hautes Études Sci. Publ. Math.* (1982), no. 56, 5–99 (1983). MR 686042 (84h:53053)
30. O. Guichard, *Composantes de Hitchin et représentations hyperconvexes de groupes de surface*, *J. Differential Geom.* **80** (2008), no. 3, 391–431. MR 2472478 (2009h:57031)
31. O. Guichard and A. Wienhard, *Convex foliated projective structures and the Hitchin component for $\mathrm{PSL}_4(\mathbf{R})$* , *Duke Math. J.* **144** (2008), no. 3, 381–445. MR 2444302 (2009k:53223)
32. ———, *Topological invariants of Anosov representations*, *J. Topol.* **3** (2010), no. 3, 578–642. MR 2684514 (2011f:57027)
33. ———, *Anosov representations: domains of discontinuity and applications*, *Invent. Math.* **190** (2012), no. 2, 357–438. MR 2981818
34. O. Hamlet, *Tight holomorphic maps, a classification*, <http://arxiv.org/abs/1110.5443>, to appear.
35. ———, *Tight maps, a classification*, <http://arxiv.org/abs/1206.4465>.
36. T. Hartnick and T. Strubel, *Cross ratios, translation lengths and maximal representations*, *Geom. Dedicata* **161** (2012), 285–322. MR 2994044
37. L. Hernández Lamonedá, *Maximal representations of surface groups in bounded symmetric domains*, *Trans. Amer. Math. Soc.* **324** (1991), no. 1, 405–420. MR 1033234 (91f:32040)
38. N. J. Hitchin, *Lie groups and Teichmüller space*, *Topology* **31** (1992), no. 3, 449–473. MR 1174252 (93e:32023)
39. F. Labourie, *Anosov flows, surface groups and curves in projective space*, *Invent. Math.* **165** (2006), no. 1, 51–114. MR 2221137 (2007c:20101)
40. G. W. Mackey, *Les ensembles boréliens et les extensions des groupes*, *J. Math. Pures Appl.* (9) **36** (1957), 171–178. MR 0089998 (19,752a)
41. W. Rudin, *Functional analysis*, second ed., *International Series in Pure and Applied Mathematics*, McGraw-Hill Inc., New York, 1991. MR 1157815 (92k:46001)
42. D. Toledo, *Representations of surface groups in complex hyperbolic space*, *J. Differential Geom.* **29** (1989), no. 1, 125–133. MR 978081 (90a:57016)
43. A. Wienhard, *Bounded cohomology and geometry*, Ph.D. thesis, *Bonner Mathematische Schriften Nr. 368*, Bonn, 2004.
44. ———, *The action of the mapping class group on maximal representations*, *Geom. Dedicata* **120** (2006), 179–191. MR 2252900 (2008g:20112)
45. D. Wigner, *Algebraic cohomology of topological groups*, *Trans. Amer. Math. Soc.* **178** (1973), 83–93. MR 0338132 (49 #2898)

E-mail address: gabi.ben-simon@math.ethz.ch

DEPARTMENT MATHEMATIK, ETH ZÜRICH, RÄMISTRASSE 101, CH-8092 ZÜRICH, SWITZERLAND

E-mail address: burger@math.ethz.ch

DEPARTMENT MATHEMATIK, ETH ZÜRICH, RÄMISTRASSE 101, CH-8092 ZÜRICH,
SWITZERLAND

E-mail address: `hartnick@tx.technion.ac.il`

MATHEMATICS DEPARTMENT, TECHNION - ISRAEL INSTITUTE OF TECHNOLOGY,
HAIFA, 32000, ISRAEL

E-mail address: `iozzi@math.ethz.ch`

DEPARTMENT MATHEMATIK, ETH ZÜRICH, RÄMISTRASSE 101, CH-8092 ZÜRICH,
SWITZERLAND

E-mail address: `wienhard@mathi.uni-heidelberg.de`

MATHEMATISCHES INSTITUT, RUPRECHT-KARLS-UNIVERSITÄT HEIDELBERG,
69120 HEIDELBERG, GERMANY
CALIFORNIA INSTITUTE OF TECHNOLOGY, 1200 E CALIFORNIA BLVD, PASADENA,
CA 91125