

# DS-GA 1014: Homework Problem Set 5

Optimization and Computational Linear Algebra for Data Science (Fall 2018)

Afonso S. Bandeira  
bandeira@cims.nyu.edu  
<http://www.cims.nyu.edu/~bandeira>

Brett D. Bernstein  
brettb@  
~brettb

Due on Friday October 12, 2018

**This homework problem set is due on October 12 on NYU Classes.**

If you have questions about the homework feel free to contact Brett Bernstein (brett.bernstein@nyu.edu) or myself, or stop by our office hours.

Unless otherwise stated all answer must be mathematically justified, even questions that ask for a computation.

Try not to look up the answers, you'll learn much more if you try to think about the problems without looking up the solutions. If you need hints, feel free to ask on Piazza.

You can work in groups but each student must write his/her own solution based on his/her own understanding of the problem. Please list, on your submission, the students you work with for the homework (this will not affect your grade).

Late submissions will be graded with a penalty of 10% per day late. Weekend days do not count, from Friday to Monday counts only 1 day.

If you need to impose extra conditions on a problem to make it easier (or consider specific cases of the question, like taking  $n$  to be 2, e.g.), state explicitly that you have done so. Solutions where extra conditions were assume, or where only special cases were treated, will also be graded (probably scored as a partial answer).

Problems with a (\*) are extra credit, they will not (directly) contribute to your score of this homework. However, for every 4 (four) extra credit questions successfully answered you get a homework "bye": your lowest homework score (or one you did not hand in) gets replaced by a perfect score.

**Problem 5.1** Given  $x \in \mathbb{R}^n$ , and  $p \geq 1$  we define

$$\|x\|_p = \left( \sum_{k=1}^n |x_k|^p \right)^{1/p}.$$

Notice that  $\|x\|_2$  is the norm we defined before. Show that

$$\frac{1}{\sqrt{n}} \|x\|_1 \leq \|x\|_2 \leq \|x\|_1.$$

**Problem 5.2** Let  $u_1, \dots, u_n \in \mathbb{R}^n$  be orthonormal, and suppose

$$v = \alpha_1 u_1 + \dots + \alpha_n u_n$$

for some  $\alpha \in \mathbb{R}^n$ . Compute  $\|v\|$  and justify your calculation.

**Problem 5.3** Given a subspace  $U \subseteq \mathbb{R}^n$  we define the orthogonal complement  $U^\perp$  of  $U$  by

$$U^\perp := \{v \in \mathbb{R}^n : \langle u, v \rangle = 0 \text{ for all } u \in U \}.$$

1. Show that  $U^\perp$  is a subspace of  $\mathbb{R}^n$ .
2. Prove that any vector  $u \in \mathbb{R}^n$  can be written as  $u = x + y$  where  $x \in U$  and  $y \in U^\perp$ .

**Problem 5.4** For any  $x \in \mathbb{R}^n$  compute  $\min\{x^T y : y \in \mathbb{R}^n, \|y\| = 1\}$  and  $\max\{x^T y : y \in \mathbb{R}^n, \|y\| = 1\}$ , and justify your calculations are correct.

(\*) **Problem 5.5 (For Extra Credit)** Let  $A \in \mathbb{R}^{m \times n}$  and define  $p, q \in \mathbb{R}$  by

$$p = \max_i \sum_{j=1}^n |A_{ij}| \quad \text{and} \quad q = \max_j \sum_{i=1}^m |A_{ij}|.$$

Prove that  $x^T A y \leq \|x\| \|y\| \sqrt{pq}$ .