

Recap

KKT (Karush-Kuhn-Tucker) conditions

necessary and sufficient conditions for a solution of the Lasso objective function

$$\begin{aligned} G_j(\hat{\beta}) &= -\text{sign}(\hat{\beta}_j)\lambda \text{ if } \hat{\beta}_j \neq 0 \\ |G_j(\hat{\beta})| &\leq \lambda \text{ if } \hat{\beta}_j = 0 \end{aligned}$$

where

$$G(\beta) = -2X^T(Y - X\beta)/n$$

(for the solution $\hat{\beta}$: sub-differential must contain the zero element)

sparsity is induced at points of non-differentiability (for each components β_j)

Coordinate descent algorithms

for optimization, exploiting the KKT conditions

path following algorithms:

compute $\{\hat{\beta}_j(\lambda)\}_{j=1}^p$ over all values of $\lambda \in \mathbb{R}^+$

the coefficient paths are typically “non-monotone” in the non-zeros

it may happen that

$$\hat{\beta}_j(\lambda) \neq 0, \hat{\beta}_j(\lambda') = 0 \text{ for } \lambda' < \lambda$$

Generalized Linear Models (GLMs)

univariate response Y , covariate $X \in \mathcal{X} \subseteq \mathbb{R}^p$

GLM: $Y_1, \dots, Y_n$ independent

$$g(\mathbb{E}[Y_i|X_i = x]) = \underbrace{\mu + \sum_{j=1}^p \beta_j x^{(j)}}_{=f(x)=f_{\mu,\beta}(x)}$$

μ in the model: one cannot simply center the data

$g(\cdot)$ real-valued, known link function

Lasso: ℓ_1 -norm regularized maximum likelihood estimation

$$\hat{\mu}, \hat{\beta} = \operatorname{argmin}_{\mu, \beta} \left(\underbrace{-\ell(\mu, \beta)}_{\text{neg. log-likelihood}} + \lambda \|\beta\|_1 \right)$$