

Effective bounds for induced size-Ramsey numbers of cycles

Domagoj Bradač

joint work with Nemanja Draganić and Benny Sudakov

(ETH Zürich)

Definition

For a positive integer k , a graph G is k -Ramsey for a graph H if every k -edge-coloring of G contains a monochromatic copy of H . We write $G \xrightarrow{k} H$.

Definition

For a positive integer k , a graph G is k -Ramsey for a graph H if every k -edge-coloring of G contains a monochromatic copy of H . We write $G \xrightarrow{k} H$.

Theorem (Ramsey's theorem)

For every k and H , there exists N such that $K_N \xrightarrow{k} H$.

Definition

For a positive integer k , a graph G is k -Ramsey for a graph H if every k -edge-coloring of G contains a monochromatic copy of H . We write $G \xrightarrow{k} H$.

Theorem (Ramsey's theorem)

For every k and H , there exists N such that $K_N \xrightarrow{k} H$.

Definition

The k -color Ramsey number of H , denoted by $r^k(H)$, is defined as $r^k(H) = \min\{v(G) \mid G \xrightarrow{k} H\}$.

Size-Ramsey numbers

Definition

The k -color size-Ramsey number of H is $\hat{r}^k(H) = \min\{e(G) \mid G \xrightarrow{k} H\}$.

Definition

The k -color size-Ramsey number of H is $\hat{r}^k(H) = \min\{e(G) \mid G \xrightarrow{k} H\}$.

For fixed k , $\hat{r}^k(H) = O(n)$ for an n -vertex graph H which is:

Definition

The k -color size-Ramsey number of H is $\hat{r}^k(H) = \min\{e(G) \mid G \xrightarrow{k} H\}$.

For fixed k , $\hat{r}^k(H) = O(n)$ for an n -vertex graph H which is:

- a path (Beck '83),
- a bounded degree tree (Friedman, Pippenger '87),
- a cycle (Haxell, Kohayakawa, Łuczak '95),
- a bounded degree graph with bounded treewidth (Kamčev, Liebenau, Wood, Yepremyan '21; Berger, Kohayakawa, Maesaka, Martins, Mendonça, Mota, Parczyk '21),
- a logarithmic subdivision of a bounded degree graph (Draganić, Krivelevich, Nenadov '22).

Definition

The k -color size-Ramsey number of H is $\hat{r}^k(H) = \min\{e(G) \mid G \xrightarrow{k} H\}$.

For fixed k , $\hat{r}^k(H) = O(n)$ for an n -vertex graph H which is:

- a path (Beck '83),
- a bounded degree tree (Friedman, Pippenger '87),
- a cycle (Haxell, Kohayakawa, Łuczak '95),
- a bounded degree graph with bounded treewidth (Kamčev, Liebenau, Wood, Yepremyan '21; Berger, Kohayakawa, Maesaka, Martins, Mendonça, Mota, Parczyk '21),
- a logarithmic subdivision of a bounded degree graph (Draganić, Krivelevich, Nenadov '22).

However, $\hat{r}^2(H)$ is not linear in $v(H)$ for all bounded degree graphs (Rödl, Szemerédi '00; Tikhomirov '22+).

Definition

A graph G is k -color induced Ramsey for H if every k -edge-coloring of G contains a monochromatic copy of H as an induced subgraph of G .

Definition

A graph G is k -color induced Ramsey for H if every k -edge-coloring of G contains a monochromatic copy of H as an induced subgraph of G .

Definition

The k -color induced Ramsey number of H is the minimum number of vertices of a graph that is k -color induced Ramsey for H .


Induced Ramsey numbers

Definition

A graph G is k -color induced Ramsey for H if every k -edge-coloring of G contains a monochromatic copy of H as an induced subgraph of G .

Definition

The k -color induced Ramsey number of H is the minimum number of vertices of a graph that is k -color induced Ramsey for H .



Definition

A graph G is k -color induced Ramsey for H if every k -edge-coloring of G contains a monochromatic copy of H as an induced subgraph of G .

Definition

The k -color induced Ramsey number of H is the minimum number of vertices of a graph that is k -color induced Ramsey for H .

- These numbers exist (Deuber '75; Erdős, Hajnal, Pósa '75; Rödl '73).

Definition

A graph G is k -color induced Ramsey for H if every k -edge-coloring of G contains a monochromatic copy of H as an induced subgraph of G .

Definition

The k -color induced Ramsey number of H is the minimum number of vertices of a graph that is k -color induced Ramsey for H .

- These numbers exist (Deuber '75; Erdős, Hajnal, Pósa '75; Rödl '73).
- $r_{\text{ind}}^2(H) = 2^{O(n \log n)}$ (Conlon, Fox, Sudakov '12).

Definition

A graph G is k -color induced Ramsey for H if every k -edge-coloring of G contains a monochromatic copy of H as an induced subgraph of G .

Definition

The k -color induced Ramsey number of H is the minimum number of vertices of a graph that is k -color induced Ramsey for H .

- These numbers exist (Deuber '75; Erdős, Hajnal, Pósa '75; Rödl '73).
- $r_{\text{ind}}^2(H) = 2^{O(n \log n)}$ (Conlon, Fox, Sudakov '12).
- Erdős conjectured $r_{\text{ind}}^2(H) = 2^{O(n)}$.

Definition

The k -color induced size-Ramsey number, $\hat{r}_{\text{ind}}^k(H)$, of H is the minimum number of edges of a graph that is k -color induced Ramsey for H .

Induced size-Ramsey numbers

Definition

The k -color induced size-Ramsey number, $\hat{r}_{\text{ind}}^k(H)$, of H is the minimum number of edges of a graph that is k -color induced Ramsey for H .

Theorem (Haxell, Kohayakawa, Łuczak '95)

For every k , there is $C = C(k)$ such that $\hat{r}_{\text{ind}}^k(P_n), \hat{r}_{\text{ind}}^k(C_n) \leq Cn$.

Induced size-Ramsey numbers

Definition

The k -color induced size-Ramsey number, $\hat{r}_{\text{ind}}^k(H)$, of H is the minimum number of edges of a graph that is k -color induced Ramsey for H .

Theorem (Haxell, Kohayakawa, Łuczak '95)

For every k , there is $C = C(k)$ such that $\hat{r}_{\text{ind}}^k(P_n), \hat{r}_{\text{ind}}^k(C_n) \leq Cn$.


Question

What is the best value of $C = C(k)$ for cycles in the Theorem above?

Previous results

	Lower bound	Upper bound
$\hat{r}^k(P_n)$	$\Omega(k^2)n$ (DP '17)	$O(k^2 \log k)n$ (K '19)
$\hat{r}_{\text{ind}}^k(P_n)$	$\Omega(k^2)n$ (DP '17)	$O(k^3 \log^4 k)n$ (DGK '22)
$\hat{r}^k(C_n), n$ even	$\Omega(k^2)n$ (DP '17)	$O(k^{120} \log^2 k)n$ (JM '23)
$\hat{r}^k(C_n), n$ odd	$2^{k-1}n$ (JM '23)	$O(2^{k^2+16 \log k})n$ (JM '23)
$\hat{r}_{\text{ind}}^k(C_n), n$ even	$\Omega(k^2)n$ (DP '17)	? (HKŁ '95)
$\hat{r}_{\text{ind}}^k(C_n), n$ odd	$2^{k-1}n$ (JM '23)	? (HKŁ '95)

Previous results

	Lower bound	Upper bound
$\hat{r}^k(P_n)$	$\Omega(k^2)n$ (DP '17)	$O(k^2 \log k)n$ (K '19)
$\hat{r}_{\text{ind}}^k(P_n)$	$\Omega(k^2)n$ (DP '17)	$O(k^3 \log^4 k)n$ (DGK '22)
$\hat{r}^k(C_n), n$ even	$\Omega(k^2)n$ (DP '17)	$O(k^{120} \log^2 k)n$ (JM '23)
$\hat{r}^k(C_n), n$ odd	$2^{k-1}n$ (JM '23)	$O(2^{k^2+16 \log k})n$ (JM '23)
$\hat{r}_{\text{ind}}^k(C_n), n$ even	$\Omega(k^2)n$ (DP '17)	
$\hat{r}_{\text{ind}}^k(C_n), n$ odd	$2^{k-1}n$ (JM '23)	

Theorem (B., Draganić, Sudakov '23+)

For any $k \geq 1$, there is n_0 such that for $n \geq n_0$, the following holds.

- $\hat{r}^k(C_n) = 2^{O(k)}n.$
- *If n is even, then $\hat{r}_{\text{ind}}^k(C_n) = O(k^{102})n.$*
- *If n is odd, then $\hat{r}_{\text{ind}}^k(C_n) = 2^{O(k \log k)}n.$*

Overview of results

	Lower bound	Upper bound
$\hat{r}^k(P_n)$	$\Omega(k^2)n$ (DP '17)	$O(k^2 \log k)n$ (K '19)
$\hat{r}_{\text{ind}}^k(P_n)$	$\Omega(k^2)n$ (DP '17)	$O(k^3 \log^4 k)n$ (DGK '22)
$\hat{r}^k(C_n), n$ even	$\Omega(k^2)n$ (DP '17)	$O(k^{102})n$ (BDS '23+)
$\hat{r}^k(C_n), n$ odd	$2^{k-1}n$ (JM '23)	$2^{O(k)}n$ (BDS '23+)
$\hat{r}_{\text{ind}}^k(C_n), n$ even	$\Omega(k^2)n$ (DP '17)	$O(k^{102})n$ (BDS '23+)
$\hat{r}_{\text{ind}}^k(C_n), n$ odd	$2^{k-1}n$ (JM '23)	$2^{O(k \log k)}n$ (BDS '23+)

Previous results: the host graph is $G(N, C_1/N)$, where $N = C_2n$.

Previous results: the host graph is $G(N, C_1/N)$, where $N = C_2n$. Choose a suitable color and find an induced C_n in this color or a large subgraph without this color.

Previous results: the host graph is $G(N, C_1/N)$, where $N = C_2n$. Choose a suitable color and find an induced C_n in this color or a large subgraph without this color.

Main idea: in this suitable color, it is easier to find a cycle of length in $[0.9n, 1.1n]$ than of length exactly n .

Previous results: the host graph is $G(N, C_1/N)$, where $N = C_2n$. Choose a suitable color and find an induced C_n in this color or a large subgraph without this color.

Main idea: in this suitable color, it is easier to find a cycle of length in $[0.9n, 1.1n]$ than of length exactly n .

Our new host graph construction is designed to exploits this.

Construction to obtain $\hat{r}_{\text{ind}}^k(C_n) = 2^{O(k \log k)} n$:

Construction to obtain $\hat{r}_{\text{ind}}^k(C_n) = 2^{O(k \log k)} n$:

- Find a small *gadget graph* $F = F(k)$ which is k -induced Ramsey for C_5 .

Host graph construction and auxiliary graph

Construction to obtain $\hat{r}_{\text{ind}}^k(C_n) = 2^{O(k \log k)} n$:


- Find a small *gadget graph* $F = F(k)$ which is k -induced Ramsey for C_5 .
- Place $C_1 N$ random copies of F , to get the host graph Γ on $N = C_2 n$ vertices.

Host graph construction and auxiliary graph

Construction to obtain $\hat{r}_{\text{ind}}^k(C_n) = 2^{O(k \log k)} n$:


- Find a small *gadget graph* $F = F(k)$ which is k -induced Ramsey for C_5 .
- Place $C_1 N$ random copies of F , to get the host graph Γ on $N = C_2 n$ vertices.
- Auxiliary graph G with $V(G) = V(\Gamma)$ and edges: for each placed copy of F , find one monochromatic induced C_5 and connect two nonadjacent vertices on this C_5 .

H




Host graph and auxiliary graph

Γ




Host graph and auxiliary graph

Γ




Host graph and auxiliary graph

Γ





Host graph and auxiliary graph

Γ



Host graph and auxiliary graph


 Γ  G 

Using the auxiliary graph

Claim: a “good” monochromatic cycle in G of any length $\ell \in [n/3, n/2]$ gives an induced monochromatic cycle of length n in Γ .


Using the auxiliary graph

Claim: a “good” monochromatic cycle in G of any length $\ell \in [n/3, n/2]$ gives an induced monochromatic cycle of length n in Γ .




Using the auxiliary graph

Claim: a “good” monochromatic cycle in G of any length $\ell \in [n/3, n/2]$ gives an induced monochromatic cycle of length n in Γ .




Using the auxiliary graph

Claim: a “good” monochromatic cycle in G of any length $\ell \in [n/3, n/2]$ gives an induced monochromatic cycle of length n in Γ .




Using the auxiliary graph

Claim: a “good” monochromatic cycle in G of any length $\ell \in [n/3, n/2]$ gives an induced monochromatic cycle of length n in Γ .




Using the auxiliary graph

Claim: a “good” monochromatic cycle in G of any length $\ell \in [n/3, n/2]$ gives an induced monochromatic cycle of length n in Γ .




Using the auxiliary graph

Claim: a “good” monochromatic cycle in G of any length $\ell \in [n/3, n/2]$ gives an induced monochromatic cycle of length n in Γ .



Using the auxiliary graph

Claim: a “good” monochromatic cycle in G of any length $\ell \in [n/3, n/2]$ gives an induced monochromatic cycle of length n in Γ .



Finding an induced cycle of approximate length

Our task essentially reduces to finding an induced cycle of length in $[n/3, n/2]$ in a locally sparse graph.

Finding an induced cycle of approximate length

Our task essentially reduces to finding an induced cycle of length in $[n/3, n/2]$ in a locally sparse graph.

Main tool: modification of the DFS algorithm for induced paths developed by Draganić, Glock and Krivelevich.

Getting all the results

For different results, we use different gadget graphs.

Getting all the results

For different results, we use different gadget graphs.

- For the odd induced case F is k -color induced Ramsey for C_5 . We take F to be Alon's dense pseudorandom triangle free graph on $2^{O(k \log k)}$ vertices.

Getting all the results

For different results, we use different gadget graphs.

- For the odd induced case F is k -color induced Ramsey for C_5 . We take F to be Alon's dense pseudorandom triangle free graph on $2^{O(k \log k)}$ vertices.
- For the even induced case F is k -color induced Ramsey for C_6 . We take F to be a dense C_4 -free bipartite on $O(k^6)$ vertices.

For different results, we use different gadget graphs.

- For the odd induced case F is k -color induced Ramsey for C_5 . We take F to be Alon's dense pseudorandom triangle free graph on $2^{O(k \log k)}$ vertices.
- For the even induced case F is k -color induced Ramsey for C_6 . We take F to be a dense C_4 -free bipartite on $O(k^6)$ vertices.
- For the odd (non-induced) we want every k -edge-coloring of F to have an odd monochromatic cycle. We take $F = K_{2^{k+1}}$.

Conjecture

For odd n , $\hat{r}_{\text{ind}}^k(C_n) = 2^{O(k)}n$.

Concluding remarks

Conjecture

For odd n , $\hat{r}_{\text{ind}}^k(C_n) = 2^{O(k)}n$.

It is enough to prove the following.

Conjecture

There is a graph F on $2^{O(k)}$ vertices such that in every k -edge-coloring of F there is an odd monochromatic cycle of length at least 5 as an induced subgraph.

Conjecture

For odd n , $\hat{r}_{\text{ind}}^k(C_n) = 2^{O(k)}n$.

It is enough to prove the following.

Conjecture

There is a graph F on $2^{O(k)}$ vertices such that in every k -edge-coloring of F there is an odd monochromatic cycle of length at least 5 as an induced subgraph.

Thank you!