

# Multivariate Extremes and the Aggregation of Dependent Risks: Examples and Counter-Examples

Paul Embrechts   Dominik D. Lambrigger   Mario V. Wüthrich  
ETH Zurich, Department of Mathematics, 8092 Zurich, Switzerland

Submitted: January 30, 2008

This version: July 29, 2008

## Abstract

Properties of risk measures for extreme risks have become an important topic of research. In the present paper we discuss sub- and superadditivity of quantile based risk measures and show how multivariate extreme value theory yields the ideal modeling environment. Numerous examples and counter-examples highlight the applicability of the main results obtained.

Keywords: multivariate extreme value theory, multivariate regular variation, risk aggregation, spectral measure, subadditivity, tail dependence, Value-at-Risk.

## 1 Introduction

In Embrechts et al. [11], the following example was worked out. Suppose  $X_1, X_2$  are independent random variables (rvs) each with a Pareto distribution function (df)

$$P(X_i > x) = x^{-1/2}, \quad x \geq 1,$$

i.e., the  $X_i$ 's have infinite mean. Consider  $X = X_1 + X_2$  and let  $\alpha \in (0, 1)$ , then

$$F_{X_1+X_2}^{-1}(\alpha) > F_{X_1}^{-1}(\alpha) + F_{X_2}^{-1}(\alpha), \quad (1.1)$$

i.e., quantiles act superadditively. This rather trivial example has far-reaching consequences in finance, where the  $X_i$ 's correspond to profits or losses, over a given fixed (holding) period, in particular markets/instruments. A quantile of such a fixed period position is referred to as Value-at-Risk (VaR); see also Definition 2.1 below. Hence (1.1) can be rewritten as

$$\text{VaR}_\alpha(X_1 + X_2) > \text{VaR}_\alpha(X_1) + \text{VaR}_\alpha(X_2). \quad (1.2)$$

The above example (1.1) and its numerous generalizations form an important topic of research in Quantitative Risk Management (QRM) as for instance discussed in McNeil et al. [23], Chapter 6. It also has important consequences within (re)insurance when modeling catastrophic risks; see Ibragimov et al. [17].

Understanding the practical relevance of situations where (1.2) holds, or indeed where subadditivity (" $\leq$ " in (1.2)) holds are crucial within the regulatory framework (so-called Basel I and II) of financial institutions; see Chapter 1 in McNeil et al. [23] and the references therein. Indeed, under the Basel II framework, the quantile risk measure  $\text{VaR}_\alpha(X)$  corresponds to regulatory (risk) capital that a financial institution has to hold in order to be able to carry the risky position  $X$  on its books. Furthermore, the quantity

$$D_\alpha(X_1, X_2) = \text{VaR}_\alpha(X_1 + X_2) - \text{VaR}_\alpha(X_1) - \text{VaR}_\alpha(X_2)$$

can be seen as a measure of *diversification*. Alternatively, the quantity

$$C_\alpha(X_1, X_2) = \frac{\text{VaR}_\alpha(X_1 + X_2)}{\text{VaR}_\alpha(X_1) + \text{VaR}_\alpha(X_2)}$$

is referred to as a measure of *concentration* within the Basel II framework. Consequently, a deeper understanding concerning the possible values of either  $D_\alpha(X_1, X_2)$  and  $C_\alpha(X_1, X_2)$  across a wide family of dfs relevant for QRM practice is important.

This paper presents several results on this topic for arbitrary dimensions  $n \geq 2$  and dependence structures, and this within the unifying framework of multivariate extreme value theory (MEVT). The MEVT approach to the above problems is by no means new. We found however that a summary of these results keeping financial applications in mind would be highly useful. Through many concrete examples

and counter-examples we show that care has to be taken concerning possible constraints/properties of the dfs of the underlying risk factors. In a wider context of QRM, these same techniques are becoming increasingly important in the analysis of high risk scenarios, see for instance Balkema and Embrechts [3], and therefore will become part of the standard toolkit of QRM.

The paper is organized as follows. Section 2 recalls the basic notion of multivariate regular variation and its link to questions like (1.1). In Section 3 we discuss three examples where (1.1) may or may not hold, stressing in particular the important difference between one-sided and two-sided risk dfs. For positive rvs, Section 4 uses the notion of spectral measure to derive additivity-type results under general portfolio assumptions. Sections 5 and 6 study the link with tail dependence concepts, whereas Section 7 concludes.

## 2 Value-at-Risk and multivariate regular variation

In this section we introduce multivariate regular variation, which provides a natural framework to discuss diversification of a portfolio under the risk measure VaR. Throughout the paper, we use the language of MEVT. For the latter, several approaches exist, like the more geometric one as presented in Balkema and Embrechts [3] or Barbe [4]. In these contributions, the geometry of the level sets of the underlying multivariate densities plays a crucial role. We choose to base our discussion on the notion of spectral measure and MEVT results in this context. Part IV in Balkema and Embrechts [3] compares some of the different approaches and highlights these in the context of high-threshold exceedances, which is akin to VaR estimation.

**DEFINITION 2.1 (VALUE-AT-RISK)** Let  $X$  be a rv with df  $F$ . The *Value-at-Risk* with respect to the level  $\alpha \in (0, 1)$  is defined as the generalized inverse of  $F$ ,  $\text{VaR}_\alpha(X) = F^{\leftarrow}(\alpha) = \inf \{x \in \mathbb{R} \mid F(x) \geq \alpha\}$ .  $\square$

In all relevant situations,  $\alpha$  is typically close to 1. We say that VaR is *asymptotically subadditive* for  $X_1, \dots, X_n$ , if

$$\lim_{\alpha \nearrow 1} \frac{\text{VaR}_\alpha(\sum_{i=1}^n X_i)}{\sum_{i=1}^n \text{VaR}_\alpha(X_i)} \leq 1, \quad (2.1)$$

provided the limit exists. VaR is called *asymptotically superadditive* for  $X_1, \dots, X_n$  if “ $\geq$ ” in (2.1) holds. We assume the reader to be familiar with univariate EVT and in particular univariate regular variation; see for instance Embrechts et al. [10] for an introduction. The following definition introduces multivariate regular variation and also the *limiting constant*  $q$ , which is of main interest in this paper; standard textbooks on multivariate EVT are for instance Resnick [30], [32], Beirlant et al. [6], de Haan and Ferreira [14] and Balkema and Embrechts [3]. A brief and very readable introduction to the field is found in Mikosch [24].

**DEFINITION 2.2 (MULTIVARIATE REGULAR VARIATION)** A random vector  $\mathbf{X} = (X_1, \dots, X_n)'$  is *multivariate regularly varying* with index  $-\beta < 0$ , if there exists a probability measure  $\mu$ , a measurable function  $b : (0, \infty) \rightarrow (0, \infty)$  with  $\lim_{t \rightarrow \infty} b(t) = \infty$  and a scalar  $q = q(b) > 0$  such that for all  $r > 0$ ,

$$\lim_{t \rightarrow \infty} t P \left( \|\mathbf{X}\| > rb(t), \frac{\mathbf{X}}{\|\mathbf{X}\|} \in G \right) = qr^{-\beta} \mu(G),$$

for any Borel set  $G \subset \mathfrak{N}_{\|\cdot\|}^{n-1} = \{\mathbf{x} = (x_1, \dots, x_n)' \in \mathbb{R}^n \mid \|\mathbf{x}\| = 1\}$ . We write  $\mathbf{X} \in \text{MRV}_n(-\beta)$ .  $\square$

The definition of multivariate regular variation is independent of the explicit choice of the norm  $\|\cdot\|$  on  $\mathbb{R}^n$ . This comes from the fact that all norms on  $\mathbb{R}^n$  are equivalent; see Lemma 2.1 in Hult and Lindskog [16] for details. Note that the limiting constant  $q$  depends on the index  $-\beta < 0$  and on the norm  $\|\cdot\|$  chosen.

The goal of this paper is to analyze the properties of the limiting constant  $q$  for random vectors  $\mathbf{X}$  with identically distributed marginals (this assumption can be relaxed using change of norms techniques; see Section 4) and with a dependence structure within the framework of multivariate regular variation. It follows from Definition 2.2 that for  $\mathbf{X} = (X_1, \dots, X_n)' \in \text{MRV}_n(-\beta)$ ,  $\beta > 0$ ,

$$q(\beta, \|\cdot\|) = \lim_{x \rightarrow \infty} \frac{P(\|\mathbf{X}\| > x)}{P(X_1 > x)} > 0;$$

see Barbe et al. [5], formula (9) and Remark 1 in Resnick [31]. An interesting choice of norm is the  $l_1$ -norm  $\|\cdot\|_1$  on  $\mathbb{R}_+^n$ , to study the sum  $X_1 + \dots + X_n$  of  $n$  risky positions.

However, also more general loss functions, say  $\Psi$ , are considered in practice.

**Lemma 2.3** *Let  $\mathbf{X} = (X_1, \dots, X_n)' \in \text{MRV}_n(-\beta)$ ,  $\beta > 0$ , with identically distributed marginals. If for a measurable function  $\Psi : \mathbb{R}^n \rightarrow \mathbb{R}$ ,*

$$\lim_{x \rightarrow \infty} \frac{P(\Psi(\mathbf{X}) > x)}{P(X_1 > x)} = q_\Psi \in (0, \infty), \quad (2.2)$$

then

$$\lim_{\alpha \nearrow 1} \frac{\text{VaR}_\alpha(\Psi(\mathbf{X}))}{\text{VaR}_\alpha(X_1)} = q_\Psi^{1/\beta}.$$

PROOF: Consider  $F_\Psi(x) = P(\Psi(\mathbf{X}) \leq x)$  and  $F(x) = P(X_1 \leq x)$ . Using (2.2) and the regular variation properties of  $X_1$ , one shows that

$$\lim_{\alpha \nearrow 1} \frac{F_\Psi^\leftarrow(\alpha)}{F^\leftarrow(\alpha)} = \lim_{x \rightarrow \infty} \frac{x}{F^\leftarrow(F_\Psi(x))} = q_\Psi^{1/\beta}.$$

The details are straightforward and therefore omitted.  $\square$

REMARK 2.4 Condition (2.2) holds for example if  $\Psi(\mathbf{X}) = \|\mathbf{X}\|$ , where  $\|\cdot\|$  is a norm on  $\mathbb{R}^n$  or if  $\Psi(\mathbf{X}) = \sum_{i=1}^n X_i$  for  $X_1, \dots, X_n$  i.i.d.; see Barbe et al. [5], formula (9) and Embrechts et al. [10], Corollary 1.3.2, respectively.  $\square$

### 3 Three examples

Many examples show that VaR properties for rvs with doubly infinite support are not easy to handle, particularly in the case of infinite mean models; see for instance Nešlehová et al. [28], Chavez-Demoulin et al. [7], Ibragimov and Walden [18]. To illustrate this, we give three basic examples:

EXAMPLE 3.1 For  $n \geq 2$ , let  $X_1, \dots, X_n$  be i.i.d. rvs, regularly varying with index  $-\beta < 0$ . In this case, it is well-known that asymptotic subadditivity holds if and only if  $\beta \geq 1$ . This follows from Lemma 2.3, yielding

$$\lim_{\alpha \nearrow 1} \frac{\text{VaR}_\alpha(\sum_{i=1}^n X_i)}{\sum_{i=1}^n \text{VaR}_\alpha(X_i)} = n^{1/\beta-1} > 1, \quad \text{for } \beta < 1,$$

because the limiting constant  $q_\Psi$  in (2.2) is equal to  $n$  for  $\Psi(\mathbf{X}) = \sum_{i=1}^n X_i$ ; see Corollary 1.3.2 in Embrechts et al. [10].  $\square$

When allowing for dependence, one has to be more careful when analyzing additivity properties of VaR; see for instance Example 6.4, Figure 4 below. As shown in Balkema and Embrechts [3] and Barbe [4], for distributions with a density, these questions are closely linked with the properties of the level sets. In the next example, we consider elliptically distributed random vectors.

**DEFINITION 3.2 (ELLIPTICAL DISTRIBUTION)** A random vector  $\mathbf{X}$  has an *elliptical distribution* with mean  $\boldsymbol{\mu} \in \mathbb{R}^n$  and dispersion matrix  $\Sigma$ , if there exist  $R, A$  and  $\mathbf{U}$  satisfying  $\mathbf{X} \stackrel{d}{=} \boldsymbol{\mu} + RA\mathbf{U}$ , with

- a)  $R \geq 0$ , a non-negative rv;
- b)  $\mathbf{U}$  uniformly distributed on the unit sphere  $\mathbb{S}_{\|\cdot\|_2}^{n-1} = \{\mathbf{z} \in \mathbb{R}^n, \|\mathbf{z}\|_2 = 1\}$ , independent of  $R$ , and
- c)  $A \in \mathbb{R}^{n \times n}$  with  $AA' = \Sigma$ .  $\square$

**EXAMPLE 3.3** Theorem 6.8 in McNeil et al. [23] states that for  $\mathbf{X} = (X_1, \dots, X_n)'$  elliptically distributed, we have for all  $\alpha \in [\frac{1}{2}, 1)$ ,

$$\text{VaR}_\alpha \left( \sum_{i=1}^n X_i \right) \leq \sum_{i=1}^n \text{VaR}_\alpha(X_i).$$

That is, in the elliptical world, subadditivity of VaR holds true for finite *and* infinite mean models.  $\square$

What is the reason for this discrepancy between Example 3.1 and Example 3.3? For  $\beta > 1$  (finite mean case) the asymptotic VaR is subadditive in both models. However, for  $\beta < 1$ , we are in the infinite mean regime and the asymptotic VaR behaves very differently in the models analyzed. The reason for this difference is connected with the behavior of the joint df (or more precisely, the *spectral measure*; see Section 4) and can not be explained by the marginal dfs alone. We will discuss risk aggregation

in the light of dependence structures describing interdependencies in the *joint tail*( $s$ ) of the distribution.

In Example 3.3 we learned that subadditivity of VaR holds for every elliptical distribution. However, asymptotic subadditivity of VaR fails for infinite mean models as soon as we weaken the influence of the negative tails by restricting for example to the positive quadrant of the elliptical distribution.

**EXAMPLE 3.4** Let  $\mathbf{X} = RA\mathbf{U}$  be a bivariate elliptical random vector with  $R \in \text{RV}_{-\beta}, \beta > 0$ ,

$$A = \begin{pmatrix} 1 & 0 \\ \varrho & \sqrt{1 - \varrho^2} \end{pmatrix},$$

and  $\mathbf{U}$  uniformly distributed on the unit sphere  $\mathfrak{N}_{\|\cdot\|_2}^1$ , i.e.,  $\mathbf{U} = (\cos W, \sin W)'$ , with  $W \sim \text{Unif}(-\pi, \pi)$ . We are interested in the behavior of  $\mathbf{X} = (X_1, X_2)'$ , restricted to the positive quadrant. We thus consider  $\widetilde{\mathbf{X}} = (\widetilde{X}_1, \widetilde{X}_2)' = \mathbf{X}|\{\mathbf{X} \geq \mathbf{0}\}$ , where the inequality has to be interpreted componentwise. We consider  $q(\beta, \|\cdot\|_1)$  as a function of  $\beta$  and  $\varrho$ . Using the Dominated Convergence Theorem in the last step below, we get

$$\begin{aligned} q(\beta, \varrho) &= \lim_{x \rightarrow \infty} \frac{P(\widetilde{X}_1 + \widetilde{X}_2 > x)}{P(\widetilde{X}_1 > x)} \\ &= \lim_{x \rightarrow \infty} \frac{P\left(R((1 + \varrho)\cos W + \sqrt{1 - \varrho^2}\sin W) > x \mid W \in [-\arcsin \varrho, \pi/2]\right)}{P(R \cos W > x \mid W \in [-\arcsin \varrho, \pi/2])} \\ &= \lim_{x \rightarrow \infty} \frac{\int_{-\arcsin \varrho}^{\pi/2} P\left(R > x / ((1 + \varrho)\cos w + \sqrt{1 - \varrho^2}\sin w)\right) dw}{\int_{-\arcsin \varrho}^{\pi/2} P(R > x / \cos w) dw} \\ &= \frac{\int_{-\arcsin \varrho}^{\pi/2} ((1 + \varrho)\cos w + \sqrt{1 - \varrho^2}\sin w)^\beta dw}{\int_{-\arcsin \varrho}^{\pi/2} \cos^\beta w dw}. \end{aligned} \tag{3.1}$$

□

**Proposition 3.5** Let  $q(\beta, \varrho)$  be defined as in Example 3.4, then

- a) for all  $\varrho \in [-1, 1]$ ,  $q(\beta, \varrho) \leq 2^\beta$  if  $\beta \geq 1$  and  $q(\beta, \varrho) \geq 2^\beta$  if  $\beta \leq 1$ ;
- b)  $\lim_{\varrho \rightarrow -1} q(\beta, \varrho) = 1 + \beta$ , and
- c)  $\lim_{\varrho \rightarrow 1} q(\beta, \varrho) = 2^\beta$ .




Figure 1: The limiting constant  $q$  in (3.1) as a function of  $\rho$  for different values of  $\beta$ .

PROOF: Define for  $f \in L^\beta([-\pi/2, \pi/2])$ ,

$$\zeta_\beta(f) = \left( \int_{-\alpha}^{\pi/2} f^\beta(w) dw \right)^{1/\beta},$$

with a fixed  $\alpha = \arcsin \rho \in [-\pi/2, \pi/2]$  and  $0 < \beta < \infty$ . From (3.1) and some standard trigonometric transformations we get

$$q(\beta, \sin \alpha)^{1/\beta} = \frac{\zeta_\beta(\cos(\cdot) + \sin(\alpha + \cdot))}{\zeta_\beta(\cos(\cdot))}.$$

Applying Minkowski's inequality for  $\beta \geq 1$ , we have

$$q(\beta, \sin \alpha)^{1/\beta} \leq 1 + \frac{\zeta_\beta(\sin(\alpha + \cdot))}{\zeta_\beta(\cos(\cdot))} = 2 \quad \text{for } \beta \geq 1.$$

For  $\beta \leq 1$ , the “ $\leq$ ” turns into a “ $\geq$ ” by Theorem 198 in Hardy et al. [15]. This proves part a). Part b) follows from the representation (3.1) and part c) is a consequence of the comonotonicity of  $X_1$  and  $X_2$  or can be calculated explicitly using (3.1).  $\square$

By part a) of Proposition 3.5 the following corollary follows from Lemma 2.3:



**Corollary 3.6** *Let  $\mathbf{X} \in \text{MRV}_2(-\beta)$ ,  $\beta > 0$ , be an elliptical random vector as in Example 3.4, and  $\widetilde{\mathbf{X}}$  the random vector  $\mathbf{X}$  restricted to the positive quadrant, then VaR is asymptotically subadditive for  $\widetilde{\mathbf{X}}$  if  $\beta \geq 1$  and asymptotically superadditive if  $\beta \leq 1$ .*

The three examples elaborated in this section show that, besides the dependence structure and the tail behavior of the marginal dfs, it is also important to differentiate between rvs with one-sided and two-sided support.

For the infinite mean multivariate  $t$ -distribution, subadditivity of VaR holds due to the dependence properties in the upper left and lower right corner. High values of one risk are compensated by low values of the other risk, turning VaR into a coherent risk measure for such infinite mean models. Of course this has important consequences in risk management. Risk managers should be aware of this property for elliptical distributions, particularly when the compensation of high losses by high gains turns out to be an inappropriate characteristic of the considered risk class.

## 4 Spectral measures for positive rvs

In the following we consider multivariate regularly varying  $\mathbb{R}_+^n$ -valued random vectors. Operations between vectors should be interpreted componentwise. Let  $\|\cdot\| : \mathbb{R}^n \rightarrow \mathbb{R}_+$  be an arbitrary norm. Denote the positive part of the unit sphere with respect to the norm  $\|\cdot\|$  by  $\mathfrak{N}_{+,\|\cdot\|}^{n-1} = \{\mathbf{z} \in \mathbb{R}_+^n \mid \|\mathbf{z}\| = 1\}$ . Note that we write  $\mathfrak{N}_{+,\|\cdot\|}^{n-1}$  for the positive part of  $\mathfrak{N}_{\|\cdot\|}^{n-1}$ . For  $\mathbb{R}_+^n$ -valued random vectors  $\mathbf{X}$ , Theorem 1 in Resnick [31] or Theorem 6.1 in Resnick [32] states that multivariate regular variation of  $\mathbf{X}$  in the sense of Definition 2.2 is equivalent to the existence of a *Radon measure*  $\nu_\beta$  such that

$$\lim_{t \rightarrow \infty} t P(\mathbf{X}/b(t) \in B) = \nu_\beta(B),$$

for all  $B \subset [0, \infty]^n \setminus \{\mathbf{0}\}$  relatively compact with  $\nu_\beta(\partial B) = 0$ . The term *Radon* means that  $\nu_\beta$  is finite for all compact subsets of  $[0, \infty]^n \setminus \{\mathbf{0}\}$ . Resnick [32] calls  $\nu_\beta$  the *limit measure* and, after normalization of the marginal dfs,  $\nu_1$  is referred to as the *exponent*

measure in de Haan and Ferreira [14]. Choosing

$$B = \{\mathbf{z} \in [0, \infty]^n \mid \|\mathbf{z}\| > r, \mathbf{z}/\|\mathbf{z}\| \in G\},$$

for  $r > 0$  and a Borel set  $G \in \mathfrak{K}_{+, \|\cdot\|}^{n-1}$ , we get from Definition 2.2,

$$q(\beta, \|\cdot\|) r^{-\beta} \mu(G) = \nu_\beta \{\mathbf{z} \in [0, \infty]^n \mid \|\mathbf{z}\| > r, \mathbf{z}/\|\mathbf{z}\| \in G\}.$$

For  $\beta = 1$  and  $r = 1$ , this defines the *spectral measure*  $S_{\|\cdot\|}$  by

$$S_{\|\cdot\|}(G) = \nu_1 \{\mathbf{z} \in [0, \infty]^n \mid \|\mathbf{z}\| > 1, \mathbf{z}/\|\mathbf{z}\| \in G\}.$$

Following Barbe et al. [5], we have  $q(\beta, \|\cdot\|) = \nu_1 \{\mathbf{z} \in [0, \infty]^n \mid \|\mathbf{z}^{1/\beta}\| > 1\}$ , and therefore the following theorem.

**Theorem 4.1**

Let  $\mathbf{X} \in \text{MRV}_n(-\beta)$ ,  $\beta > 0$ , be a  $\mathbb{R}_+^n$ -valued random vector with identically distributed marginals, then

$$q(\beta, \|\cdot\|) = \lim_{x \rightarrow \infty} \frac{P(\|\mathbf{X}\| > x)}{P(X_1 > x)} = \int_{\mathfrak{K}_{+, \|\cdot\|}^{n-1}} \|\mathbf{z}^{1/\beta}\|^\beta S_{\|\cdot\|}(d\mathbf{z}).$$

PROOF: Barbe et al. [5] give an explicit proof when  $\|\cdot\|$  is the  $l_1$ -norm in  $\mathbb{R}^n$ . The same proof holds true for general norms in  $\mathbb{R}^n$ , as was certainly noticed by these authors. We therefore refrain from giving the details.  $\square$

Theorem 4.1 shows that  $\beta = 1$  plays an important role in this context. Regardless of our choice of the norm, we have  $q(\beta = 1, \|\cdot\|) = S_{\|\cdot\|}(\mathfrak{K}_{+, \|\cdot\|}^{n-1})$ . If we consider the  $l_1$ -norm, we can give the following result.

**Corollary 4.2** Let  $\mathbf{X} \in \text{MRV}_n(-\beta)$ ,  $\beta > 0$ , be a  $\mathbb{R}_+^n$ -valued random vector with identically distributed marginals. Let  $\|\cdot\|_1$  be the  $l_1$ -norm in  $\mathbb{R}^n$ , then

$$\begin{aligned} n &\leq q(\beta, \|\cdot\|_1) \leq n^\beta, & \text{for } \beta \geq 1, \\ n &\geq q(\beta, \|\cdot\|_1) \geq n^\beta, & \text{for } \beta \leq 1. \end{aligned}$$

PROOF: Proposition 2.2 in Barbe et al. [5] states that  $q(\beta, \|\cdot\|_1)$  is increasing in  $\beta$ . Further,  $q(1, \|\cdot\|_1) = S_{\|\cdot\|_1}(\mathbb{N}_{+, \|\cdot\|_1}^{n-1}) = n$ , because  $S_{\|\cdot\|_1}/n$  is a probability measure. This proves the LHS of the statements. For the RHS, consider the functional

$$\tilde{\zeta}_\beta(f) = \left( \int_{\mathbb{N}_{+, \|\cdot\|_1}^{n-1}} f^\beta(\mathbf{z}) S_{\|\cdot\|_1}(d\mathbf{z}) \right)^{1/\beta},$$

for non-negative functions  $f \in L^\beta(\mathbb{N}_{+, \|\cdot\|_1}^{n-1}, S_{\|\cdot\|_1})$ . Note that for  $\beta \geq 1$ , by Minkowski's inequality (note the slight abuse of notation),

$$(q(\beta, \|\cdot\|_1))^{1/\beta} = \tilde{\zeta}_\beta \left( \sum_{i=1}^n z_i^{1/\beta} \right) \leq \sum_{i=1}^n \tilde{\zeta}_\beta(z_i^{1/\beta}) = \sum_{i=1}^n \left( \int_{\mathbb{N}_{+, \|\cdot\|_1}^{n-1}} z_i S_{\|\cdot\|_1}(d\mathbf{z}) \right)^{1/\beta} = n.$$

For  $\beta \leq 1$  the “ $\leq$ ” turns into a “ $\geq$ ” by Theorem 198 in Hardy et al. [15].  $\square$

### Theorem 4.3

Let  $\mathbf{X} \in \text{MRV}_n(-\beta)$ ,  $\beta > 0$ , be a  $\mathbb{R}_+^n$ -valued random vector with identically distributed marginals, then  $\text{VaR}_\alpha$  is asymptotically subadditive for  $\mathbf{X}$  if  $\beta \geq 1$  and asymptotically superadditive if  $\beta \leq 1$ .

PROOF: Lemma 2.3 and Corollary 4.2 yield the result.  $\square$

Asymptotic subadditivity for bivariate regularly varying random vectors with  $\beta \geq 1$  has already been proven in Daniélsson et al. [8], Proposition 1.

REMARK 4.4 Note that all components of  $\mathbf{X}$  in Theorem 4.3 need to be positive. If this assumption is not fulfilled, subadditivity also for infinite mean models may occur, for example for elliptical distributed random vectors; see Example 3.3.  $\square$

The norm  $\|\mathbf{z}\|_1 = |z_1| + \dots + |z_n|$  is a natural choice, because it allows for the study of sums of dependent, positive risks and in particular for an analysis of the additivity properties of VaR; see Theorem 4.3. Sometimes however, spectral measures with respect to other norms are chosen; for instance in Stărică [35] and Hult and Lindskog [16], the spectral measure with respect to  $\|\cdot\|_2$  and  $\|\cdot\|_\infty$ , respectively, is more convenient in their context.

Also when one deals with elliptical types of distributions, where (after a linear transformation of  $\mathbf{X}$ ) the spectral measure with respect to the Euclidean norm  $\|\cdot\|_2$  is uniformly distributed on  $\mathfrak{N}_{\|\cdot\|_2}^{n-1}$ , a change of measure could be appropriate. It is thus important to express the spectral measure with respect to one norm in terms of the spectral measure with respect to another norm. This can always be done; see for instance formula (8.38) in Beirlant et al. [6], which we formulate in the following lemma.

**Lemma 4.5** *Let  $S_{\|\cdot\|}$  and  $S_{\|\cdot\|'}$  be the spectral measure with respect to the norms  $\|\cdot\|$  and  $\|\cdot\|'$ , respectively, then*

$$S_{\|\cdot\|}(G) = \int_{\mathfrak{N}_{+,\|\cdot\|'}^{n-1}} \mathbf{1}_{\{\mathbf{z}/\|\mathbf{z}\| \in G\}} \|\mathbf{z}\| S_{\|\cdot\|'}(d\mathbf{z}),$$

for any Borel set  $G \subset \mathfrak{N}_{+,\|\cdot\|}^{n-1}$ .

We call a  $\mathbb{R}_+^n$ -valued multivariate regularly varying random vector asymptotically independent, if the spectral measure  $S_{\|\cdot\|}$  is concentrated on the points  $\mathbf{e}_i/\|\mathbf{e}_i\|$ ,  $i = 1, \dots, n$ , with  $\mathbf{e}_i$  the  $i$ th basis vector of the canonical basis in  $\mathbb{R}^n$ ; it is called asymptotically fully dependent, if the spectral measure  $S_{\|\cdot\|}$  is concentrated on  $\mathbf{1}/\|\mathbf{1}\|$ , with  $\mathbf{1} = (1, \dots, 1)'$ ; see Resnick [31]. Note that by Lemma 4.5 asymptotic independence as well as asymptotic full dependence is well-defined.

**Proposition 4.6** *Let  $\mathbf{X} \in \text{MRV}_n(-\beta)$ ,  $\beta > 0$ , be an asymptotically independent  $\mathbb{R}_+^n$ -valued random vector with identically distributed marginals, then  $q(\beta, \|\cdot\|) = \sum_{i=1}^n \|\mathbf{e}_i\|^\beta$  and in particular, if  $\|\mathbf{e}_i\| = 1$  for all  $i = 1, \dots, n$ ,  $q(\beta, \|\cdot\|) = n$ .*

PROOF: Theorem 4.1 yields

$$q(\beta, \|\cdot\|) = \sum_{i=1}^n \|\mathbf{e}_i/\|\mathbf{e}_i\|\|^{1/\beta} \|\mathbf{e}_i/\|\mathbf{e}_i\|\|^\beta S_{\|\cdot\|}(\mathbf{e}_i/\|\mathbf{e}_i\|) = \sum_{i=1}^n \|\mathbf{e}_i\|^{\beta-1} S_{\|\cdot\|}(\mathbf{e}_i/\|\mathbf{e}_i\|),$$

with  $S_{\|\cdot\|}(\mathbf{e}_i/\|\mathbf{e}_i\|) = \|\mathbf{e}_i\| S_{\|\cdot\|_1}(\mathbf{e}_i) = \|\mathbf{e}_i\|$ , by Lemma 4.5 and because  $S_{\|\cdot\|_1}/n$  is a probability measure.  $\square$

This proposition generalizes Lemma 2.1 in Davis and Resnick [9] to arbitrary norms; see also Lemma 3.1 in Jessen and Mikosch [20], where the result from [9] is generalized to rvs with doubly infinite support.

**Proposition 4.7** *Let  $\mathbf{X} \in \text{MRV}_n(-\beta)$ ,  $\beta > 0$ , be an asymptotically fully dependent  $\mathbb{R}_+^n$ -valued random vector with identically distributed marginals, then  $q(\beta, \|\cdot\|) = \|\mathbf{1}\|^\beta$ .*

PROOF: Theorem 4.1 yields

$$q(\beta, \|\cdot\|) = \|(\mathbf{1}/\|\mathbf{1}\|)^{1/\beta}\|^\beta S_{\|\cdot\|}(\mathbf{1}/\|\mathbf{1}\|) = \|\mathbf{1}\|^{\beta-1} S_{\|\cdot\|}(\mathbf{1}/\|\mathbf{1}\|),$$

with  $S_{\|\cdot\|}(\mathbf{1}/\|\mathbf{1}\|) = \|\mathbf{1}/n\| S_{\|\cdot\|_1}(\mathbf{1}/n) = \|\mathbf{1}\|$ , by Lemma 4.5 and because  $S_{\|\cdot\|_1}/n$  is a probability measure.  $\square$

**Proposition 4.8** *Let  $\mathbf{X} \in \text{MRV}_n(-\beta)$ ,  $\beta > 0$ , be a  $\mathbb{R}_+^n$ -valued random vector with identically distributed marginals. Let  $S_{\|\cdot\|_\infty}$  be the spectral measure with respect to  $\|\cdot\|_\infty$ , the maximum-norm in  $\mathbb{R}^n$ , then*

$$q(\beta, \|\cdot\|_\infty) = S_{\|\cdot\|_\infty}(\mathbb{N}_{+, \|\cdot\|_\infty}^{n-1}) = \int_{\mathbb{N}_{+, \|\cdot\|_\infty}^{n-1}} \bigvee_{i=1}^n z_i S_{\|\cdot\|_\infty}(d\mathbf{z}).$$

PROOF: Note that  $\|\mathbf{z}\|_\infty^{1/\beta} = 1$  on  $\mathbb{N}_{+, \|\cdot\|_\infty}^{n-1}$ , for all  $\beta > 0$ . Hence, the first equality follows from Theorem 4.1. Using Lemma 4.5 the second equality follows.  $\square$

The following well-known result characterizes asymptotic independence and full dependence.

**Corollary 4.9** *Let  $\mathbf{X} \in \text{MRV}_n(-\beta)$ ,  $\beta > 0$ , be a  $\mathbb{R}_+^n$ -valued random vector with identically distributed marginals, then*

*i)  $\mathbf{X}$  is asymptotically independent if and only if*

$$\int_{\mathbb{N}_{+, \|\cdot\|_\infty}^{n-1}} \bigvee_{i=1}^n z_i S_{\|\cdot\|_\infty}(d\mathbf{z}) = n.$$

ii)  $\mathbf{X}$  is asymptotically fully dependent if and only if

$$\int_{\mathbb{R}_{+, \|\cdot\|}^{n-1}} \prod_{i=1}^n z_i S_{\|\cdot\|}(d\mathbf{z}) = 1.$$

PROOF: The “ $\Rightarrow$ ”-part is straightforward from the definition of asymptotic independence and full dependence, but also a consequence of Propositions 4.6, 4.7 and 4.8. For the converse, see Beirlant et al. [6], Section 8.2.7.  $\square$

By Proposition 4.8 and Corollary 4.9, it suffices to evaluate  $q(\beta, \|\cdot\|_\infty)$  in order to test for asymptotic independence and full dependence, respectively.

## 5 Tail dependence and asymptotic independence

In Sections 5 and 6, we will discuss further examples and counter-examples for sub-additivity of VaR. We restrict ourselves to the bivariate case and only sums of rvs are considered. Since the marginal dfs have equal asymptotic behavior in the different infinite mean models in Examples 3.1 and 3.3, the asymptotic VaR behavior for the sum of the risks must follow from the different dependence structures (copulas, spectral measures). We exemplify this issue through the notions of asymptotic dependence coefficients in the (four) tails of the underlying bivariate distribution.

DEFINITION 5.1 Let  $(X_1, X_2)'$  be a bivariate random vector, with marginal dfs  $F_{X_1}$  and  $F_{X_2}$ . The *positive upper* ( $\lambda_u^+$ ), *positive lower* ( $\lambda_l^+$ ), *negative upper* ( $\lambda_u^-$ ) and *negative lower* ( $\lambda_l^-$ ) *tail dependence coefficients* are defined as

$$\begin{aligned} \lambda_u^+ &= \lim_{\alpha \nearrow 1} P(X_2 > F_{X_2}^{\leftarrow}(\alpha) | X_1 > F_{X_1}^{\leftarrow}(\alpha)), \\ \lambda_l^+ &= \lim_{\alpha \searrow 0} P(X_2 \leq F_{X_2}^{\leftarrow}(\alpha) | X_1 \leq F_{X_1}^{\leftarrow}(\alpha)), \\ \lambda_u^- &= \lim_{\alpha \nearrow 1} P(X_2 > F_{X_2}^{\leftarrow}(\alpha) | X_1 \leq F_{X_1}^{\leftarrow}(1 - \alpha)), \\ \lambda_l^- &= \lim_{\alpha \searrow 0} P(X_2 \leq F_{X_2}^{\leftarrow}(\alpha) | X_1 > F_{X_1}^{\leftarrow}(1 - \alpha)), \end{aligned}$$

provided the limits exist in  $[0, 1]$ .  $\square$

A sufficient condition for the existence of the tail dependence coefficient is bivariate regular variation of  $(f(X_1), f(X_2))'$  for some strictly increasing transformation  $f$ ; see Mikosch [25] and references therein for weaker conditions on  $(X_1, X_2)'$ .

Note that for  $(X_1, X_2)' \in \text{MRV}_2(-\beta)$ ,  $\beta > 0$ , a  $\mathbb{R}_+^2$ -valued random vector with identically distributed marginals, we have

$$\lambda_u^+ = \lim_{x \rightarrow \infty} P(X_2 > x | X_1 > x) = 2 - \lim_{x \rightarrow \infty} \frac{P(\max(X_1, X_2) > x)}{P(X_1 > x)} = 2 - q(\beta, \|\cdot\|_\infty),$$

and hence by Proposition 4.8,  $\lambda_u^+ = 2 - S_{\|\cdot\|_\infty}(\mathfrak{N}_{+, \|\cdot\|_\infty}^1)$ .

**Proposition 5.2** *Let  $(X_1, X_2)' \in \text{MRV}_2(-\beta)$ ,  $\beta > 0$ , be a  $\mathbb{R}_+^2$ -valued random vector with identically distributed marginals, then*

$$\lambda_u^+ = 0 \iff (X_1, X_2)' \text{ asymptotically independent.}$$

**PROOF:** By Proposition 4.8 and Corollary 4.9, asymptotic independence of the random vector  $(X_1, X_2)'$  is equivalent to  $2 = q(\beta, \|\cdot\|_\infty)$ . This is equivalent to  $\lambda_u^+ = 0$ .  $\square$

**REMARK 5.3** The concept of positive tail dependence is well-known and often used in risk management, in particular for describing so-called spillover events; see for instance Straetmans [33]. However, negative tail dependence  $\lambda_u^-, \lambda_l^-$ , i.e., the probability that high values of  $X_1$  are compensated by low values of  $X_2$  and vice versa, did not draw risk managers' attention so far. We will see its importance in the sequel. High negative tail dependence might not always be reasonable in reality. If it is not appropriate that high losses are compensated by high gains with positive probability, then more conservative models should be considered. Recently Zhang [36] introduced negative tail dependence in order to define a novel dependence measure called *total tail dependence*, which is a  $(2 \times 2)$ -matrix with components  $\lambda_u^+, \lambda_l^+, \lambda_u^-, \lambda_l^-$ .  $\square$

The tail dependence coefficients do not depend on the marginal dfs and thus can be written in terms of the corresponding copulas.

**Proposition 5.4** *Let  $F_{X_1}$  and  $F_{X_2}$  from Definition 5.1 be continuous dfs and  $C$  the corresponding copula, then*

$$\lambda_u^+ = \lim_{\alpha \nearrow 1} \frac{1 - 2\alpha + C(\alpha, \alpha)}{1 - \alpha}, \quad (5.1)$$

$$\lambda_l^+ = \lim_{\alpha \searrow 0} \frac{C(\alpha, \alpha)}{\alpha}, \quad (5.2)$$

$$\lambda_u^- = 1 - \lim_{\alpha \nearrow 1} \frac{C(1 - \alpha, \alpha)}{1 - \alpha}, \quad (5.3)$$

$$\lambda_l^- = 1 - \lim_{\alpha \searrow 0} \frac{C(1 - \alpha, \alpha)}{\alpha}. \quad (5.4)$$

PROOF: See for instance Joe [21], Section 2.1.10 or McNeil et al. [23], Section 5.2.3 for the proof of (5.1) and (5.2). The proof of (5.3) and (5.4) is completely analogous.  $\square$

In the case of regularly varying elliptical distributions, the four tail dependence coefficients can be calculated explicitly.

**Proposition 5.5** *Let  $\mathbf{X} \stackrel{d}{=} \boldsymbol{\mu} + R\mathbf{A}\mathbf{U}$  be a bivariate elliptically distributed regularly varying random vector with index  $-\beta < 0$ , as defined in Definition 3.2, then*

$$\lambda_u^+ = \lambda_l^+ = \frac{\int_{a_+}^{\pi/2} \cos^\beta t \, dt}{\int_0^{\pi/2} \cos^\beta t \, dt}, \quad (5.5)$$

$$\lambda_u^- = \lambda_l^- = \frac{\int_{a_-}^{\pi/2} \cos^\beta t \, dt}{\int_0^{\pi/2} \cos^\beta t \, dt}, \quad (5.6)$$

with  $a_+ = (\pi/2 - \arcsin \varrho)/2$ ,  $a_- = (\pi/2 + \arcsin \varrho)/2$ , and where  $\varrho = \sigma_{12}/\sqrt{\sigma_{11}\sigma_{22}}$  with  $(\sigma_{ij})_{1 \leq i, j \leq 2} = \Sigma = AA'$ .

PROOF: Equation (5.5) has been proven in Hult and Lindskog [16]. By considering the map  $\mathbf{X} \mapsto D\mathbf{X}$ , with

$$D = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

(5.6) follows from (5.5).  $\square$



Even for rvs with a positive linear correlation coefficient,  $\lambda_u^-$  can be significantly larger than zero. Another important consequence of the previous proposition is that there exists *no* elliptical distribution without negative tail dependence and with heavy (i.e., regularly varying) tails, provided  $\rho < 1$ .

**REMARK 5.6** The class of *Archimedean* copulas contains several dependence structures important for practical purposes; see Nelsen [27] Section 4.1, for a definition of an Archimedean copula and further results. An interesting connection between Archimedean copulas and so-called  $l_1$ -norm symmetric distributions is established by McNeil and Nešlehová [29]. One can show that bivariate dfs with continuous marginals and with certain Archimedean copulas (e.g., with strict generator; see Nelsen [27]) have no negative tail dependence, that is  $\lambda_u^- = \lambda_l^- = 0$ . Therefore, they stand in violent contrast to elliptical distributions, where (unless in the comonotonic case)  $\lambda_u^-$  and  $\lambda_l^-$  are always positive.  $\square$

For every elliptical copula one can always find a (strict) Archimedean copula with the same positive upper tail dependence coefficient  $\lambda_u^+$ . However, the asymptotic VaRs behave very differently; see also Embrechts et al. [12]. We hence conclude that the positive upper tail dependence coefficient in an infinite mean model is not able to explain the sub-/superadditive behavior of VaR.

In the next section we show that a crucial role is indeed played by  $\lambda_u^-$  and  $\lambda_l^-$  whenever  $X_1$  and  $X_2$  have doubly infinite support.

## 6 Tail dependence and subadditivity

The simplest model incorporating independence as well as co- and countermonotonicity is the *Fréchet family*. Therefore, we combine the independent copula  $C_{0,0}(u, v) = uv$ , the comonotonic copula  $C_{1,0}(u, v) = u \wedge v = \min(u, v)$  and the countermonotonic copula  $C_{0,1}(u, v) = (u + v - 1)^+$ ; see Nelsen [27], Exercise 2.4. Let  $C_{p_1, p_2}$  be a convex combination of these copulas,

$$C_{p_1, p_2}(u, v) = p_1(u \wedge v) + p_2(u + v - 1)^+ + (1 - p_1 - p_2)uv, \quad (6.1)$$

for  $p_1 \in [0, 1]$  and  $p_2 \in [0, 1 - p_1]$ . The copula family  $C_{p_1, p_2}$  is referred to as the *Fréchet family*. Let  $X_1, X_2$  be two identically distributed regularly varying rvs with index  $-\beta < 0$  and with marginal df  $F_\beta$  (i.e.,  $F_\beta(x) = x^{-\beta}L(x)$ , with  $L$  slowly varying) and copula  $C_{p_1, p_2}$ . The bivariate df of  $(X_1, X_2)'$  is then given by

$$F_{X_1, X_2}(x_1, x_2) = C_{p_1, p_2}(F_\beta(x_1), F_\beta(x_2)). \quad (6.2)$$

In Figure 2 we give a typical random sample from the copula  $C_{p_1, p_2}$  in (6.1) and the bivariate df in (6.2).




Figure 2: Left panel: 500 realizations of the copula  $C_{p_1, p_2}$  in (6.1) with parameters  $p_1 = p_2 = 0.1$ . Right panel: the realizations are transformed according to (6.2) where  $F_\beta$  is a  $t$  distribution with 6 degrees of freedom.

In the following, we only look at symmetric marginals, i.e., where  $X \stackrel{d}{=} -X$ . In order to investigate subadditivity properties of VaR for the df (6.2), we consider  $q_\Psi$  as a function of  $p_1, p_2$  and with  $\Psi(\mathbf{X}) = X_1 + X_2$ ,

$$q_\Psi(\beta, (p_1, p_2)) = \lim_{x \rightarrow \infty} \frac{P(X_1 + X_2 > x)}{P(X_1 > x)}.$$

In the symmetric case,  $q_\Psi$  can be calculated explicitly.

**Proposition 6.1** *Let  $(X_1, X_2)'$  be a bivariate random vector defined by (6.2) with identically distributed, symmetric, regularly varying marginals with index  $-\beta < 0$ ,*

then

$$q_{\Psi}(\beta, (p_1, p_2)) = 2^{\beta}p_1 + 2(1 - p_1 - p_2).$$

PROOF: Due to the linearity of (6.1) it is sufficient to check the independent, the comonotonic and the countermonotonic case separately. For  $X_1, X_2$  independent, we have  $q_{\Psi}(\beta, (0, 0)) = 2$ , which does not depend on  $\beta$ . For  $X_1, X_2$  comonotonic and  $X_1 \stackrel{d}{=} X_2$ , we have  $X_1 = X_2$   $P$ -a.s. and hence  $q_{\Psi}(\beta, (1, 0)) = \lim_{x \rightarrow \infty} (x/2)^{-\beta} / x^{-\beta} = 2^{\beta}$ . For  $X_1, X_2$  countermonotonic and  $X_1 \stackrel{d}{=} X_2$ , we have  $X_1 = -X_2$   $P$ -a.s. and hence  $q_{\Psi}(\beta, (0, 1)) = 0$ .  $\square$

Note that the positive upper tail dependence coefficient of the model (6.2) is given by

$$\lambda_u^+ = \lim_{u \nearrow 1} \frac{1 - 2u + C(u, u)}{1 - u} = p_1.$$

Equivalently, we have  $\lambda_l^+ = p_1$  and  $\lambda_u^- = \lambda_l^- = p_2$ . Thus, we have the following corollary.

**Corollary 6.2** *Let  $(X_1, X_2)'$  be a bivariate random vector with a copula from the Fréchet family defined by (6.1) and identically distributed, symmetric, regularly varying marginals with index  $-\beta < 0$ , then  $q_{\Psi}(\beta, (\lambda_u^+, \lambda_u^-)) = 2^{\beta}\lambda_u^+ + 2(1 - \lambda_u^+ - \lambda_u^-)$ , with  $\lambda_u^+, \lambda_u^- \geq 0$  and  $\lambda_u^+ + \lambda_u^- \leq 1$ .*

If  $\beta \geq 1$ , then of course, asymptotic subadditivity always holds. This follows from the fact that  $q_{\Psi}(\beta, (\lambda_u^+, \lambda_u^-)) = 2^{\beta}\lambda_u^+ + 2(1 - \lambda_u^+ - \lambda_u^-) \leq 2^{\beta}(\lambda_u^+ + (1 - \lambda_u^+ - \lambda_u^-)) \leq 2^{\beta}$ , together with Lemma 2.3, yielding that

$$\lim_{\alpha \nearrow 1} \frac{\text{VaR}_{\alpha}(X_1 + X_2)}{\text{VaR}_{\alpha}(X_1) + \text{VaR}_{\alpha}(X_2)} \leq 1.$$

In the case  $\beta < 1$ ,  $q_{\Psi}(\beta, (\lambda_u^+, \lambda_u^-))$  can be smaller or larger than  $2^{\beta}$ , thus depending on the values  $\lambda_u^+, \lambda_u^-$ , subadditivity may hold or fail. To analyze this infinite mean model in more detail, we exclude the trivial case  $\lambda_u^+ = 1$  and introduce the *relative*

*negative tail dependence coefficient*

$$\gamma = \frac{\lambda_u^-}{1 - \lambda_u^+}.$$

Because  $\lambda_u^+ + \lambda_u^-$  is always smaller than 1,  $\gamma$  takes values only in  $[0, 1]$  and is interpreted as the amount of negative tail dependence, relative to the possible maximal negative tail dependence coefficient  $1 - \lambda_u^+$ . We then have the following theorem.

**Theorem 6.3**

*Let  $(X_1, X_2)'$  be a bivariate random vector with a copula from the Fréchet family defined by (6.1) with  $p_1 < 1$  and identically distributed, symmetric, regularly varying marginals with index  $-\beta < 0$ . Then asymptotic subadditivity of VaR holds if and only if  $\gamma \geq 1 - 2^{\beta-1}$ .*

PROOF: This follows immediately from Corollary 6.2 and Lemma 2.3. □




Figure 3: This figure shows the impact of negative tail dependence on subadditivity properties of VaR for infinite mean models. Inside of the hatched area subadditivity holds, whereas outside superadditivity holds.

Theorem 6.3 provides a simple criterion for asymptotic subadditivity in the case of

the Fréchet family model (6.2). Only for sufficiently small values of  $\gamma$  superadditivity occurs. For large values of  $\gamma$  subadditivity always holds. The interpretation of this behavior is that if the negative tail dependence coefficient is sufficiently large, then positive extreme values in one coordinate are compensated by negative extreme values in the other coordinate. This effect can be so strong that we obtain asymptotic subadditivity. In Figure 3, we plot the range, where subadditivity occurs as a function of  $\beta \in [0, 1]$  and the relative negative tail dependence coefficient  $\gamma$ .

Several authors mention the substantial influence of the transition from a finite to an infinite mean model on the additivity properties of VaR; see for instance Nešlehová et al. [28], Embrechts et al. [12], Ibragimov and Walden [18] and Jang and Jho [19]. An early statement in the finance literature that *diversification does not in general reduce its effects on the dispersion of the portfolio return* is found already in 1972 in Fama and Miller [13]; these authors based their conclusion on properties of Lévy-stable dfs.

Theorem 6.3 also shows that  $\beta = 1$  plays a fundamental role for independent rvs and more generally if (in the Fréchet model) the relative negative tail dependence coefficient  $\gamma$  is 0. However, as soon as  $\gamma > 0$ , it is likely that high losses are compensated by high gains and therefore the transition from sub- to superadditivity will be located at  $\beta$  strictly smaller than 1; see again Figure 3.

Theorem 6.3 of course does not hold in general (outside the Fréchet family model). However it gives first insight in the still open problem of the characterization of asymptotic sub- and superadditivity in an infinite mean model. A more in depth analysis would presumably need to be based on properties of the level sets of the densities. Consider as an example a bivariate distribution with identical  $t_\nu$  marginals and a  $t_4$ -copula with dependence  $\rho = 0$ . This is an example from the so-called class of meta- $t_4$ -distributions. McNeil et al. [23], Paragraph 5.1.3 contains a general definition of meta-distributions, as well as a discussion of applications to risk management. A simulation with  $10^5$  realizations then shows that the transition from sub- to superadditivity is located at a value  $\nu_0$  in the interval  $(0.8, 0.9)$ , significantly below 1. If Theorem 6.3 would hold in general, it would indicate a (theoretical) transition located at  $\nu_0 = \log_2(1 - \gamma) + 1 \approx 0.877 \in (0.8, 0.9)$ , in agreement with our empirical result. This simulation-based statement we have included for illustrative purposes, because to the best of our knowledge one does not have a satisfactory explanation

for this anomaly at the moment.

In Section 3, we mentioned that in an infinite mean model sub- as well as superadditivity of VaR may occur in a completely arbitrary (or somewhat chaotic) way. In the following example we construct such an (artificial) model.

**EXAMPLE 6.4** Corollary 6.2 shows that there exist bivariate rvs where the positive upper tail dependence coefficient is not the main driver leading to sub- or superadditivity of VaR.




Figure 4:  $q_{\Psi}(\beta, \lambda_u^+)$  as a function of  $\lambda_u^+$  for the Fréchet family copula (6.1) with parameters  $p_1$  and  $p_2$  from (6.3) and (6.4), respectively. Subadditivity occurs below the horizontal line, superadditivity above.

Indeed, by choosing  $\beta = 1/2$  (infinite mean model) and for instance

$$p_1 = \lambda_u^+, \quad (6.3)$$

$$p_2 = (1 - \lambda_u^+) \sin^2 \left( 4\pi \frac{\lambda_u^+}{1 - \lambda_u^+} \right), \quad (6.4)$$

the plot of  $q_{\Psi}(\beta, \lambda_u^+)$  in Figure 4 clearly shows that there is no obvious connection between the positive upper tail dependence coefficient and subadditivity of VaR. Asymptotic subadditivity occurs if and only if  $q_{\Psi}(\beta = 1/2, \lambda_u^+) \leq \sqrt{2}$  (horizontal

line). One can always choose  $p_2 = \lambda_u^-$  such that for an arbitrary value of  $\lambda_u^+ < 1$ ,  $q_\Psi(\beta, \lambda_u^+)$  is greater *or* smaller than  $\sqrt{2}$ . Sub- as well as superadditivity occurs in a completely arbitrary way.  $\square$

Note that in Theorem 6.3 we assume the rvs to have doubly infinite support. We give an explicit counter-example, when this assumption is not fulfilled.

**EXAMPLE 6.5** Let  $X_1, X_2$  be two countermonotonic (i.e.,  $\gamma = 1$ ) Pareto distributed rvs with marginal dfs

$$F_{X_1}(x) = F_{X_2}(x) = 1 - x^{-1/2}, \quad x \geq 1.$$

Using countermonotonicity of  $X_1$  and  $X_2$ , we deduce that the df of  $X_1 + X_2$  for all  $x \geq 8$  is,

$$F_{X_1+X_2}(x) = \sqrt{1 + 4(1 - \sqrt{1+x})/x};$$

see for instance Strassburger and Pfeifer [34], Lemma 5.1. Hence,

$$F_{X_1+X_2}(x) = 1 - 2x^{-1/2} + O(x^{-3/2}), \quad x \rightarrow \infty,$$

and therefore

$$\lim_{x \rightarrow \infty} \frac{P(X_1 + X_2 > x)}{P(X_1 > x)} = 2.$$

Thus by Lemma 2.3, asymptotic subadditivity does not hold.  $\square$

For  $X'_1, X'_2 \stackrel{i.i.d.}{\sim} \text{Pareto}(1/2)$ , for  $x \geq 1$ ,

$$F_{X'_1+X'_2}(x) = 1 - 2\frac{\sqrt{x-1}}{x}.$$

Hence,

$$F_{X'_1+X'_2}(x) = 1 - 2x^{-1/2} + O(x^{-3/2}), \quad x \rightarrow \infty,$$

and therefore  $X_1 + X_2$  in Example 6.5 and  $X'_1 + X'_2$  are tail-equivalent. Hence, full diversification in the sense of countermonotonicity is as bad as independence (and therefore worse than no diversification in the sense of comonotonicity). For infinite mean models, diversification clearly goes the wrong way. More generally, we have the

following proposition; see for instance Davis and Resnick [9], Lemma 2.1 or Albrecher et al. [2], Corollary 3.2.

**Proposition 6.6** *Let  $X_1, X_2 > 0$  be two identically distributed regularly varying rvs with index  $-\beta < 0$  and with  $\lambda_u^+ = 0$ , i.e., such that  $X_1, X_2$  are tail-independent in the positive upper tail, then*

$$\lim_{x \rightarrow \infty} \frac{P(X_1 + X_2 > x)}{P(X_1 > x)} = 2.$$

This proposition is indeed a special case of Proposition 4.6. It has the nice interpretation that the sum of tail-independent (in the positive upper tail) rvs behaves asymptotically as if the summands were independent. According to the above remarks, Proposition 6.6 yields the following result. Let  $X_1, X_2, X'_1, X'_2 > 0$  be identically distributed regularly varying rvs with  $X_1, X_2$  countermonotonic and  $X'_1, X'_2$  independent, then  $\text{VaR}_\alpha(X_1 + X_2) \sim \text{VaR}_\alpha(X'_1 + X'_2)$ , for  $\alpha \rightarrow 1$ . That is, the VaR of the sum of highly “diversified” positive rvs is asymptotically equal to the VaR of the sum of independent rvs for  $\alpha$  large.

## 7 Conclusion

Under the current regulatory guidelines for banking and insurance, risk diversification, concentration and aggregation play a prominent role. Within and between “nice” risk categories, with elliptical dfs, say, these concepts are easily modeled and particular solutions can be readily worked out. However, for skew, heavy-tailed risk rvs, diversification and aggregation have to be handled with care.

In this paper we show that the interplay between existence, non-existence of a finite moment, one- or two-sidedness and symmetry versus asymmetry of the underlying risk dfs have to be carefully balanced in order to be able to conclude sub- or superadditivity of quantile based risk measures like Value-at-Risk.

We have highlighted in the paper that MEVT offers the canonical language for analyzing from an asymptotic point of view questions of the above type for heavy-tailed dfs. That answers to these questions are relevant for practice can for instance be seen in applied publications like Moscadelli [26] and Aas et al. [1]. Though we



obtained a better understanding of the diversification-concentration-aggregation issue for VaR, many questions still remain unsolved and further research is no doubt needed, especially for two-sided skew rvs. For two-sided rvs the sum operator is not a norm but only a so-called gauge function and this makes their analysis much more delicate; see for instance Balkema and Embrechts [3] for details on MEVT in this context.

The following recent preprint came to our attention during the refereeing process, Mainik and Rüschendorf [22], in which statistical estimation procedures for problems presented in Section 4 are discussed.

### Acknowledgments

The authors would like to thank Johanna Nešlehová and two anonymous referees for their useful comments.

## References

- [1] Aas, K., Dimakos, X. K. and Øksendal, A. (2007) Risk capital aggregation. *Risk Management* **9**, 82–107.
- [2] Albrecher, H., Asmussen, S. and Kortschak, D. (2006) Tail asymptotics for the sum of two heavy-tailed dependent risks. *Extremes* **9**, 107–130.
- [3] Balkema, G. and Embrechts, P. (2007) *High Risk Scenarios and Extremes. A geometric approach*. Zurich Lectures in Advanced Mathematics, European Mathematical Society Publishing House, Zurich.
- [4] Barbe, P. (2003) Approximation of integrals over asymptotic sets with applications to statistics and probability. <http://arxiv.org/abs/math/0312132>.
- [5] Barbe, P., Fougères, A. and Genest, C. (2006) On the tail behavior of sums of dependent risks. *Astin Bulletin* **36**, 361–373.
- [6] Beirlant, J., Goegebeur, Y., Segers, J. and Teugels, J. (2004) *Statistics of Extremes - Theory and Applications*. Wiley, Chichester.

- [7] Chavez-Demoulin, V., Embrechts, P. and Nešlehová, J. (2006) Quantitative models for operational risk: extremes, dependence and aggregation. *Journal of Banking and Finance* **30**(10), 2635–2658.
- [8] Daníelsson, J., Jorgensen, B. N., Samorodnitsky, G., Sarma, M. and de Vries, C. G. (2005) Subadditivity re-examined: the case for Value-at-Risk. FMG Discussion Papers, London School of Economics.
- [9] Davis, R. A. and Resnick, S. I. (1996) Limit theory for bilinear processes with heavy-tailed noise. *The Annals of Applied Probability* **6**, 1191–1210.
- [10] Embrechts, P., Klüppelberg, C. and Mikosch, T. (1997) *Modelling Extremal Events for Insurance and Finance*. Springer, Berlin.
- [11] Embrechts, P., McNeil, A. J. and Straumann, D. (2002) Correlation and dependence in risk management: properties and pitfalls. Dempster, M. (Ed.), *Risk management: Value at Risk and Beyond*, pp. 176–223, Cambridge University Press, Cambridge.
- [12] Embrechts, P., Nešlehová, J. and Wüthrich, M. V. (2008) Additivity properties for Value-at-Risk under Archimedean dependence and heavy-tailedness. *Insurance: Mathematics and Economics*. To appear.
- [13] Fama, E. T. and Miller, M. H. (1972) *The Theory of Finance*. Dryden Press, Hinsdale.
- [14] Haan, L. de and Ferreira, A. (2006) *Extreme Value Theory, An Introduction*. Springer, New York.
- [15] Hardy, G. H., Littlewood, J. E. and Pólya, G. (1934) *Inequalities*. Cambridge University Press, Cambridge.
- [16] Hult, H. and Lindskog, F. (2002) Multivariate extremes, aggregation and dependence in elliptical distributions. *Advances in Applied Probability* **34**, 587–608.
- [17] Ibragimov, R., Jaffee, D. and Walden, J. (2008) Nondiversification traps in catastrophe insurance markets. *Review of Financial Studies*. To appear.

- [18] Ibragimov, R. and Walden, J. (2007) The limits of diversification when losses may be large. *Journal of Banking and Finance* **31**(8), 2551–2569.
- [19] Jang, J. and Jho, J. H. (2007) Asymptotic super(sub)additivity of Value-at-Risk of regularly varying dependent variables. Preprint MacQuarie University, Sydney.
- [20] Jessen, H., A. and Mikosch, T. (2006) Regularly varying functions. *Publications de l'Institut Mathématique, Nouvelle Série* **80**, 171–192.
- [21] Joe, H. (1997) *Multivariate Models and Dependence Concepts*. Chapman and Hall, London.
- [22] Mainik, G. and Rüschendorf, L. (2008) On optimal portfolio diversification with respect to extreme risks. Preprint Albert-Ludwigs-Universität Freiburg; April 28, 2008.
- [23] McNeil, A. J., Frey, R. and Embrechts, P. (2005) *Quantitative Risk Management: Concepts, Techniques, Tools*. Princeton University Press, Princeton.
- [24] Mikosch, T. (2004) Modeling dependence and tails of financial time series. In: Finkelstädt, B. and Rootzén, H. (Eds.), *Extreme Values in Finance, Telecommunications and the Environment*, pp. 185–286, Chapman & Hall, Boca Raton.
- [25] Mikosch, T. (2006) Copulas: Tales and facts. *Extremes* **9**, 3–20.
- [26] Moscadelli, M. (2004) The modelling of operational risk: experience with the analysis of the data collected by the Basel Committee. Bank of Italy, Working Paper No 517.
- [27] Nelsen, R. B. (2007) *An Introduction to Copulas*. Springer, New York.
- [28] Nešlehová, J., Embrechts, P. and Chavez-Demoulin, V. (2006) Infinite mean models and the LDA for operational risk. *Journal of Operational Risk* **1**(1), 3–25.
- [29] Nešlehová, J. and McNeil, A. J. (2008) Multivariate Archimedean copulas,  $d$ -monotone functions and  $l_1$ -norm symmetric distributions. *Annals of Statistics*. To appear.

- [30] Resnick, S. I. (1987) *Extreme Values, Regular Variation, and Point Processes*. Springer, New York.
- [31] Resnick, S. I. (2004) The extremal dependence measure and asymptotic independence. *Stochastic Models* **20**, 205–227.
- [32] Resnick, S. I. (2007) *Heavy-Tail Phenomena: Probabilistic and Statistical Modeling*. Springer, New York.
- [33] Straetmans, S. (1998) *Extreme financial returns and their comovements*. Tinbergen Institute Research Series nr. 181, Thesis Publishers Amsterdam.
- [34] Strassburger, D. and Pfeifer, D. (2005) Dependence matters. 36th International Astin Colloquium, ETH Zurich.
- [35] Stărică, C. (1999) Multivariate extremes for models with constant conditional correlations. *Journal of Empirical Finance* **6**, 515–553.
- [36] Zhang, M.-H. (2008) Modelling total tail dependence along diagonals. *Insurance: Mathematics and Economics* **42**, 73–80.