

Closed Chains of Conics Carrying Poncelet Triangles

Lorenz Halbeisen

Department of Mathematics, ETH Zentrum, Rämistrasse 101, 8092 Zürich, Switzerland
lorenz.halbeisen@math.ethz.ch

Norbert Hungerbühler

Department of Mathematics, ETH Zentrum, Rämistrasse 101, 8092 Zürich, Switzerland
norbert.hungerbuehler@math.ethz.ch

key-words: conic sections, Poncelet theorem, conjugate conics, projective maps

2010 Mathematics Subject Classification: **51A05** 51A10 51A20

Abstract

We investigate closed chains of conics which carry Poncelet triangles. In particular, we show that every chain of conics which carries Poncelet triangles can be closed. Furthermore, for $k = 3$ and $k = 4$ we show that there are closed chains of pairwise conjugate conics which carry Poncelet k -gons such that the contact points of each k -gon are the vertices of the next k -gon—such miraculous chains of conics do not exist for $5 \leq k \leq 23$.

1 Introduction

In 1813, while Poncelet was in captivity as war prisoner in the Russian city of Saratov, he discovered his famous closing theorem which, in its simplest form, reads as follows (see [21]): *Let K and C be two conics in general position. Suppose there is a k -sided polygon inscribed in K and circumscribed about C . Then for any point P on K which is an exterior point of C , there exists a (possibly degenerate) k -sided polygon, also inscribed in K and circumscribed about C , which has P as one of its vertices.* See for example Dragović and Radnović [10] or Flatto [12] for classical overviews about Poncelet's Theorem, or Halbeisen and Hungerbühler [14] for a new elementary proof based only on Pascal's Theorem.

Recent years have seen a flourishing revitalization of Poncelet's closure theorem. We just mention some of these developments: Bos, Kers, Oort, and Raven [2] gave a comprehensive overview reaching from the pre-history of Poncelet's Porism to the modern approaches using elliptic curves. In their paper they use the language of algebraic geometry to analyze steps in the proofs of Poncelet and Jacobi with the help of modern notation and methods. Del Centina [8, 9] comments on the full range of the historical genesis of Poncelet's Theorem, its proofs, variants and relatives, and puts the results in a modern context. He does not miss to mention lesser known aspects like Nicola Trudi's approach or George Halphen's and Francesco Gerbaldi's discovery of a relation to continued fractions, or the applications of Poncelet's closure theorem and its generalizations to the theory of integrable system,

billiard dynamics, PDEs and statistical mechanics. In the further development of Poncelet's Porism, even very classical results came to new life: Chapple [5] found a relation for the radii and the distance of the centers of two nested circles which carry Poncelet triangles. This was generalized to Poncelet n -gons without self-intersections for small values of n by Euler [11], Fuss [13] and Steiner [23, p.96], [24, p.289]. These results have recently been extended by Cieřlak, Martini, Mozgawa [6] who formulated a relation for 5-gons with self-intersections and found a link to the rotation index of bar billiards and special functions leading to a new series expansion of π . Another approach to Poncelet's Porism uses measure theory: King [17] observed that there exists a measure on the conic which carries the vertices, which is invariant under tangential projection with respect to the inscribed conic. This easily proves Poncelet's Theorem. A similar construction has been proposed by Bryant [3], and Cieřlak and Mozgawa [7] found a nice geometric interpretation of the corresponding measure on the inscribed conic. Vallès [26] devised a new proof of Poncelet's Porism and of the related Darboux Theorem based upon vector bundles by exploiting a link between Schwarzenberger bundles and Poncelet curves.

In the present paper we investigate chains of conics G_0, \dots, G_{n-1} such that for each $0 \leq i \leq n-2$ there is a *Poncelet triangle* Δ_i with vertices on G_i whose sides are tangent to G_{i+1} . It will be shown in Section 4 that each such chain can be closed by adding a suitable conic G_n , i.e., there is a Poncelet triangle Δ_{n-1} with vertices on G_{n-1} and sides tangent to G_n , and G_n carries the vertices of a Poncelet triangle Δ_n whose sides are tangent to G_0 . The crucial point in the proof will be, that in specific cases, two conics can be projectively mapped simultaneously into two circles. This technical point is of some interest in its own right and will be investigated in Section 3.

We then show in Section 5 that there are closed chains of arbitrary length of conics G_0, \dots, G_{n-1} carrying Poncelet triangles, such that the contact points of the Poncelet triangle Δ_i (having its vertices on G_i) are the vertices of the Poncelet triangle Δ_{i+1} (having its vertices on G_{i+1} , where indices are taken modulo n).

Finally, in Section 6 we investigate two very peculiar chains of conics carrying Poncelet triangles or Poncelet quadrilaterals, respectively, which move synchronously while keeping contact with the neighbouring polygons. We also show that, up to projective transformations, these two chains are unique. Moreover, we show that there are no such chains of conics carrying k -gons for $5 \leq k \leq 23$.

2 Preliminary

In order to make this text self-contained, we start with a brief description of the general setting and establish the notation. An extensive survey about algebraic representations of conics in the real projective plane can be found in [1] or [18].

2.1 Projective plane and conics

In this paper, we mostly work in the standard model of the real projective plane. For this, we consider \mathbb{R}^3 and its dual space $(\mathbb{R}^3)^*$ of linear functionals on \mathbb{R}^3 . The set of points is

$\mathbb{P} = \mathbb{R}^3 \setminus \{0\} / \sim$, where $x \sim y \in \mathbb{R}^3 \setminus \{0\}$ are equivalent, if $x = \lambda y$ for some $\lambda \in \mathbb{R}$. The set of lines is $\mathbb{B} = (\mathbb{R}^3)^* \setminus \{0\} / \sim$, where $g \sim h \in (\mathbb{R}^3)^* \setminus \{0\}$ are equivalent, if $g = \lambda h$ for some $\lambda \in \mathbb{R}$. Finally, we say a point $[x]$ and a line $[g]$ are incident if $g(x) = 0$, where we denoted equivalence classes by square brackets. In the sequel we will identify \mathbb{R}^3 and $(\mathbb{R}^3)^*$ by the standard inner product $\langle \cdot, \cdot \rangle$ which allows to express the incidence through the relation $\langle x, g \rangle = 0$.

As usual, a line $[g]$ can be identified with the set of points which are incident with it. Vice versa a point $[x]$ can be identified with the set of lines which pass through it. The affine plane \mathbb{R}^2 is embedded in the present model of the projective plane by the map

$$\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \mapsto \left[\begin{pmatrix} x_1 \\ x_2 \\ 1 \end{pmatrix} \right].$$

The projective general linear group $PGL(3, \mathbb{R})$ consists of equivalence classes $[A]$ of regular matrices $A \in \mathbb{R}^{3 \times 3}$ representing maps $\mathbb{P} \rightarrow \mathbb{P}, [x] \mapsto [Ax]$, where two matrices are equivalent, $A_1 \sim A_2$, if $A_1 = \lambda A_2$ for some $\lambda \in \mathbb{R}$.

A *conic* in the constructed model is an equivalence class of a regular, linear, selfadjoint map $A : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ with mixed signature, i.e., A has eigenvalues of both signs. It is convenient to say, a matrix A is a conic, instead of A is a representative of a conic. We may identify a conic by the set of points $[x]$ such that $\langle x, Ax \rangle = 0$, or by the set of lines $[g]$ for which $\langle A^{-1}g, g \rangle = 0$ (see below). Notice that, in this interpretation, a conic cannot be empty: Since A has positive and negative eigenvalues, there are points $[p], [q]$ with $\langle p, Ap \rangle > 0$ and $\langle q, Aq \rangle < 0$. Hence a continuity argument guarantees the existence of points $[x]$ satisfying $\langle x, Ax \rangle = 0$.

From now on, we will only distinguish in the notation between an equivalence class and a representative if necessary.

FACT 2.1. *Let x be a point on the conic A . Then the line Ax is tangent to the conic A with contact point x .*

Proof. We show that the line Ax meets the conic A only in x . Suppose otherwise, that $y \not\sim x$ is a point on the conic, i.e., $\langle y, Ay \rangle = 0$, and at the same time on the line Ax , i.e., $\langle y, Ax \rangle = 0$. By assumption, we have $\langle x, Ax \rangle = 0$. Note, that $Ax \not\sim Ay$ since A is regular, and $\langle Ay, x \rangle = 0$ since A is selfadjoint. Hence x and y both are perpendicular to the plane spanned by Ax and Ay , which contradicts $y \not\sim x$. *q.e.d.*

In other words, the set of tangents of a conic A is the image of the points on the conic under the map A . And consequently, a line g is a tangent of the conic iff $A^{-1}g$ is a point on the conic, i.e., if and only if $\langle A^{-1}g, g \rangle = 0$.

DEFINITION 2.2. *If P is a point, the line AP is called its polar with respect to a conic A . If g is a line, the point $A^{-1}g$ is called its pole with respect to the conic A .*

Obviously, the pole of the polar of a point P is again P , and the polar of the pole of a line g is again g . Moreover:

FACT 2.3. *If the polar of point P with respect to a conic A intersects the conic in a point x , then the tangent in x passes through P .*

Proof. For x , we have $\langle x, Ax \rangle = 0$ since x is a point on the conic, and $\langle x, AP \rangle = 0$ since x is a point on the polar of P . The tangent in x is the line Ax , and indeed, P lies on this line, since $\langle P, Ax \rangle = \langle AP, x \rangle = 0$. q.e.d.

2.2 Coordinate transformations and projective maps

An element $T \in PGL(3, \mathbb{R})$ can be interpreted as a change of coordinates $x = Ty$: If for example $\langle x, Ax \rangle = 0$ is a conic in x -coordinates, then $0 = \langle x, Ax \rangle = \langle Ty, ATy \rangle = \langle y, T^\top ATy \rangle$ is the same conic in y -coordinates (i.e., the transformed conic is represented by the matrix $T^\top AT$). Similarly, if $\langle g, x \rangle = 0$ is a line, then $0 = \langle g, x \rangle = \langle T^\top g, y \rangle$, i.e., the transformed line is represented by $T^\top g$.

Instead of considering T as a coordinate transformation, we can equivalently interpret $T : \mathbb{P} \rightarrow \mathbb{P}$ as a projective map: Then, if y is a point on the conic $T^\top AT$, i.e., if $\langle y, T^\top ATy \rangle = 0$, then the image $x := Ty$ is a point of the conic $\langle x, Ax \rangle = 0$.

2.3 Rank one and rank two matrices

For completeness we recall the following basic facts:

LEMMA 2.4. *Let $A \in \mathbb{R}^{n \times n}$ be symmetric. Then:*

- *A is of rank 1 iff $A = \alpha aa^\top$ for a unit vector $a \in \mathbb{R}^n$ and some $\alpha \in \mathbb{R} \setminus \{0\}$. Moreover, if $A = \alpha aa^\top$ then, the vector a is an eigenvector and α the corresponding eigenvalue of A , and $\langle x, Ax \rangle = \alpha \langle a, x \rangle^2$.*
- *A is of rank 2 iff $A = \alpha aa^\top + \beta bb^\top$ for two orthonormal vectors $a, b \in \mathbb{R}^n$ and some $\alpha, \beta \in \mathbb{R} \setminus \{0\}$. Moreover, if $A = \alpha aa^\top + \beta bb^\top$ then, a and b are eigenvectors and α, β the corresponding eigenvalues of A . Moreover, if A is of rank 2, $A = uv^\top + vu^\top$ for two vectors $u, v \in \mathbb{R}^n \setminus \{0\}$ iff A has mixed signature. Finally, if $A = \alpha aa^\top + \beta bb^\top$, then $\langle x, Ax \rangle = \alpha \langle x, a \rangle^2 + \beta \langle x, b \rangle^2$, and if $A = uv^\top + vu^\top$, then $\langle x, Ax \rangle = 2 \langle x, u \rangle \langle x, v \rangle$.*

Proof. Since A is symmetric there exists an orthonormal eigenbasis $t^{(1)}, \dots, t^{(n)}$, i.e., if T is the matrix with columns $t^{(1)}, \dots, t^{(n)}$, then $T^\top AT = \text{diag}(\lambda_1, \dots, \lambda_n) =: D$, where λ_i is the eigenvalue corresponding to $t^{(i)}$. Therefore $A = TDT^\top$. If A is of rank 1, exactly one eigenvalue $\lambda_i \neq 0$, and hence $A = \lambda_i t^{(i)} t^{(i)\top}$. If A is of rank 2, exactly two eigenvalues $\lambda_i, \lambda_j \neq 0$, and hence $A = \lambda_i t^{(i)} t^{(i)\top} + \lambda_j t^{(j)} t^{(j)\top}$.

On the other hand, if $A = \alpha aa^\top$ for a unit vector a and $\alpha \in \mathbb{R} \setminus \{0\}$, it is clear that $A = A^\top$, that all columns of A are multiples of a and hence that A is of rank 1. Moreover, $Aa = \alpha aa^\top a = \alpha a$. If $A = \alpha aa^\top + \beta bb^\top$ for orthonormal vectors a, b and $\alpha, \beta \in \mathbb{R} \setminus \{0\}$, then $A = A^\top$, $Aa = (\alpha aa^\top + \beta bb^\top)a = \alpha a$ and $Ab = (\alpha aa^\top + \beta bb^\top)b = \beta b$. Hence, A

has two orthonormal eigenvectors and therefore $\text{rank } A \geq 2$. On the other hand, $Ax = 0$ whenever $x \perp \text{span}(a, b)$, i.e., $\text{rank } A = 2$.

Finally, let A be of rank 2, i.e., $A = \alpha aa^\top + \beta bb^\top$, and suppose $\alpha > 0 > \beta$. Then, for $u = \mu a + \nu b, v = \mu a - \nu b$ we have $A = uv^\top + vu^\top$ if we choose $\mu = \sqrt{\alpha/2}, \nu = \sqrt{-\beta/2}$. On the other hand, if $A = uv^\top + vu^\top$ for two non-zero vectors $u \neq v \in \mathbb{R}^n$, then A has the eigenvalues

$$0, \langle u, v \rangle + \|u\|\|v\|, \langle u, v \rangle - \|u\|\|v\|.$$

Thus, by the Cauchy-Schwarz inequality, the two non-zero eigenvalues have opposite sign.

The formulas for the quadratic forms follow immediately.

q.e.d.

LEMMA 2.5. *Let A be a conic and $C \in \mathbb{R}^{3 \times 3}$ symmetric of rank 1. Suppose, $A + C$ is also a conic, and p a point on A and on $A + C$. Then, A and $A + C$ have a common tangent in p .*

Proof. According to Lemma 2.4, C is of the form γcc^\top for some $c \in \mathbb{R}^3 \setminus \{0\}$, $\gamma \neq 0$. We have

$$0 = \langle p, (A + C)p \rangle = \langle p, Ap \rangle + \langle p, Cp \rangle = \langle p, Cp \rangle = \gamma \langle c, p \rangle^2,$$

hence $\langle c, p \rangle = 0$. The tangent in p at A is given by $0 = \langle x, Ap \rangle$. The tangent in p at $A + C$ is given by the same equation: $0 = \langle x, (A + C)p \rangle = \langle x, Ap + \gamma cc^\top p \rangle = \langle x, Ap \rangle$. *q.e.d.*

2.4 Projective maps leaving a circle invariant

Let $K = \text{diag}(1, 1, -1)$ be the affine unit circle and \mathfrak{G}_K the subgroup of all projective maps which leave K invariant, i.e.,

$$\mathfrak{G}_K := \{T \in PGL(3, \mathbb{R}) \mid T^\top K T \sim K\}.$$

LEMMA 2.6. *If $T \in \mathfrak{G}_K$ has the fixed points $(0, 0, 1)^\top$ and $(1, 0, 1)^\top$, then T is the identity or $T = \text{diag}(1, -1, 1) =: S$.*

Proof. Since $(0, 0, 1)^\top$ is a fixed point of T , we have

$$T = \begin{pmatrix} * & * & 0 \\ * & * & 0 \\ t_{31} & * & t_{33} \end{pmatrix}.$$

Then, because $(1, 0, 1)^\top$ is also a fixed point, it follows

$$T = \begin{pmatrix} t_{31} + t_{33} & * & 0 \\ 0 & * & 0 \\ t_{31} & * & t_{33} \end{pmatrix}$$

i.e., $t_{11} = t_{31} + t_{33}$. On the other hand, from $K \sim T^\top K T$, it now follows

$$K \sim \begin{pmatrix} * & * & -t_{31}t_{33} \\ * & * & -t_{32}t_{33} \\ * & * & -t_{33}^2 \end{pmatrix}$$

which implies $t_{31} = t_{32} = 0$. Using again $K \sim T^\top K T$ we find

$$K \sim \begin{pmatrix} * & t_{12}t_{33} & 0 \\ * & * & 0 \\ 0 & 0 & -t_{33}^2 \end{pmatrix}$$

from which we deduce $t_{12} = 0$. Then, finally,

$$K \sim \begin{pmatrix} t_{33}^2 & 0 & 0 \\ 0 & t_{22}^2 & 0 \\ 0 & 0 & -t_{33}^2 \end{pmatrix}.$$

Hence $t_{22} = \pm 1$ and w.l.o.g. we may choose $t_{33} = 1$.

q.e.d.

LEMMA 2.7. *If $T \in \mathfrak{G}_K$ has the fixed point $(0, 0, 1)^\top$ and*

$$T(1, 0, 1)^\top = (\cos \phi, \sin \phi, 1)^\top,$$

then $T = D_\phi$ or $T = D_\phi S =: S_\phi$, where $S = \text{diag}(1, -1, 1)$ and

$$D_\phi := \begin{pmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Proof. Because of Lemma 2.6, we have $D_\phi^{-1}T = \mathbb{I}$ or $D_\phi^{-1}T = S$.

q.e.d.

LEMMA 2.8. *If $T \in \mathfrak{G}_K$ has the fixed point $(0, 0, 1)^\top$ and $T(\cos \phi, \sin \phi, 1)^\top = (\cos \psi, \sin \psi, 1)^\top$, then $T = D_\psi D_\phi^{-1} = D_{\psi-\phi}$ or $T = S_\psi D_\phi^{-1} = S_{\psi+\phi}$.*

Proof. The projective map $TD_\phi \in \mathfrak{G}_K$ has the fixed point $(0, 0, 1)^\top$, and $TD_\phi(1, 0, 1)^\top = (\cos \psi, \sin \psi, 1)^\top$. Therefore, by Lemma 2.7, $TD_\phi = D_\psi$ or $TD_\phi = S_\psi$. *q.e.d.*

Summarizing, we have the following: If $T \neq \text{id}$ is a projective map with fixed point $(0, 0, 1)^\top$ which maps the unit circle K to itself, then T is either a rotation D_η (namely if $(0, 0, 1)^\top$ is the only fixed point) or T is a reflection S_η . In particular: If $T \in \mathfrak{G}_K$ has the fixed points $(0, 0, 1)^\top$ and $(1, 0, 1)^\top$, then T is either the identity or the reflection with respect to the line $(0, 1, 0)^\top$.

A conic C divides the points \mathbb{P} of the projective plane into three disjoint sets: The points on C (for those points there is exactly one tangent to C), the exterior points (for those points there are two tangents to C) and the inner points (which are not incident with a tangent of C). Observe, that this classification is invariant under projective maps. In particular, if $T \in \mathfrak{G}_K$ and if z is an inner point of K , then Tz is also inner point of K . For example, with

$$U_\psi := \begin{pmatrix} \cosh \psi & 0 & \sinh \psi \\ 0 & 1 & 0 \\ \sinh \psi & 0 & \cosh \psi \end{pmatrix} \in \mathfrak{G}_K,$$

we have $U_\psi(0, 0, 1)^\top \sim (\tanh \psi, 0, 1)^\top$.

Let $T \in \mathfrak{G}_K$ and $T(0, 0, 1)^\top = (\tanh \psi \cos \phi, \tanh \psi \sin \phi, 1)^\top$ be an arbitrary inner point of K . Then $U_\psi^{-1}D_\phi^{-1}T \in \mathfrak{G}_K$ has the fixed point $(0, 0, 1)^\top$ and is hence either a rotation D_η or a reflection S_η . Therefore, $T = D_\phi U_\psi D_\eta$ or $T = D_\phi U_\psi S_\eta$. Thus, we have:

THEOREM 2.9. *The group \mathfrak{G}_K is generated by*

$$\begin{pmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} \cosh \psi & 0 & \sinh \psi \\ 0 & 1 & 0 \\ \sinh \psi & 0 & \cosh \psi \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

where $\phi \in [0, 2\pi[$, $\psi \in]-\infty, \infty[$.

For later use we add the following two lemmata:

LEMMA 2.10. *Let $K = \text{diag}(1, 1, -1)$ be the affine unit circle and g a line which misses K . Then, g can be written as $g = (\tanh \psi \cos \phi, \tanh \psi \sin \phi, 1)^\top$ for some $\psi \geq 0$, $\phi \in [0, 2\pi[$, and $T = U_{-\psi}^{-1}D_\phi^{-1} \in \mathfrak{G}_K$ maps g to the ideal line $(0, 0, 1)$.*

Proof. The pole $P = K^{-1}g \sim (\tanh(-\psi) \cos \phi, \tanh(-\psi) \sin \phi, 1)^\top$ of g with respect to K is an inner point of K and $T = U_{-\psi}^{-1}D_\phi^{-1} \in \mathfrak{G}_K$ maps P to the origin, and hence g to the ideal line. q.e.d.

LEMMA 2.11. *Let P_1, P_2, P_3 and Q_1, Q_2, Q_3 be two triples of points on the affine unit circle K . Then, there is a projective map $T \in \mathfrak{G}_K$ which maps P_i to Q_i , $i \in \{1, 2, 3\}$.*

Proof. It suffices to show the lemma for $Q_1 = (1, 0, 1)^\top$, $Q_2 = (-1, 0, 1)^\top$, $Q_3 = (0, 1, 1)^\top$. By using a suitable rotation D_ϕ we may assume, that P_1 and P_2 have the same x_1 component: $P_{1,2} = (\cos \phi, \pm \sin \phi, 1)^\top$. Then, U_ψ maps $P_{1,2}$ to the points $(0, \pm 1, 1)^\top$ for $\psi = -\text{artanh} \cos \phi$. Another rotation maps these points to $Q_1 = (1, 0, 1)^\top$ and $Q_3 = (-1, 0, 1)^\top$, and the image of P_3 under the previous operations is some point $(\cos \eta, \sin \eta, 1)^\top$. Then, a suitable map U_ξ maps this point to either $(0, 1, 1)^\top$ or $(0, -1, 1)^\top$ and leaves the points Q_1 and Q_2 invariant. If necessary, a reflection S with respect to the x_1 axis completes the proof. q.e.d.

The circle $(x_1 - a_1 x_3)^2 + (x_2 - a_2 x_3)^2 - r^3 x_3^2 = 0$ with radius r and center (a_1, a_2) in the affine plane corresponds to the matrix

$$\begin{pmatrix} 1 & 0 & -a_1 \\ 0 & 1 & -a_2 \\ -a_1 & -a_2 & a_1^2 + a_2^2 - r^2 \end{pmatrix}.$$

Vice versa, the conic

$$\begin{pmatrix} 1 & 0 & \alpha \\ 0 & 1 & \beta \\ \alpha & \beta & \gamma \end{pmatrix}$$

corresponds to a circle with center $(-\alpha, -\beta)$ and radius $r^2 = \alpha^2 + \beta^2 - \gamma$ if this number is strictly positive.

For two different circles

$$K_i = \begin{pmatrix} 1 & 0 & -a_i \\ 0 & 1 & -b_i \\ -a_i & -b_i & a_i^2 + b_i^2 - r_i^2 \end{pmatrix},$$

$i \in \{1, 2\}$, the difference

$$K_1 - K_2 = \begin{pmatrix} 0 & 0 & a_2 - a_1 \\ 0 & 0 & b_2 - b_1 \\ a_2 - a_1 & b_2 - b_1 & a_2^2 - a_1^2 + b_2^2 - b_1^2 + r_2^2 - r_1^2 \end{pmatrix}$$

is a matrix of rank < 3 and represents two lines

$$\langle x, (K_1 - K_2)x \rangle = \left\langle \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, x \right\rangle \left\langle \begin{pmatrix} 2(a_2 - a_1) \\ 2(b_2 - b_1) \\ a_1^2 - a_2^2 + b_1^2 - b_2^2 + r_2^2 - r_1^2 \end{pmatrix}, x \right\rangle.$$

The line that corresponds to the first factor is the ideal line, the second line is the radical axis of the two circles (which is different from the ideal line if K_1 and K_2 have different centers). Vice versa, we have:

THEOREM 2.12. *Let K be the affine unit circle*

$$K = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix},$$

and A a conic. Suppose, the pencil $K + \lambda A$ of K and A contains the ideal line $(0, 0, 1)^\top$, then A is a circle.

Proof. Let $e = (0, 0, 1)^\top$. Then, for certain $\lambda, \mu \in \mathbb{R}$ and a line g , the quadratic form

$$\langle x, (K + \lambda A)x \rangle = 2\mu \langle e, x \rangle \langle g, x \rangle = 2\mu x_3 (g_1 x_1 + g_2 x_2 + g_3 x_3) \quad (1)$$

is represented by

$$K + \lambda A = \mu \begin{pmatrix} 0 & 0 & g_1 \\ 0 & 0 & g_2 \\ g_1 & g_2 & 2g_3 \end{pmatrix}.$$

Obviously, $\lambda \neq 0$ and

$$A \sim \begin{pmatrix} 1 & 0 & -\mu g_1 \\ 0 & 1 & -\mu g_2 \\ -\mu g_1 & -\mu g_2 & -1 - 2\mu g_3 \end{pmatrix}.$$

which is a circle in the affine plane with radius $r^2 = 1 + 2\mu g_3 + \mu^2(g_1^2 + g_2^2)$ and center $\mu(g_1, g_2)$. *q.e.d.*

As a consequence of Lemma 2.10 and Theorem 2.12 we get:

COROLLARY 2.13. *Let K be the unit circle and A a conic such that the pencil of K and A contains a line g which misses both K and A . Then, the projective map T which leaves K invariant and sends g to the ideal line maps A to a circle.*

Proof. According to Lemma 2.10 there exists a map $T \in \mathfrak{G}_K$ which sends g to the ideal line. Observe, that T maps the pencil of K and A to the pencil of their images, K and $\tilde{T} := T^{-\top}AT^{-1}$. Therefore, the ideal line belongs to the pencil of K and \tilde{T} and thus, according to Theorem 2.12, \tilde{T} is a circle. q.e.d.

3 How to transform two conics into two circles

Let us first recall, how one can find coordinates for which *one* given conic is a circle. For this, consider a conic $A \in \mathbb{R}^{3 \times 3}$, symmetric, regular and with mixed signature. The signature of A can be determined from the characteristic polynomial:

LEMMA 3.1. *Let $A \in \mathbb{R}^{3 \times 3}$ be regular and symmetric with characteristic polynomial $p_A(\lambda) = \det(A - \lambda I) = a_0 + a_1\lambda + a_2\lambda^2 - \lambda^3$. Then the following holds:*

- *If all eigenvalues of A are positive, we have $a_0 > 0, a_1 < 0, a_2 > 0$.*
- *If all eigenvalues of A are negative, we have $a_0 < 0, a_1 < 0, a_2 < 0$.*
- *If A has two negative and one positive eigenvalue, then $a_0 > 0$, and if $a_1 < 0$ then $a_2 < 0$.*
- *If A has two positive and one negative eigenvalue, then $a_0 < 0$, and if $a_1 < 0$ then $a_2 > 0$.*

Proof. Since A is symmetric, all eigenvalues $\lambda_1, \lambda_2, \lambda_3$ are real and $p_A(\lambda) = (\lambda_1 - \lambda)(\lambda_2 - \lambda)(\lambda_3 - \lambda)$. Hence, $a_0 = \lambda_1\lambda_2\lambda_3 \neq 0$, $a_1 = -(\lambda_1\lambda_2 + \lambda_1\lambda_3 + \lambda_2\lambda_3)$ and $a_2 = \lambda_1 + \lambda_2 + \lambda_3$. It is then easy to check the four cases. q.e.d.

The transformation of A to a unit circle is then as follows:


LEMMA 3.2. *Let $A \in \mathbb{R}^{3 \times 3}$ be a conic. Without loss of generality, A has two positive and one negative eigenvalue (otherwise take the representative $-A$). Then there is a regular matrix T such that $T^\top AT = \text{diag}(1, 1, -1)$. This corresponds to the unit circle in the affine plane.*

Proof. Let $\lambda_1, \lambda_2 > 0$ be the positive eigenvalues, $\lambda_3 < 0$ the negative eigenvalue, and x_1, x_2, x_3 be the corresponding orthonormal eigenvectors. Then the matrix T with columns $\frac{1}{\sqrt{\lambda_1}}x_1, \frac{1}{\sqrt{\lambda_2}}x_2, \frac{1}{\sqrt{-\lambda_3}}x_3$ has the desired property. q.e.d.


In his seminal work [21, Section I, Chapitre III, no. 121, p. 59] Poncelet claims that two conics can, in general, be considered as projective image of two circles. Actually, referring

to this statement, he reduces the proof of Poncelet's Porism to the case of a pencil of circles in [21, Section IV, Chapitre II, no. 530, p. 311 ff.] The question whether two conics are the projective image of two circles also occurs in [22] as a proposed problem, with alleged solution in [25]. However, it is clear that for example two conics which intersect in four points *cannot* be the projective image of two circles.


It turns out that two conics can lie in 8 different positions relative to each other (see [20]):




Case 1: four intersections




Case 2: no intersections




Case 3: two intersections




Case 4: two intersections,
one 1st order contact




Case 5: one 1st order contact



Case 6: two 1st order
contacts



Case 7: one intersection, one
2nd order contact



Case 8: one 3rd order contact

Projective transformations do not change the number of intersections nor the order of contact, hence the cases 1, 4, 6, 7 and 8 cannot be the projective image of two circles. In fact we have:

THEOREM 3.3. *Two conics are the projective image of two circles if and only if they (i) intersect in two points, (ii) have one 1st order contact, or (iii) are disjoint.*

Proof. By the remark above, we only need to show one implication. Without loss of generality, we may assume that one of the conics is the circle $A = \text{diag}(1, 1, -1)$. $B \in \mathbb{R}^{3 \times 3}$ is an arbitrary conic different from A . According to Corollary 2.13, it suffices to show in each of the three cases (i)–(iii) that the pencil of A and B contains a line that misses both conics.

(i) Suppose A and B intersect in two points z_1, z_2 . The line $g = z_1 \times z_2$ is the common secant of A and B . Let p be a point on g different from z_1, z_2 and

$$r = -\frac{\langle p, Ap \rangle}{\langle p, Bp \rangle} \neq 0. \quad (2)$$

Then, z_1, z_2 and p are solutions of $\langle x, (A + rB)x \rangle = 0$. Since z_1, z_2 and p are collinear, we conclude $\text{rank}(A + rB) < 3$. On the other hand, A and B are different conics, hence $\text{rank}(A + rB) > 0$. Moreover, $\text{rank}(A + rB) \neq 1$, because otherwise, $-A$ and $-A + (A + rB) = rB$ would be tangential in z_1 and z_2 (see Lemma 2.5). Thus, $\text{rank} A = 2$ and by Lemma 2.4, we have $A + rB = \alpha aa^\top + \beta bb^\top$ for two orthonormal vectors a, b and $\alpha, \beta \in \mathbb{R}^3 \setminus \{0\}$. If the eigenvalues α, β would both be positive or both be negative, $\langle x, (A + rB)x \rangle = 0$ would have only one solution (see Lemma 2.4), but we have at least three, namely z_1, z_2, p . Therefore, $A + rB$ has mixed signature and hence, according to Lemma 2.4, $A + rB = uv^\top + vu^\top$ for two non-zero vectors $u \neq v \in \mathbb{R}^3$, and $\langle x, (A + rB)x \rangle = 2\langle x, u \rangle \langle x, v \rangle = 0$ consists of two lines: One of them is $g = u$, the other, $v \neq u$, has no common point with the conics A and B : Indeed, suppose we have a point p on v and A or B , then p would solve $\langle p, Ap \rangle = 0$ and $\langle p, Bp \rangle = 0$. Hence, w.l.o.g., $p = z_1$. But then, v cuts A or B in a second point, which then is necessarily z_2 . This contradicts $u \neq v$.

(ii) Suppose, A and B have one contact point, say z_0 , in common. Let $p \neq z_0$ be a point on the tangent g in z_0 and r as above in (2). Then, p and z_0 are solutions of $\langle x, (A + rB)x \rangle = 0$. Suppose $A + rB$ has full rank. Then, the tangent g in z_0 at the conic $A + rB$ does not contain another point of that same conic. But this is not true, since p is a point of g . Hence, $\text{rank}(A + rB) < 3$. Since, by assumption, A and B are different conics, $\text{rank}(A + rB) > 0$. Suppose, $\text{rank}(A + rB) = 1$. We may assume, that $z_0 = (0, 1, 1)^\top$ and thus $g = (0, -1, 1)^\top$. According to Lemma 2.4 $A + rB \sim aa^\top$ for a unit vector a . Hence, $g \sim (A + rB)z_0 \sim aa^\top z_0 \sim a$, and we conclude

$$B \sim A + \mu gg^\top \sim A + \mu \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & -1 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 0 \\ 0 & \mu + 1 & -\mu \\ 0 & -\mu & \mu - 1 \end{pmatrix}$$

and $\langle x, Bx \rangle = x_1^2 + x_2^2(1 + \mu) - x_3^2(1 - \mu) - 2x_2x_3\mu$. It is then easy to check, that A and B have a 3rd order contact in z_0 , which, by assumption, is not the case. Therefore, $\text{rank}(A + rB) = 2$, and, by the same reasoning as above in (i), we conclude, that $\langle x, (A + rB)x \rangle = 2\langle x, g \rangle \langle x, v \rangle = 0$ consists of two lines, $g \neq v$, and that v has no common point with A or B .

(iii) Suppose, A and B are disjoint. Then, there exist coordinates for which both conics are diagonal (see for example [19] or [16]): W.l.o.g.

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & b \end{pmatrix}$$

where $a, b \neq 0$ are not both positive. Then, for the values $r \in \{-1, -\frac{1}{a}, \frac{1}{b}\}$, we have for $A + rB$:

$$A - B = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 - a & 0 \\ 0 & 0 & -1 - b \end{pmatrix}, \quad A - \frac{1}{a}B = \begin{pmatrix} 1 - \frac{1}{a} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 - \frac{b}{a} \end{pmatrix},$$

$$A + \frac{1}{b}B = \begin{pmatrix} 1 + \frac{1}{b} & 0 & 0 \\ 0 & 1 + \frac{a}{b} & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

For all values of a and b one of these three matrices has mixed signature and is of rank 2. The corresponding quadratic form $\langle x, (A + rB)x \rangle = 2\langle x, g \rangle \langle x, h \rangle = 0$ represents two lines which both miss A and B . *q.e.d.*

4 Weakly connected closed chains of Poncelet triangles

Let us first recall the Cayley criterion for Poncelet polygons:

THEOREM 4.1 (Cayley criterion, see [4]). *Let A, H be conics, $D(\lambda) = \det(A + \lambda H)$, and*

$$\sqrt{D(\lambda)} = c_0 + c_1\lambda + c_2\lambda^2 + c_3\lambda^3 + \dots$$

Then, there exists a Poncelet k -gon with vertices on H and tangent to A if and only if

$$\det \begin{pmatrix} c_3 & c_4 & \dots & c_{p+1} \\ c_4 & c_5 & \dots & c_{p+2} \\ \dots & & & \\ c_{p+1} & c_{p+2} & \dots & c_{2p-1} \end{pmatrix} = 0 \quad \text{for } k = 2p,$$

or

$$\det \begin{pmatrix} c_2 & c_3 & \dots & c_{p+1} \\ c_3 & c_4 & \dots & c_{p+2} \\ \dots & & & \\ c_{p+1} & c_{p+2} & \dots & c_{2p} \end{pmatrix} = 0 \quad \text{for } k = 2p + 1.$$

In this section, we will consider chains of conics, where each pair of consecutive conics is a Poncelet pair for triangles. The main result of this section is that every such chain can be closed. Let us start with the following definition:

A chain of conics G_0, G_1, \dots, G_n , $n \geq 1$, is called **weakly connected Poncelet chain for triangles** if, for $i \in \{0, 1, \dots, n-1\}$, (G_i, G_{i+1}) is a Poncelet pair for triangles, i.e., G_i carries vertices of triangles whose sides are tangential to G_{i+1} . The chain is called **closed** and of length n , if $G_n = G_0$.

The central lemma which allows to close a chain is the following:

LEMMA 4.2. *Let G and H be conics. Then, there exists a conic A such that (H, A) and (A, G) are Poncelet pairs for triangles: H carries vertices of triangles that are tangent to A , and A carries vertices of triangles that are tangent to G .*

We call A a **lock** between G and H .

The six coefficients of A have to satisfy two polynomial equations of order 2 and 4 respectively (the Cayley conditions), A needs to have mixed signature, and it has to have points outside of G and inside of H .

Proof of Lemma 4.2. We have to consider the 8 cases listed in Section 3.

Case 1. G and H have four intersections P_1, P_2, P_3, P_4 . By Lemma 3.2 we may assume, that G is the unit circle K . Let P be the intersection of the lines P_1P_3 and P_2P_4 joining opposite points. Then we choose $T \in \mathfrak{G}_K$ such that T maps P to $(0, 0, 1)^\top$ (see Theorem 2.9). Thus, the image of H is a conic with center $(0, 0, 1)^\top$, either a hyperbola or an ellipse. By a suitable rotation D_ϕ , we may further assume that H is symmetric with respect to the x_1 - and x_2 -axis.

Case 1a. A circle and a concentric hyperbola:

$$G = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \text{ and } H = \begin{pmatrix} a^2 & 0 & 0 \\ 0 & -b^2 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

We choose

$$A = \begin{pmatrix} 0 & x & x \\ x & 1 & -1 \\ x & -1 & 1 \end{pmatrix} \text{ with } x = \frac{4a}{\sqrt{b^2 + 1}}.$$

Then a direct computation shows that

$$\left. \frac{d^2}{d\lambda^2} \sqrt{\det(G + \lambda A)} \right|_{\lambda=0} = 0 \quad \text{and} \quad \left. \frac{d^2}{d\lambda^2} \sqrt{\det(A + \lambda H)} \right|_{\lambda=0} = 0 \quad (3)$$

which shows that the Cayley criterion in Theorem 4.1 for triangles is satisfied for both pairs H, A and A, G .

The eigenvalues of A are $\{2, \sqrt{2}x, -\sqrt{2}x\}$, i.e., A has mixed signature. As A is a hyperbola, it has points outside G . A passes through the point $(0, 1, 1)$ and has the asymptote $(0, 1, 1)^\top$. Therefore, the intersections of this asymptote with H are outer points of A .

Case 1b. A circle and a concentric ellipse:

$$G = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \text{ and } H = \begin{pmatrix} a^2 & 0 & 0 \\ 0 & b^2 & 0 \\ 0 & 0 & -1 \end{pmatrix}, b^2 \neq 1.$$


We choose

$$A = \begin{pmatrix} y & x & x \\ x & y & 0 \\ x & 0 & 0 \end{pmatrix}$$

with

$$x = \sqrt{1 + 2a^2 + b^2 + 2\sqrt{(1 + a^2)(a^2 + b^2)}}, \quad y = b^2 - 1.$$

Again, the Cayley criterion (3) for triangles is easily verified. By Lemma 3.1 it can be checked, that A has mixed signature. In fact, A is a hyperbola through $(0, 0, 1)^\top$ and has therefore points outside G , and H has points outside A .



Lock in Case 1a (left) and 1b (right): H carries vertices of Poncelet triangles that are tangent to A , A carries vertices of Poncelet triangles that are tangent to G .

Cases 2 and 3. G and H have either no common points or two intersections. By Theorem 3.3, we may assume, that G is the affine unit circle, and H another circle having its center on the x_1 -axis:

$$G = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \text{ and } H = \begin{pmatrix} 1 & 0 & -a \\ 0 & 1 & 0 \\ -a & 0 & a^2 - r^2 \end{pmatrix} \text{ with } a \geq 0.$$

We choose

$$A = \begin{pmatrix} 0 & 0 & x \\ 0 & 1 & 0 \\ x & 0 & y \end{pmatrix}$$

with

$$x = \sqrt{1 + (1 + a - r)^2} + r - 1 - a, \quad y = 2\sqrt{1 + (1 + a - r)^2} + 2r - 3 - 2a.$$

Then the Cayley criterion (3) for triangles is easily verified. A is a parabola symmetric to the x_1 -axis with vertex inside H and has therefore points outside G , and H has points outside A .

Case 4. G and H have two intersections and one first order contact. We may again assume that G is the affine unit circle $\text{diag}(1, 1, -1)$, and, by Lemma 2.11, that the first order contact is in $(1, 0, 1)^\top$ and that the two intersections are $(0, \pm 1, 1)^\top$. Then, the pencil of G and H contains two lines $(1, \pm 1, -1)^\top$, and according to Lemma 2.4

$$H = G + \mu \begin{pmatrix} 1 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 1 \end{pmatrix} = \begin{pmatrix} \mu + 1 & 0 & -\mu \\ 0 & 1 - \mu & 0 \\ -\mu & 0 & \mu - 1 \end{pmatrix}, \quad \mu \notin \{0, 1\}.$$

We choose

$$A = \begin{pmatrix} 0 & 0 & p \\ 0 & 1 & 0 \\ p & 0 & q \end{pmatrix}$$

with

$$p = \frac{2\mu - \sqrt{1 + 3\mu^2}}{\mu - 1}, \quad q = \frac{1 + 3\mu - 2\sqrt{1 + 3\mu^2}}{\mu - 1}.$$

A direct calculation shows that the Cayley criterion (3) is satisfied. A is a parabola which is symmetric to the x_1 -axis. Its vertex $(x_1, 0, 1)^\top$ satisfies $x_1 < 1$. In particular, A has points outside G . H is also symmetric to the x_1 -axis: For $\mu < -1$ and $\mu > 1$ it is a hyperbola, for $-1 < \mu < 1$ an ellipse, and for $\mu = -1$ a parabola. In each case, H has points outside A .

Case 5. G and H have one first order contact: See Cases 2 and 3.

Case 6. G and H have two first order contacts. As above, we may assume, that G is the affine unit circle $\text{diag}(1, 1, -1)$, and, by Lemma 2.11, that the contact points are $(\pm 1, 0, 1)^\top$. The pencil of G and H contains the double line $(0, 1, 0)^\top$ and according to Lemma 2.4:

$$H = G - \mu \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 - \mu & 0 \\ 0 & 0 & -1 \end{pmatrix}, \mu \neq 0.$$

If $\mu > 0$, we choose

$$A = \begin{pmatrix} 0 & q & q \\ q & 1 & -1 \\ q & -1 & 1 \end{pmatrix} \text{ with } q = \frac{4}{\sqrt{\mu}}.$$

If $\mu < 0$, we choose

$$A = \begin{pmatrix} p & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 2 - q \end{pmatrix} \text{ with } p = 1 - \frac{1}{\sqrt{1 - \mu}} + \sqrt{\frac{\mu - 3 + 2\sqrt{1 - \mu}}{\mu - 1}}.$$

In both cases, it is easy to check, that the Cayley criterion (3) is satisfied, that A has mixed signature and has points outside G . Moreover, H has points outside A .

Case 7. G and H have one intersection and one first order contact. We may again assume, that G is the affine unit circle K . By Lemma 2.11 we may assume, that the first order contact is in $(1, 0, 1)^\top$ and the intersection in $(-1, 0, 1)^\top$. Then the lines $(-1, 0, 1)^\top$ and $(0, 1, 0)^\top$ belong to the pencil of G and H and according to Lemma 2.4:

$$H = K - \mu \begin{pmatrix} 0 & -1 & 0 \\ -1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & -\mu & 0 \\ -\mu & 1 & \mu \\ 0 & \mu & -1 \end{pmatrix}, \mu \neq 0.$$

Then, the choice

$$A = \begin{pmatrix} 0 & q & q \\ q & 1 & -1 \\ q & -1 & 1 \end{pmatrix} \text{ with } q = \frac{2}{\mu} \left(2 + \mu + \sqrt{4 - 2\mu + \mu^2} \right)$$

yields the desired lock.

Case 8. G and H have one third order contact. We may assume that G is the affine unit circle $(1, 1, -1)^\top$, and, by a suitable rotation, that the point of contact is $(0, 1, 1)^\top$. As discussed in the proof of Theorem 3.3 the pencil of G and H contains the double line $(0, -1, 1)^\top$. Thus, by Lemma 2.4,

$$H = G + \mu \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & -1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \mu + 1 & -\mu \\ 0 & -\mu & \mu - 1 \end{pmatrix}.$$

Here, if $\mu \neq -1$, we choose

$$A = \begin{pmatrix} 1 & 0 & a \\ 0 & -1 & b \\ a & b & 0 \end{pmatrix} \text{ with } a = \frac{\sqrt{1 + \mu + 4\mu^2} - 2\mu}{1 + \mu}, b = \frac{2\mu}{1 + \mu} - \frac{\sqrt{1 + \mu + 4\mu^2}}{1 + \mu} - 2$$

and, if $\mu = -1$,

$$A = \begin{pmatrix} 1 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{pmatrix}$$

q.e.d.

Using the above lemma, we are now able to prove the following

THEOREM 4.3. *Let G_0, G_1, \dots, G_{n-2} be a weakly connected Poncelet chain for triangles. Then, there exists a conic G_{n-1} such that $G_0, G_1, \dots, G_{n-1}, G_n = G_0$ is a closed weakly connected Poncelet chain for triangles.*

Proof. Choose G_{n-1} to be a lock between G_{n-2} and G_0 , as described in LEMMA 4.2.

q.e.d.

As an immediate consequence we get

COROLLARY 4.4. *There are closed weakly connected Poncelet chains for triangles of arbitrary length $n \geq 2$.*

Proof. For $n \geq 3$, this follows directly from THEOREM 4.3. For $n = 2$, the two conics

$$G_0 = \begin{pmatrix} 1 & 0 & \sqrt{3}/2 \\ 0 & 1 & 0 \\ \sqrt{3}/2 & 0 & -1/4 \end{pmatrix}, \quad G_1 = \begin{pmatrix} 1 & 0 & -\sqrt{3}/2 \\ 0 & 1 & 0 \\ -\sqrt{3}/2 & 0 & -1/4 \end{pmatrix},$$

form a closed chain of length 2.

q.e.d.

5 Rigid closed chains of Poncelet triangles

In a closed weakly connected Poncelet chain G_0, G_1, \dots, G_{n-1} of length n for triangles, as seen in Section 4, the Poncelet triangles Δ_i sitting between two consecutive conics (G_i, G_{i+1}) of the chain are not related to the neighbouring triangles. In this section, we will require, that the contact points of the sides of Δ_i on G_{i+1} are the vertices of the next triangle Δ_{i+1} . Then, the problem of finding closed chains G_0, \dots, G_{n-1} with a chain of Poncelet triangles $\Delta_0, \dots, \Delta_{n-1}$ satisfying this additional requirement, becomes more subtle.

A closed chain of triangles $\Delta_0, \dots, \Delta_{n-1}$ of length n is called a **rigid chain of Poncelet triangles** if there are conics G_0, \dots, G_{n-1} such that Δ_i has its vertices on G_i and is tangent to G_{i+1} , where the contact points are the vertices of Δ_{i+1} (where we take indices modulo n).

If we move one of the triangles, say Δ_0 , such that two of its sides are still tangent to G_1 , then, by Poncelet's Theorem, also the third side is tangent to G_1 . However, the contact points of the new triangle, which are points on G_1 , are not necessarily the vertices of a triangle whose sides are tangent to G_2 . So, in general we cannot move a rigid chain of Poncelet triangles (*i.e.*, the Poncelet triangles sit rigidly in the corresponding chain of conics).

By Corollary 4.4 we know that there are closed weakly connected Poncelet chains for triangles of arbitrary length; a similar result we get also for rigid closed chains of Poncelet triangles.

THEOREM 5.1. *There are rigid closed chains of Poncelet triangles of arbitrary length $n \geq 3$.*

Proof. We first give an example of a rigid chain of Poncelet triangles of length $n = 3$. Let


$$G_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad G_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad G_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 1 \end{pmatrix}.$$

It is not hard to see that the three triangles ΔABC , ΔPQR , ΔUVW on G_0 , G_1 , G_2 , respectively, where

$$A = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad B = \begin{pmatrix} -1 \\ -1 \\ 1 \end{pmatrix} \quad C = \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$$

$$P = \begin{pmatrix} 0 \\ -1 \\ 1 \end{pmatrix} \quad Q = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \quad R = \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}$$


$$U = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \quad V = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \quad W = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$$



A rigid chain of Poncelet triangles of length $n = 3$.

form a rigid closed chain of Poncelet triangles of length $n = 3$.

Now, if we enlarge the parabola and pinch the circle to an ellipse, we can plug in two additional ellipses between the parabola and the pinched circle in such a way, that the two additional triangles we get fit in the chain of pairwise Poncelet triangles. Notice that the hyperbola and the two triangles which are tangent to the hyperbola and the parabola respectively remain unchanged.




Two ellipses between the parabola and the pinched circle.

By a similar construction, we can plug in as many pairs of ellipses as we like. So, we can construct rigid closed chains of Poncelet triangles of arbitrarily odd length $n \geq 3$.

In order to construct rigid chains of Poncelet triangles of even length, we proceed as follows: We start with the same closed chain of Poncelet triangles as above. Then we enlarge and reflect the parabola and pinch the circle to an ellipse, such that we can plug

in an additional ellipse between the parabola and the pinched circle in such a way, that the additional triangle fits in the chain of pairwise Poncelet triangles. Notice that the hyperbola and the two triangles which are tangent to the hyperbola and the parabola respectively remain unchanged (see figure below).



An ellipse between the reflected parabola and the pinched circle.

Like above, we can now plug in arbitrarily many pairs of ellipses. So, we can construct rigid chains of Poncelet triangles of arbitrarily even length $n \geq 4$. *q.e.d.*

6 Miraculous chains of Poncelet polygons

In general, the triangles Δ_i in a closed rigid chain of Poncelet triangles satisfy the condition, that the contact points of Δ_i on G_{i+1} are the vertices of Δ_{i+1} , only in *one* particular position. In this section, we investigate the question, whether closed rigid chains exist such that the contact condition is satisfied in *every* position. For this, we first recall some relevant facts and notations.

If a point x moves along a conic G_0 , then each polar of x with respect to a second conic G_1 is tangent to one particular conic G_2 , which is called the **conjugate of G_0 with respect to G_1** (see [15, Theorem 1.5]). In particular, if K_0 is a Poncelet k -gon, inscribed in G_0 and circumscribed about G_1 , then the k -gon K_1 whose vertices are the contact points of K_0 on G_1 is tangent to the conjugate conic G_2 of G_0 with respect to G_1 . Hence K_1 is itself a Poncelet k -gon for the pair G_1 and G_2 . Obviously, this process can be iterated. Astonishingly, there are very particular configurations, where this process closes after a finite number of steps, *i.e.*, the n -th k -gon K_n is equal to K_0 .

For an integer $k \geq 3$, a closed chain of pairwise conjugate conics G_0, \dots, G_{n-1} (*i.e.*, G_i and G_{i+2} are conjugate with respect to G_{i+1}) is a **miraculous chain for Poncelet k -gons** if for each $0 \leq i < n$, G_i carries the vertices of a k -gon K_i which is tangent to G_{i+1} , where

the contact points are the vertices of K_{i+1} (where we take indices modulo n).

Since we require that the conics G_0, \dots, G_{n-1} are pairwise conjugate, we can move the vertices of the k -gons without losing the property that the contact points of a k -gon are the vertices of the next k -gon.

We shall see that up to projective transformations, there is exactly one miraculous chain of Poncelet triangles and one miraculous chain for Poncelet quadrilaterals.

6.1 Closed chains of conjugate conics

From [15, Theorem 2.1 & 2.2]), we infer the following: Let G_0, G_1, \dots be a sequence of conics such that G_{i+2} is the conjugate of G_i with respect to G_{i+1} , for all indices $i \geq 0$. Then, $G_{i+2} \sim G_1(G_0^{-1}G_1)^{i+1}$.

Let I denote the 3×3 identity matrix. A chain $G_0, G_1, G_2, \dots, G_{n-1}, G_n = G_0$ of length n of conjugate conics is closed *iff* there are representatives of the conics such that

$$(G_0^{-1}G_1)^n = I.$$

and n is minimal with this property.

We now want to investigate, in which of the 8 cases listed in Section 3, closed chains may exist. For this, let G_0, \dots, G_{n-1} , for some $n \geq 3$, be a closed chain of conjugate conics. We may assume, that G_0, G_1 are in standard form, as in the proof of Lemma 4.2.

Case 1a and b.

$$G_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad \text{and} \quad G_1 = \begin{pmatrix} a^2 & 0 & 0 \\ 0 & \pm b^2 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad a, b > 0.$$

Then, $(G_0^{-1}G_1)^n = \text{diag}(a^{2n}, (\pm b^2)^n, 1)$. This is the identity matrix only for $a = b = 1$ which is excluded. Thus, if G_0, G_1 have four intersections, they cannot be part of a closed chain of conjugate conics.

Cases 2. By Theorem 3.3 we may assume that

$$G_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad \text{and} \quad G_1 = \begin{pmatrix} 1 & 0 & -a \\ 0 & 1 & 0 \\ -a & 0 & a^2 - r^2 \end{pmatrix}$$

for some $0 \leq a$ and $r > 0$. So, G_0 is the unit circle and G_1 is a circle with center $(a, 0, 1)$ and radius r . Observe, that for any k the second element of the diagonal of $(G_0^{-1}G_1)^k$ equals 1, and $\det(G_0^{-1}G_1) = r^2$. Hence, it follows from $(G_0^{-1}G_1)^n = I$, that $r = 1$ and hence $a > 2$. But then, G_{i+1} is nested inside G_i for each $i > 1$ and the chain cannot be closed.

Case 3. As in Case 2, we may assume

$$G_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad \text{and} \quad G_1 = \begin{pmatrix} 1 & 0 & -a \\ 0 & 1 & 0 \\ -a & 0 & a^2 - r^2 \end{pmatrix}, \quad a \geq 0$$

and as above, we conclude that $r = 1$, and therefore $a \neq 0$. Hence,

$$G_1 = \begin{pmatrix} 1 & 0 & -a \\ 0 & 1 & 0 \\ -a & 0 & a^2 - 1 \end{pmatrix}.$$

Next, we move the two unit circles $\frac{a}{2}$ to the left and exchange the axes x_2 and x_3 . Then, since we assume that the two conics intersect in two points, we have $a < 2$ and by a suitable scaling of the x_1 and x_2 axes, we arrive at the following form:

$$\tilde{G}_0 = \begin{pmatrix} \frac{\sqrt{4-a^2}}{2} & \frac{a}{2} & 0 \\ \frac{a}{2} & -\frac{\sqrt{4-a^2}}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad \tilde{G}_1 = \begin{pmatrix} \frac{\sqrt{4-a^2}}{2} & -\frac{a}{2} & 0 \\ -\frac{a}{2} & -\frac{\sqrt{4-a^2}}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Finally, a rotation in the first two variables leads to the final form of two hyperbolas

$$H_0 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad H_\alpha = \begin{pmatrix} -(1 - \frac{a^2}{2}) & \frac{a}{2}\sqrt{4-a^2} & 0 \\ \frac{a}{2}\sqrt{4-a^2} & 1 - \frac{a^2}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} -\cos(\alpha) & \sin(\alpha) & 0 \\ \sin(\alpha) & \cos(\alpha) & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

where $0 < \alpha < \pi$. Now, the condition $(H_0^{-1}H_\alpha)^n = I$, and n minimal with this property, is equivalent to $\alpha = \frac{2\pi}{n}$. So, this is the standard form if the two initial conics have two points of intersection.

Case 4. Here,

$$G_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad \text{and} \quad G_1 = \begin{pmatrix} \mu + 1 & 0 & -\mu \\ 0 & 1 - \mu & 0 \\ -\mu & 0 & \mu - 1 \end{pmatrix}, \quad \mu \notin \{0, 1\}.$$

Then, for each n , $(G_0^{-1}G_1)^n =: H_n \neq I$, since the third element of the first column of H_n equals $n\mu$.

Case 5. By the same reasoning as in Case 2 above, we may exclude that a closed chain of conjugate conics exist such that two consecutive conics have one first order contact.

Case 6. Here,

$$G_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad \text{and} \quad G_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 - \mu & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad \mu \neq 0.$$

Then, $(G_0^{-1}G_1)^n = \text{diag}(1, (1 - \mu)^n, 1) = I$ iff $(1 - \mu)^n = 1$. Then either $\mu = 0$ (which is excluded) or $\mu = 2$ and $n = 2$. (Recall that n is supposed to be minimal). By exchanging the x_2 - and the x_3 -axis, we find

$$G_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad G_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

which as standard form of the only existing closed chain of conjugate conics such that two consecutive conics have two first order contacts. Such a chain has necessarily length 2.

Case 7. Here,

$$G_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad \text{and} \quad G_1 = \begin{pmatrix} 1 & -\mu & 0 \\ -\mu & 1 & \mu \\ 0 & \mu & -1 \end{pmatrix}, \quad \mu \neq 0.$$

Here, the second element of the first row of $(G_0^{-1}G_1)^n$ equals $-n\mu \neq 0$. Hence in this situation, no closed chain of conjugate conics exists.

Case 8. For

$$G_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad \text{and} \quad G_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \mu + 1 & -\mu \\ 0 & -\mu & \mu - 1 \end{pmatrix}, \quad \mu \neq 0.$$

Here, the third element of the second row of $(G_0^{-1}G_1)^n$ equals $-n\mu \neq 0$. Hence in this situation, no closed chain of conjugate conics exists either.

6.2 Miraculous chains of Poncelet triangles and quadrilaterals

It is geometrically easy to see that the closed chain of conjugate conics of length 2 in Case 6 in the previous section cannot carry Poncelet polygons. Hence, the only chance to find closed chains of conjugate conics which carry Poncelet polygons are H_0 and H_1 from Case 3 in the previous section.

For $D(\lambda) = \det(H_0 + \lambda H_1)$ and $\sqrt{D(\lambda)} = c_0 + c_1\lambda + c_2\lambda^2 + c_3\lambda^3 + \dots$ we compute

$$c_0 = 1, \quad c_1 = \frac{1}{2} + \cos(\alpha), \quad c_2 = -\frac{1}{8} + \frac{\cos(\alpha)}{2} + \frac{\sin(\alpha)^2}{2},$$

$$c_3 = \frac{1}{16} - \frac{\cos(\alpha)}{8} + \frac{\sin(\alpha)^2}{4} - \frac{\cos(\alpha)\sin(\alpha)^2}{2}, \dots$$


In order to get a miraculous chain of Poncelet triangles, we must find $0 < \alpha < \pi$ such that $\alpha = \frac{2\pi}{n}$ (for some $n \geq 3$) and $c_2 = 0$. If we replace $\cos(\alpha)$ with t , then the equation $c_2 = 0$ is equivalent to the equation

$$(-3 + 2t)(1 + 2t) = 0.$$


As solutions we get $t_1 = \frac{3}{2}$ and $t_2 = -\frac{1}{2}$. Since $-1 \leq \cos(\alpha) \leq 1$, there is no $0 < \alpha < \pi$ such that $t_1 = \cos(\alpha)$. On the other hand, for $\alpha = \frac{2\pi}{3}$ we get $t_2 = \cos(\alpha)$. Hence, up to projective transformations, there is exactly one miraculous chain of Poncelet triangles. The figure below shows the canonical miraculous chain with three hyperbolas and three triangles.

In order to get a miraculous chain of Poncelet quadrilateral, we must find $0 < \alpha < \pi$ such that $\alpha = \frac{2\pi}{n}$ (for some $n \geq 3$) and $c_3 = 0$. If we replace $\cos(\alpha)$ again with t , then the equation $c_3 = 0$ is equivalent to the equation

$$(-1 + 2t)(-5 + 4t^2) = 0.$$



Canonical miraculous chain of Poncelet triangles



Canonical miraculous chain of Poncelet quadrilaterals

The only solution with $-1 < t < 1$ is $t = \frac{1}{2}$, which gives us $\alpha = \frac{\pi}{3}$. Hence, up to projective transformations, there is exactly one miraculous chain of Poncelet quadrilaterals. The figure above shows the canonical miraculous chain with six hyperbolas and two of the six quadrilaterals.

6.3 Are there other miraculous chains of Poncelet polygons?

In order to get a miraculous chain of Poncelet pentagons, we must find $0 < \alpha < \pi$ such that $\alpha = \frac{2\pi}{n}$ (for some $n \geq 3$) and $c_2c_4 - c_3^2 = 0$. For $t = \cos(\alpha)$, the latter equation is equivalent to the equation

$$(17 - 10t - 20t^2 + 8t^3)(-11 + 6t - 4t^2 + 8t^3) = 0$$

where the two factors are irreducible. Now, if $\cos(\alpha)$ is a root of one of these factors, where $\alpha = \frac{2\pi}{n}$ (for some $n \geq 3$), then, since cyclotomic polynomials are irreducible, all roots of that factor must be in the open interval $(-1, 1)$. With Sturm's Theorem one can now count, how many roots of the factors $17 - 10t - 20t^2 + 8t^3$ and $-11 + 6t - 4t^2 + 8t^3$ respectively, belong to $[-1, 1]$. Since this number is less than 3 (which is the degree of the factors), we can be sure that the factor does not have a root of the form $\cos(\frac{2\pi}{n})$, i.e., there is no miraculous chain of Poncelet pentagons.

In order to show that there is no miraculous chain of Poncelet hexagons, we consider the equation $c_3c_5 - c_4^2 = 0$, which is equivalent to the equation

$$(-3 + 2t)(1 + 2t)(-7 + 4t + 4t^2)(-41 + 64t - 8t^2 - 64t^3 + 48t^4) = 0.$$

Since the root of the factor $1 + 2t$ leads to $\alpha = \frac{2\pi}{3}$, the hexagon is just a triangle run through twice. Thus, we have just to consider the other three factors and with the same arguments as above we can show that there are no miraculous chains of Poncelet hexagons.

With this technique, we have shown that except for $k = 3$ and $k = 4$, there are no miraculous chains of Poncelet k -gons for $k \leq 23$. This motivates the following

Conjecture: There are no miraculous chains of Poncelet k -gons for $k \geq 5$.

References

- [1] ROBERT BIX, *Conics and Cubics: A Concrete Introduction to Algebraic Curves*, Undergraduate Texts in Mathematics, Springer New York, 2006.
- [2] HENK J. M. BOS, CEES KERS, FRANS OORT AND DIEDERICK W. RAVEN, *Poncelet's closure theorem*, *Exposition. Math.*, 5 (1987), no. 4, 289–364.
- [3] ROBERT L. BRYANT, *Poncelet's theorem*, http://mathcircle.berkeley.edu/archivedocs/2010__2011/lectures/1011lecturespdf/PonceletforBMC.pdf, 2010.
- [4] ARTUR CAYLEY, *Developments on the porism of the in-and-circumscribed polygon*, *Philosophical Magazine*, 7 (1854), 339–345.
- [5] WILLIAM CHAPPLE, *An essay on the properties of triangles inscribed in and circumscribed about two given circles*, *Miscellanea Curiosa Mathematica* (1746), no. 4, 117–124.

- [6] WALDEMAR CIEŚLAK, HORST MARTINI AND WITOLD MOZGAWA, *On the rotation index of bar billiards and Poncelet's porism*, **Bull. Belg. Math. Soc. Simon Stevin**, 20 (2013), no. 2, 287–300.
- [7] WALDEMAR CIEŚLAK AND WITOLD MOZGAWA, *In search of a measure in Poncelet's porism*, **Acta Math. Hungar.**, 149 (2016), no. 2, 338–345.
- [8] ANDREA DEL CENTINA, *Poncelet's porism: a long story of renewed discoveries, I*, **Arch. Hist. Exact Sci.**, 70 (2016), no. 1, 1–122.
- [9] ———, *Poncelet's porism: a long story of renewed discoveries, II*, **Arch. Hist. Exact Sci.**, 70 (2016), no. 2, 123–173.
- [10] VLADIMIR DRAGOVIĆ AND MILENA RADNOVIĆ, ***Poncelet Porisms and Beyond: Integrable Billiards, Hyperelliptic Jacobians and Pencils of Quadrics***, Frontiers in Mathematics, Springer, 2011.
- [11] LEONHARD EULER, *Solutio facilis problematum quorundam geometricorum difficiliorum*, **Novi Commentarii academiae scientiarum imperialis Petropolitanae** (1767), no. 11, 103–123.
- [12] LEOPOLD FLATTO, ***Poncelet's Theorem***, American Mathematical Society, Providence, RI, 2009.
- [13] NICOLAS FUSS, *De polygonis symmetrice irregularibus circulo simul inscriptis et circumscriptis*, **Nova Acta Petropol** (1798), no. 13, 166–189.
- [14] LORENZ HALBEISEN AND NORBERT HUNGERBÜHLER, *A simple proof of Poncelet's theorem (on the occasion of its bicentennial)*, **American Mathematical Monthly**, 122 (2015), 537–551.
- [15] ———, *Conjugate conics and closed chains of Poncelet polygons*, **Mitteilungen der Mathematischen Gesellschaft in Hamburg**, 36 (2016), 5–28.
- [16] YOO PYO HONG, ROGER A. HORN AND CHARLES R. JOHNSON, *On the reduction of pairs of Hermitian or symmetric matrices to diagonal form by congruence*, **Linear Algebra Appl.**, 73 (1986), 213–226.
- [17] JONATHAN L. KING, *Three problems in search of a measure*, **American Mathematical Monthly**, 101 (1994), 609–628.
- [18] ERIC LORD, ***Symmetry and Pattern in Projective Geometry***, Springer London, 2012.
- [19] ERKKI PESONEN, *Über die Spektraldarstellung quadratischer Formen in linearen Räumen mit indefiniter Metrik*, **Ann. Acad. Sci. Fenn. Ser. A. I.**, 227 (1956), 31.
- [20] SYLVAIN PETITJEAN, *Invariant-based characterization of the relative position of two projective conics*, Nonlinear computational geometry, IMA Vol. Math. Appl., 151, Springer, New York, 2010, pp. 189–220.

- [21] JEAN-VICTOR PONCELET, *Traité des propriétés projectives des figures*, Bachelier, Paris, 1822.
- [22] QUESTIONS PROPOSÉES, *Problème de géométrie*, *Ann. Math. Pures Appl.*, tome 7 (1816–1817), 128.
- [23] JACOB STEINER, *Geometrische Aufgaben und Lehrsätze*, *J. Reine Angew. Math.*, 2 (1827), 96–98.
- [24] ———, *Geometrische Lehrsätze*, *J. Reine Angew. Math.*, 2 (1827), 287–292.
- [25] PAR UN ABONNÉ, *Questions résolues. Ébauche de solution du problème de géométrie proposé à la page 128 de ce recueil*, *Ann. Math. Pures Appl.*, tome 7 (1816–1817), 373–379.
- [26] JEAN VALLÈS, *A vector bundle proof of Poncelet's closure theorem*, *Expo. Math.*, 30 (2012), no. 4, 399–405.