

MATHIAS ABSOLUTENESS AND THE RAMSEY PROPERTY

Lorenz Halbeisen
ETH Zürich
Switzerland

Haim Judah¹
Bar-Ilan University
Israel

Abstract

In this article we give a forcing characterization for the Ramsey property of Σ_2^1 -sets of reals. This research was motivated by the well-known forcing characterizations for Lebesgue measurability and the Baire property of Σ_2^1 -sets of reals. Further we will show the relationship between higher degrees of forcing absoluteness and the Ramsey property of projective sets of reals.

1 Notations and Definitions

Most of our set-theoretical notations and notations of forcings are standard and can be found in [Je 2] or [Ku]. An exception is, that we will write A^B for the set of all functions from B to A , instead of ${}^B A$ because we never use ordinal arithmetic. $A^{<\omega}$ is the set of all partial functions f from ω to A , such that the cardinality of $\text{dom}(f)$ is finite.

First we will give the definitions of the sets we will consider as the real numbers.

Let $[x]^\kappa := \{y \subseteq x : |y| = \kappa\}$ and $[x]^{<\kappa} := \{y \subseteq x : |y| < \kappa\}$, where $|y|$ denotes the cardinality of y . For $x \in [\omega]^\omega$, we will consider $[x]^{<\omega}$ as the set of strictly increasing, finite sequences in x and $[x]^\omega$ as the set of strictly increasing, infinite sequences in x . For $x \in [\omega]^\omega$ and $n \in \omega$ let $x(n)$ be such that $x(n) \in x$ and $|x(n) \cap x| = n$.

We can consider $[\omega]^\omega$ also as a set of infinite 0-1-sequences

$$\begin{aligned} [\omega]^\omega &\longrightarrow 2^\omega \\ x &\longmapsto f \text{ such that } f(n) = 1 \text{ iff } n \in x, \end{aligned}$$

or as the infinite sequences in ω

$$\begin{aligned} [\omega]^\omega &\longrightarrow \omega^\omega \\ x &\longmapsto \langle a_n : n \in \omega \rangle \text{ such that: } a_0 := x(0) \text{ and} \\ & a_{n+1} := x(n+1) - x(n) - 1. \end{aligned}$$

Note that these two mappings are bijective.

¹The research for this paper was partially supported by the Emmy Nöther Institute in mathematics of Bar-Ilan and the Minerva foundation, Germany.

The Baire space

The *Baire space* is the space ω^ω of all infinite sequences of natural numbers, $\langle a_n : n \in \omega \rangle$, with the following topology: For every finite sequence $s = \langle a_k : k < n \rangle$, let

$$U_s := \{f \in \omega^\omega : s \subset f\} = \{\langle c_k : k \in \omega \rangle : \forall k < n (c_k = a_k)\}.$$

The sets U_s ($s \in \omega^{<\omega}$) form a basis for the topology of ω^ω . Note that each U_s is also closed. The Baire space is homeomorphic to the space of all irrational numbers in $[0, 1]$ with the topology of the real line (cf. [Je 2] p. 36).

Because the mapping given above between $[\omega]^\omega$ and ω^ω is bijective, we can endow $[\omega]^\omega$ with the induced topology and will not distinguish between the two spaces $[\omega]^\omega$ and ω^ω . The same holds for the sets $[\omega]^\omega$ and 2^ω .

Three properties of sets of reals

Let us work in the topological space $[\omega]^\omega$.

A set $R \subseteq [\omega]^\omega$ is *rare* (or *nowhere dense*) if the complement of R contains a dense open set and a set $M \subseteq [\omega]^\omega$ is *meager* (or of *first category*) if M is the union of countably many rare sets. A nonmeager set is also called a set of *second category*. A set $A \subseteq [\omega]^\omega$ has the *Baire property* if there exists an open set $G \subseteq [\omega]^\omega$ such that the symmetric difference $A\Delta G = (A \setminus G) \cup (G \setminus A)$ is meager.

A set $N \subseteq [\omega]^\omega$ is *null* if N considered as a set of reals has Lebesgue measure zero. A set $A \subseteq [\omega]^\omega$ is *Lebesgue measurable* if there is a Borel set B such that the symmetric difference $A\Delta B$ is null.

A set $A \subseteq [\omega]^\omega$ has the *Ramsey property* (or is *Ramsey*) if $\exists x \in [\omega]^\omega ([x]^\omega \subseteq A \vee [x]^\omega \cap A = \emptyset)$. If there exists an x such that $[x]^\omega \cap A = \emptyset$ we call A a *Ramsey_o* set and if $[x]^\omega \subseteq A$ we call A a *co-Ramsey_o* set. Note that A can also be both. A set $A \subseteq [\omega]^\omega$ is called *uniformly Ramsey_o* if, for each $x \in [\omega]^\omega$ there is a $y \in [x]^\omega$ such that $[y]^\omega \cap A = \emptyset$.

The hierarchy of projective sets

We always consider the boldface Σ_n^1 hierarchy (see [Je 2] p.510). A Σ_1^1 -set is the projection of a closed set. The Σ_1^1 -sets are also called *analytic* sets. The Π_1^1 -sets are the complements of the analytic sets. A Σ_{n+1}^1 -set is the projection of a Π_n^1 -set and the Π_{n+1}^1 -sets are the complements of the Σ_{n+1}^1 -sets. A set is Δ_n^1 if it is Σ_n^1 and Π_n^1 . For the normal form of the formulas representing projective sets and relations cf. [Je 2] Section 40. Further we will consider a Σ_n^1 -relation without free variables as a Σ_n^1 -sentence.

If all Σ_n^1 -sets with parameters in $V \cap W$ are Ramsey, (are Lebesgue measurable, have the Baire property, respectively), with respect to V , we will write $V \models \Sigma_n^1(\mathcal{R})_W$ ($V \models \Sigma_n^1(\mathcal{L})_W$, $V \models \Sigma_n^1(\mathcal{B})_W$, respectively). If $V = W$, then we do not write the index W . The notations $\Delta_n^1(\mathcal{R})_W$, $\Delta_n^1(\mathcal{L})_W$, $\Delta_n^1(\mathcal{B})_W$, $\Pi_n^1(\mathcal{R})_W$, $\Pi_n^1(\mathcal{L})_W$ and $\Pi_n^1(\mathcal{B})_W$ are similar. Note that because of the three properties are closed under complements, the statements $\Sigma_n^1(\mathcal{R})$, $\Sigma_n^1(\mathcal{L})$ and $\Sigma_n^1(\mathcal{B})$ are equivalent to $\Pi_n^1(\mathcal{R})$, $\Pi_n^1(\mathcal{L})$ and $\Pi_n^1(\mathcal{B})$, respectively.

Filters and Families on ω

$\mathcal{F} \subseteq [\omega]^\omega$ is a *Ramsey family* if for all $\pi \in 2^{[\omega]^2}$ there is an $h \in \mathcal{F}$ such that $\pi|_{[h]^2}$ is constant.

$\mathcal{F} \subseteq [\omega]^\omega$ is a *dominating family* if for all $x \in [\omega]^\omega$ there is a $d \in \mathcal{F}$ and a natural number $n \in \omega$ such that for all $k \geq n$: $d(k) \geq x(k)$.

$\mathcal{F} \subseteq [\omega]^\omega$ is *dominated* by the real d if for all $f \in \mathcal{F}$ there is a natural number $n \in \omega$ such that for all $k \geq n$: $d(k) \geq f(k)$. (In this case we call d a *dominating real* with respect to \mathcal{F}).

$\mathcal{F} \subseteq [\omega]^{\leq \omega}$ is a *filter* (on ω) if $\omega \in \mathcal{F}$ and for all $x, y \in [\omega]^{\leq \omega}$: if $x, y \in \mathcal{F}$ then $x \cap y \in \mathcal{F}$ and if $x \in \mathcal{F}$, $x \subseteq y$ then $y \in \mathcal{F}$.

A filter \mathcal{F} is *proper* if $\emptyset \notin \mathcal{F}$.

A filter \mathcal{F} is an *ultrafilter* if it is proper and for every $x \in [\omega]^{\leq \omega}$, either $x \in \mathcal{F}$ or $\omega \setminus x \in \mathcal{F}$. The filter $\mathcal{F} = \{x \in [\omega]^\omega : |\omega \setminus x| < \omega\}$ is called the *Fréchet filter*.

A Ramsey ultrafilter is a Ramsey family which is also an ultrafilter. We consider only filters which are proper and contain the Fréchet filter.

Some notions of forcing

We recall the definition of the following seven notions of forcing.

(i) The Amoeba (measure) forcing **A**:

$$p \in \mathbf{A} \Leftrightarrow p \subseteq 2^\omega \text{ is a perfect tree } \wedge \mu(p) > \frac{1}{2},$$

$$p \leq q \Leftrightarrow p \subseteq q.$$

(ii) The Random forcing **B**:

$$p \in \mathbf{B} \Leftrightarrow p \subseteq 2^\omega \text{ is a perfect tree } \wedge \mu(p) > 0,$$

$$p \leq q \Leftrightarrow p \subseteq q.$$

(iii) The Cohen forcing **C**:

$$p \in \mathbf{C} \Leftrightarrow p \in 2^{< \omega},$$

$$p \leq q \Leftrightarrow p \text{ extends } q.$$

(iv) The Hechler forcing **D**:

$$\langle n, f \rangle \in \mathbf{D} \Leftrightarrow n \in \omega \wedge f \in \omega^\omega,$$

$$\langle n, f \rangle \leq \langle m, g \rangle \Leftrightarrow n \geq m \wedge f|_m = g|_m \wedge \forall k (f(k) \geq g(k)).$$

(v) The Mathias forcing **M**:

$$\langle s, S \rangle \in \mathbf{M} \Leftrightarrow s \in [\omega]^{< \omega} \wedge S \in [\omega]^\omega \wedge \max(\text{range}(s)) < \min(S),$$

$$\langle s, S \rangle \leq \langle t, T \rangle \Leftrightarrow s \text{ extends } t \wedge S \subseteq T \wedge \forall i \in \text{dom}(s) \setminus \text{dom}(t) (s(i) \in T).$$

(vi) The forcing notion $\mathbf{P}(D)$ for an ultrafilter D :

$$p_s \in \mathbf{P}(D) \Leftrightarrow p_s \subseteq [\omega]^{< \omega} \text{ is a tree and there is an } s \in p_s \text{ such that}$$

$$\forall t \in p_s ((s \subseteq t \vee t \subseteq s) \wedge (s \subseteq t \rightarrow \{n : t \frown n \in p_s\} \in D)),$$

$$p_s \leq q_t \Leftrightarrow p_s \subseteq q_t.$$

(vii) The forcing notion \mathbf{P}_D for an ultrafilter D :

$$\langle s, a \rangle \in \mathbf{P}_D \Leftrightarrow s \in [\omega]^{< \omega} \wedge a \in [\omega]^\omega \wedge a \in D \wedge \max(\text{range}(s)) < \min(a),$$

$$\langle s, a \rangle \leq \langle t, b \rangle \Leftrightarrow s \text{ extends } t \wedge a \subseteq b \wedge \forall i \in (\text{dom}(s) \setminus \text{dom}(t)) (s(i) \in b).$$

In the forcing notions (v),(vi) and (vii) we call s the *stem* of the condition $\langle s, S \rangle$, p_s and $\langle s, a \rangle$, respectively. A generic object over one of these seven forcing notions can be considered as a generic real and we will handle the generic reals like the corresponding

generic objects. For example if $G_{\mathbf{M}}$ is Mathias generic and $p \in G_{\mathbf{M}}$ (for a Mathias condition p), then we write $p \in m$ (for m Mathias generic real) and if p has empty stem ($p = \langle \emptyset, S \rangle$), we also write $m \subseteq p$. Note that the conditions of these seven forcing notions can also be considered as reals, (and the meaning of $r_1 \leq r_2$ is clear). Let p, q are Mathias conditions, then we write $p \leq^0 q$ to say that p and q have the same stem and $p \leq q$. Names in the forcing language are denoted with a “ \sim ” over the letter. Canonical names for generic objects are usually denoted by boldface letters and canonical names for objects in the ground model we denote with a “ $\check{\cdot}$ ” over the letter.

Forcing-absoluteness

Let \mathbf{P} be a notion of forcing. We define

$$V^{\mathbf{P}} \models \Phi \iff V \models \mathbf{1} \Vdash_{\mathbf{P}} \Phi$$

where Φ is a formula with parameters in V and $\mathbf{1}$ is the weakest condition of \mathbf{P} .

Now we say V is Σ_n^1 - \mathbf{P} -absolute if for all Σ_n^1 -sentences φ with parameters in V ,

$$V^{\mathbf{P}} \models \varphi \text{ iff } V \models \varphi.$$

Or equivalently, if for all \mathbf{P} -generic objects $G_{\mathbf{P}}$ over V :

$$V[G_{\mathbf{P}}] \models \varphi \text{ iff } V \models \varphi.$$

2 Introduction

In this section we give a list of results. Some of them are well-known, others gave the motivation to this work.

Characterizations with generic reals

Because the canonical well-ordering of constructible reals is Δ_2^1 (cf. [Je 2] Theorem 97), Gödel’s constructible universe L is neither a model for $\Delta_2^1(\mathcal{B})$ nor $\Delta_2^1(\mathcal{L})$ nor $\Delta_2^1(\mathcal{R})$. Hence, a model V of set theory in which one of these properties holds, has to be larger than L . In fact, V has to contain even some reals which are generic over L .

THEOREM 2.1 (i) $V \models \Delta_2^1(\mathcal{B})$ if and only if for all reals $r \in V$ the set of reals in V which are Cohen over $L[r]$ is not empty.

(ii) $V \models \Delta_2^1(\mathcal{L})$ if and only if for all reals $r \in V$ the set of reals in V which are random over $L[r]$ is not empty.

(iii) $V \models \Delta_2^1(\mathcal{R})$ if and only if for all reals $r \in V$ the set of reals in V which are Ramsey over $L[r]$ is not empty.

PROOF: All three results were proved in [JS 1]. ↯

A similar characterization we also have for Σ_2^1 -sets.

THEOREM 2.2 (i) $V \models \Sigma_2^1(\mathcal{B})$ if and only if for all reals $r \in V$ the set of reals in V which are Cohen over $L[r]$ is co-meager.

(ii) $V \models \Sigma_2^1(\mathcal{L})$ if and only if for all reals $r \in V$ the set of reals in V which are random over $L[r]$ has measure 1.

(iii) $V \models \Sigma_2^1(\mathcal{R})$ if and only if for all reals $r \in V$ the set of reals in V which are Ramsey over $L[r]$ is co-Ramsey_o.

PROOF: A proof can be found in [BJ]. For the third result see also [JS 1]. ↯

Characterizations with forcing absoluteness

For the Σ_2^1 -sets we also find a characterization with forcing absoluteness.

THEOREM 2.3 (i) $V \models \Sigma_2^1(\mathcal{B})$ if and only if V is Σ_3^1 -Hechler-absolute.
(ii) $V \models \Sigma_2^1(\mathcal{L})$ if and only if V is Σ_3^1 -Amoeba-absolute.
(iii) $V \models \Sigma_2^1(\mathcal{R})$ if and only if V is Σ_3^1 -Mathias-absolute.

PROOF: The first two results were proved in [Ju 1] and [Ju 2]. A proof of the last one will be given in this work, Theorem 4.1. \dashv

For higher levels in the projective hierarchy, we lose the forcing characterization with Mathias forcing for the Ramsey property. We will show in Theorem 5.3 that

$$\Sigma_4^1\text{-Mathias-absoluteness} \implies \Sigma_3^1(\mathcal{R})$$

but (Theorem 5.2)

$$\Delta_3^1(\mathcal{R}) \not\implies \Sigma_4^1\text{-Mathias-absoluteness}.$$

The reason for this is, that if V is Σ_4^1 -Mathias-absolute, then ω_1^V is inaccessible in L . On the other hand we can build a model in which $\Delta_3^1(\mathcal{R})$ holds without using inaccessible cardinals.

We will show further (Corollary 6.1) that

$$\Sigma_5^1\text{-Mathias-absoluteness} \implies \Delta_4^1(\mathcal{R}),$$

and moreover (Corollary 6.5)

$$\Sigma_6^1\text{-Mathias-absoluteness} \implies \Delta_5^1(\mathcal{R}).$$

3 The Ramsey Property and Mathias Forcing

Basic facts about the Ramsey property

FACT 3.1 If $A \subseteq [\omega]^\omega$ is Ramsey and $C \subseteq [\omega]^\omega$ is uniformly Ramsey_o (e.g. countable), then both, $A \cup C$ and $A \setminus C$ are Ramsey.

PROOF: To see that $A \cup C$ is Ramsey, first note that if there is an $x \in [\omega]^\omega$ such that $[x]^\omega \subseteq A$, we are done. Otherwise, pick x such that $[x]^\omega \cap A = \emptyset$ and pick $y \in [x]^\omega$ such that $[y]^\omega \cap C = \emptyset$. Now $[y]^\omega \cap (A \cup C) = \emptyset$.

To see that $A \setminus C$ is Ramsey, again note that if there is an x such that $[x]^\omega \cap A = \emptyset$, we are done. Otherwise, pick x such that $[x]^\omega \subseteq A$. Now there is a $y \in [x]^\omega$ such that $[y]^\omega \cap C = \emptyset$ and $[y]^\omega \subseteq (A \setminus C)$. \dashv

FACT 3.2 The axiom of choice implies that there are sets without the Ramsey property.

PROOF: Define on $[\omega]^\omega$ an equivalence-relation as follows:

$$x \sim y \quad \text{iff} \quad |x \Delta y| \text{ is finite.}$$

Now choose from each equivalence class x^\sim an element c_x . Further define:

$$f(x) := \begin{cases} 1 & \text{if } |x \Delta c_x| \text{ is odd,} \\ 0 & \text{otherwise.} \end{cases}$$

Then the set $\{x : f(x) = 1\}$ is evidently not Ramsey. \dashv

The first example of a set which does not have the Ramsey property is given in [ER]. A lot of other examples can be found in [Bn] and [Co].

FACT 3.3 *Analytic sets (these are the Σ_1^1 -sets) are Ramsey.*

PROOF: A proof can be found in [El] and [Si]. \dashv

The forcing notions $\mathbf{P}(D)$, \mathbf{P}_D and \mathbf{M}

(Compare also with [Ma]).

Let $\mathcal{J} = [\omega]^{<\omega}$ be the ideal of finite sets and let $\langle \mathcal{P}(\omega)/\mathcal{J}, \leq \rangle =: \mathbf{U}$ be the partial order defined as follows:

$$\begin{aligned} p \in \mathbf{U} &\Leftrightarrow p \in [\omega]^\omega, \\ p \leq q &\Leftrightarrow p \setminus q \in \mathcal{J} \quad (\text{this is } p \subseteq^* q). \end{aligned}$$

FACT 3.4 *Let D be \mathbf{U} -generic over V , then D is a Ramsey ultrafilter in $V[D]$.*

PROOF: First note that \mathbf{U} is \aleph_0 -closed, hence adds no new reals to V , (cf.[Je 2] Lemma 19.6). Let $\pi \in 2^{[\omega]^2}$, then by the Ramsey Theorem (cf.[Je 2] Lemma 29.1) for each $p \in [\omega]^\omega$ there exists a $q \subseteq^* p$ such that π is constant on $[q]^2$. Therefore $H_\pi := \{q \in [\omega]^\omega : \pi|_{[q]^2} \text{ is constant}\}$ is dense in \mathbf{U} , hence $H_\pi \cap D \neq \emptyset$. \dashv

LEMMA 3.5 *Let \tilde{D} be the canonical \mathbf{U} -name for the \mathbf{U} -generic object, then*

$$\mathbf{U} * \mathbf{P}_{\tilde{D}} \approx \mathbf{M}.$$

PROOF:

$$\begin{aligned} \mathbf{U} * \mathbf{P}_{\tilde{D}} &= \{ \langle p, \langle \tilde{s}, \tilde{a} \rangle \rangle : p \in \mathbf{U} \wedge p \Vdash_{\mathbf{U}} \langle \tilde{s}, \tilde{a} \rangle \in \mathbf{P}_{\tilde{D}} \} \\ &= \{ \langle p, \langle \tilde{s}, \tilde{a} \rangle \rangle : p \in [\omega]^\omega \wedge p \Vdash_{\mathbf{U}} (\tilde{a} \in \tilde{D} \wedge \max(\text{range}(\tilde{s})) < \min(\tilde{a})) \}. \end{aligned}$$

Now the embedding

$$\begin{aligned} h : \quad \mathbf{M} &\longrightarrow \mathbf{U} * \mathbf{P}_{\tilde{D}} \\ \langle s, a \rangle &\longmapsto \langle a, \langle \check{s}, \check{a} \rangle \rangle \end{aligned}$$

is a dense embedding (see [Go] Definition 0.8):

1. It is easy to see, that h preserves the order relation \leq .
2. Let $\langle p, \langle \check{s}, \check{a} \rangle \rangle \in \mathbf{U} * \mathbf{P}_{\tilde{D}}$. Because \mathbf{U} is \aleph_0 -closed, there is a condition $q \leq p$ and $s \in [\omega]^{<\omega}$, $a \in [\omega]^\omega$ such that $q \Vdash_{\mathbf{U}} \check{s} = \tilde{s} \wedge \check{a} = \tilde{a}$. It is obvious that $\langle q, \langle \check{s}, \check{a} \rangle \rangle \in \mathbf{U} * \mathbf{P}_{\tilde{D}}$ is stronger than $\langle p, \langle \check{s}, \check{a} \rangle \rangle$. Now let $b := q \cap a$, then $h(\langle s, b \rangle) \leq \langle p, \langle \check{s}, \check{a} \rangle \rangle$. \dashv

LEMMA 3.6 $\mathbf{P}_D \approx \mathbf{P}(D)$ *if and only if D is a Ramsey ultrafilter.*

PROOF: See [JS 1] Theorem 1.20. \dashv

LEMMA 3.7 *The Mathias forcing \mathbf{M} is flexible.*

PROOF AND DEFINITION: For the notation see [Je 2] p. 153 and [Ku] p. 224.

A set $T \subseteq \omega^{<\omega}$ is called a *Laver-tree* if

$$T \text{ is a tree and } \exists \tau \in T \forall \sigma \in T (\sigma \subseteq \tau \vee (\tau \subseteq \sigma \wedge |\{n : \sigma \hat{\ } n \in T\}| = \omega)).$$

(We call τ the stem of T . For $\sigma \in T$ we let $\text{succ}_T(\sigma) := \{n : \sigma \hat{\ } n \in T\}$, (the successors of σ in T) and $T_\rho := \{\sigma \in T : \sigma \subseteq \rho \wedge \rho \subseteq \sigma\}$.)

A Laver-tree T is *uniform* if there exists $u_T \in [\omega]^\omega$ such that $\forall \sigma \supseteq \text{stem}(T) (\{n : \sigma \hat{\ } n \in T\} = u_T \setminus (\max(\sigma) + 1))$.

For a Laver-tree T , we say $A \subseteq T$ is a *front* if $\sigma \neq \tau$ in A implies $\sigma \not\subseteq \tau$ and for all $f \in [T]$ there is an $n \in \omega$ such that $f|_n \in A$.

The meaning of $p \leq \llbracket \Phi \rrbracket$ and $p \cap \llbracket \Phi \rrbracket$ are $U_p \subseteq \llbracket \Phi \rrbracket$ and $U_p \cap \llbracket \Phi \rrbracket$, respectively.

1. We say a forcing notion \mathbf{P} is *Laver-like* if there is a \mathbf{P} -name \tilde{r} for a dominating real such that
 - (i) the complete Boolean algebra generated by the family $\{\llbracket \tilde{r}(i) = n \rrbracket : i, n \in \omega\}$ equals r.o. (\mathbf{P}) , and
 - (ii) for each condition $p \in \mathbf{P}$ there exists a Laver-tree $T \subseteq \omega^{<\omega}$ so that

$$\forall \sigma \in T \left(p(T_\sigma) := \prod_{n \in \omega} \sum_{\tau \in T_\sigma} \{p \cap \llbracket \tilde{r}|_{\text{lg}(\tau)} = \tau \rrbracket : \text{lg}(\tau) = n\} \in \text{r.o. } (\mathbf{P}) \setminus \{\mathbf{0}\} \right).$$

We express this by saying $p(T) \neq \emptyset$ where $p(T) := p(T_{\text{stem}(T)})$.

\mathbf{M} is Laver-like:

Let \mathbf{m} be the canonical \mathbf{M} -name for the Mathias real, then \mathbf{m} is dominating (cf. [Je 1] Part I, Lemma 3.15) and further let $p = \langle s, S \rangle \in \mathbf{M}$ with $\text{lg}(s) = n$ and $S = \{a(j) : j \in \omega\}$. Then $U_p = \prod_{k \in n} \llbracket \mathbf{m}(k) = s(k) \rrbracket \cdot \prod_{i \in \omega} \sum_{j \in \omega} \llbracket \mathbf{m}(\check{n} + \check{i}) = a(j) \rrbracket$, which gives a proof of (i).

For (ii) consider $T \subseteq \omega^{<\omega}$ defined as follows:

$$\sigma \in T \text{ iff } \begin{array}{l} \sigma \text{ strictly increasing and} \\ \sigma \subseteq s \vee (s \subseteq \sigma \wedge \text{range}(\sigma) \setminus \text{range}(s) \subseteq S). \end{array}$$

This T has the desired property and is even a uniform Laver-tree. \dashv

2. If \tilde{r} is a \mathbf{P} -name that witnesses that \mathbf{P} is Laver-like, we say that \mathbf{P} has *strong fusion* if for countably many open dense sets $D_n \subseteq \mathbf{P}$ and for $p \in \mathbf{P}$, there is a Laver-tree T such that $p(T) \neq \emptyset$ and for each n :

$$\{\sigma \in T : p(T) \cap \llbracket \tilde{r}|_{\text{lg}(\sigma)} = \sigma \rrbracket \in D_n\}$$

contains a front.

\mathbf{M} has strong fusion:

Let $D \subseteq \mathbf{M}$ be dense open and $p = \langle s, S \rangle$ an \mathbf{M} -condition. For each σ such that $\sigma \subseteq s$ or $(s \subseteq \sigma \wedge \sigma \setminus s \subseteq S)$ we define the rank of σ , $\text{rk}_D(\sigma)$ as follows:

$$\begin{aligned} \text{rk}_D(\sigma) = 0 &\Leftrightarrow \exists A \in [S]^\omega (\langle \sigma, A \rangle \in D), \\ \text{rk}_D(\sigma) = \alpha &\Leftrightarrow \neg \exists \beta < \alpha (\text{rk}_D(\sigma) = \beta) \text{ and} \\ &|\{n : n \in S \wedge \text{rk}_D(\sigma \hat{\ } n) < \alpha\}| = \omega. \end{aligned}$$

If $\text{rk}_D(\sigma)$ is undefined, we put $\text{rk}_D(\sigma) = \infty$.

Note that if $\sigma \in \text{dom}(\text{rk}_D)$, then $\text{rk}_D(\sigma) < \infty$. Otherwise almost all successors (in S) of σ have rank $= \infty$, hence the complement of $S_0 := \{n : n \in S \wedge \text{rk}_D(\sigma \hat{\ } n) = \infty\}$ with respect to S is finite. Let $s_n := \min(S_n)$, then the complement of

$$S_{n+1} := \{n : n \in S_n \wedge \text{rk}_D(\sigma \hat{\ } \tau \hat{\ } s_n \hat{\ } n) = \infty \text{ for all } \tau \in [\{s_0, \dots, s_{n-1}\}]^{<n}\}$$

with respect to S_n is finite. Let $A := \{s_i : i \in \omega\} \subseteq S$ and take $\langle \rho, A' \rangle \leq \langle \sigma, A \rangle$ such that $\langle \rho, A' \rangle \in D$. Then $\rho = \sigma \hat{\ } \tau \hat{\ } s_n$ (for an n) and $A' \in [A]^\omega$, hence $\text{rk}_D(\rho) = \infty$, a contradiction.

For two uniform Laver-trees T and T' , the expression $T \leq_n T'$ means that the first n elements of u_T and $u_{T'}$ are the same. Let T_0 be the uniform Laver-tree constructed in the proof of part (ii) above.

Define a uniform Laver-tree T_{n+1} and the corresponding set $u_{T_{n+1}}$ recursively such that $T_{n+1} \leq_n T_n$ and if $\sigma \in T_{n+1}$ then one of the following is true:

$$\max(\sigma) \leq u_{T_n}(n) \wedge \sigma \in T_n$$

$$(\text{rk}_{D_n}(\sigma) = 0 \wedge \forall k < \text{lg}(\sigma)(\text{rk}_{D_n}(\sigma|_k) > 0)) \rightarrow \langle \sigma, u_{T_{n+1}} \setminus (\max(\sigma) + 1) \rangle \in D_n$$

$$\text{rk}_{D_n}(\sigma) > 0 \wedge \forall k \in u_{T_{n+1}} \setminus (\max(\sigma) + 1)(\text{rk}_{D_n}(\sigma) > \text{rk}_{D_n}(\sigma \hat{\ } k))$$

Now T_{n+1} is a uniform Laver-tree and $T := \bigcap_{n \in \omega} T_n$ is also uniform, $p(T) \neq \emptyset$ and $\{\sigma \in T : p \cap [\mathbf{m}|_{\text{lg}(\sigma)} = \sigma] \in D_n\}$ contains a front, (consider rk_{D_n}). \dashv

3. A Laver-like \mathbf{P} is *closed under finite changes* if given a $p \in \mathbf{P}$ and Laver trees T and T' so that for all $\sigma \in T' : |\text{succ}_T(\sigma) \setminus \text{succ}_{T'}(\sigma)| < \omega$, if $p(T) \neq \emptyset$, then $p(T') \neq \emptyset$, too.

\mathbf{M} is closed under finite changes:

Use a standard fusion argument. \dashv

4. We call \mathbf{P} a *flexible* forcing notion iff \mathbf{P} is Laver-like, has strong fusion and is closed under finite changes.

Hence, the Mathias forcing \mathbf{M} is flexible. \dashv

Essential Theorems about Δ_2^1 -sets of reals

Now we will give the relationship between the Ramsey property and Mathias forcing.

FACTS 3.8

1. [JS 1] Theorem 1.7:

For every $\mathbf{P}(D)$ -sentence Φ and for all $p \in \mathbf{P}(D)$ there exists a $q \in \mathbf{P}(D)$ such that $q \leq p$, $\text{stem}(p) = \text{stem}(q)$ and

$$q \Vdash_{\mathbf{P}(D)} \Phi \text{ or } q \Vdash_{\mathbf{P}(D)} \neg \Phi \quad (q \text{ decides } \Phi).$$

(This is known as pure decision.)

2. [JS 1] Theorem 1.14:

If $V \subseteq V' \subseteq V''$ are models of ZFC and $D \in V$ is an ultrafilter and $x \in V'$ is $\mathbf{P}(D)$ -generic over V , then for every $y \in [x]^\omega \cap V''$, y is $\mathbf{P}(D)$ -generic over V , too.

3. [JS 1] Theorem 1.15:

If $D \in V$ and g is $\mathbf{P}(D)$ -generic over V , then

$$V[g] \models \Sigma_2^1(\mathcal{R})_V.$$

4. [JS 1] Theorem 1.16:

If $D \in V$, then

$r \in [\omega]^\omega$ is $\mathbf{P}(D)$ -generic over V if and only if

$$\forall a \in D(r \subseteq^* a) \text{ and } \forall \pi \in 2^{[\omega]^2} \cap V : \exists n \in \omega \text{ such that } \pi|_{[r \setminus n]^2} \text{ is constant.}$$

5. [JS 1] Theorem 2.7:

$$V \models \Delta_2^1(\mathcal{R}) \text{ if and only if } V \models \Sigma_2^1(\mathcal{R}).$$

6. [JS 1] Theorem 2.11:

For an $s \in [\omega]^\omega$ define $D_s := \{a \in [\omega]^\omega : s \subseteq^* a\}$ (where $s \subseteq^* a$ means $|s \setminus a| < \omega$) and $D^s := D_s \cap L[D_s]$. If D^s is an ultrafilter in $L[D^s]$ and $r \subseteq^* s$, then $D^r = D^s$ and we write \mathbf{P}_s for the forcing notion $\mathbf{P}(D^s)$ in $L[D^s]$.

$$V \models \Delta_2^1(\mathcal{R})_{L[u]} \Leftrightarrow \forall r \in L[u] \exists s \in [r]^\omega \cap V (s \text{ is } \mathbf{P}_s\text{-generic over } L[u][D^s]).$$

Some properties of Mathias forcing

FACTS 3.9

1. Using the Fact 3.8 3. and the Lemmas 3.5 and 3.6 we see that if m is Mathias over V , then $V[m] \models \Sigma_2^1(\mathcal{R})_V$. Thus, (with [Ku] Lemma 5.14 on p.276) an ω_1 -iteration of Mathias forcing with countable support gives a model in which each Σ_2^1 -set is Ramsey.

2. We call r a *Ramsey real over V* if and only if there exists a $D \in V$ such that:

(i) D is an ultrafilter, $\forall a \in D(r \subseteq^* a)$ and

(ii) for all $\pi \in 2^{[\omega]^2}$, $\pi \in V$ there is an $n \in \omega$ such that $\pi|_{[r \setminus n]^2}$ is constant.

(See also [JS 1] Definition 1.17).

Now we see that if s is \mathbf{P}_s -generic over $L[u][D^s]$, then (by 3.8 4.) it is Ramsey over $L[u][D^s]$ and even a dominating real with respect to $L[u][D^s]$.

PROOF: To each real $r \in L[u][D^s]$ consider the function $\pi_r \in 2^{[\omega]^2}$ (which also belongs to $L[u][D^s]$) defined as follows:

$$\pi_r(\{i, j\}) = 0 \iff \exists k(r(2^k) < i, j \leq r(2^{k+1})).$$

Because s is \mathbf{P}_s -generic and by 3.8 4. we have

$$\exists n \in \omega (\pi|_{[s \setminus n]^2} \text{ is constant}).$$

Thus, because $s \setminus n$ is infinite, $\pi|_{[s \setminus n]^2} \equiv 1$ and for $k \geq 2n$ we get $s(k) > r(k)$, hence

$$\forall r \in L[u][D^s] \cap [\omega]^\omega \exists l \in \omega \forall k \geq l (s(k) > r(k))$$

which says, that the reals of $L[u][D^s]$ are dominated by s . \dashv

We close this section by mentioning two useful corollaries.

COROLLARY 3.10 *If p is an \mathbf{M} -condition and \tilde{x} is an \mathbf{M} -name for a real, then there exists an \mathbf{M} -condition $q \leq^0 p$ and a real $\bar{x} \in V$ such that $V \models "q \Vdash_{\mathbf{M}} \bar{x} = \tilde{x}"$.*

PROOF: Let \tilde{x} be an \mathbf{M} -name for a real. Each real can be considered as an infinite 0-1-sequence, so \tilde{x} is such that for all natural numbers n :

$$\tilde{x}(\check{n}) = \check{1} \quad \text{or} \quad \tilde{x}(\check{n}) = \check{0}.$$

Take $p = \langle s, X \rangle$. Because Mathias forcing has pure decision (by the Lemmas 3.5, 3.6 and Fact 3.8 1., or by [Ba] Theorem 9.3) in V there is a condition $\langle s, X_0 \rangle$ such that $X_0 \subseteq X$ which decides $\tilde{x}(\check{0})$. Let a_0 be the least member of X_0 , then there are Y, X_1 such that $X_0 \setminus \{a_0\} \supseteq Y \supseteq X_1$ and $\langle s \hat{\ } a_0, Y \rangle, \langle s, X_1 \rangle$ both decide $\tilde{x}(\check{1})$. Let now a_1 be the least member of X_1 . There are Y_1, Y_2, Y_3, X_2 such that $X_1 \setminus \{a_1\} \supseteq Y_1 \supseteq \dots \supseteq X_2$ and $\langle s \hat{\ } a_0 \hat{\ } a_1, Y_1 \rangle, \langle s \hat{\ } a_1, Y_2 \rangle, \langle s \hat{\ } a_0, Y_3 \rangle, \langle s, X_2 \rangle$ all decide $\tilde{x}(\check{2})$. Now let a_2 be the least member of X_2 and so on. Define $r := \{a_i : i \in \omega\}$. We encode now \tilde{x} by $\bar{x} := \{s \hat{\ } t : t \in [r]^{<\omega} \wedge \langle s \hat{\ } t, r \setminus (\max(t) + 1) \rangle \Vdash_{\mathbf{M}} \tilde{x}(\lg(t))^\vee = \check{1}\}$. Then \bar{x} is a real and if m is a Mathias real over V such that $\langle s, r \rangle \in m$ then $\tilde{x}[m] = \bar{x}[m]$, where $\bar{x}[m](n) = 1$ if and only if $m|_n \in \bar{x}$. \dashv

COROLLARY 3.11 *If p is an \mathbf{M} -condition and $V \models "p \Vdash_{\mathbf{M}} \exists x \Phi(x)"$, then there is an \mathbf{M} -condition $q \leq^0 p$ and an \mathbf{M} -name \tilde{x} for a real such that $V \models "q \Vdash_{\mathbf{M}} \Phi(\tilde{x})"$.*

PROOF: We will follow the proofs of [Ba] Theorems 9.1 and 9.3.

Assume $p = \langle s, A \rangle \Vdash_{\mathbf{M}} \exists x \Phi(x)$. First we prove that there is a $B \subseteq A$ such that if $\langle t, C \rangle \leq \langle s, B \rangle$, \tilde{x} an \mathbf{M} -name and $\langle t, C \rangle \Vdash_{\mathbf{M}} \Phi(\tilde{x})$, then we find an \mathbf{M} -name \tilde{y} such that $\langle t, B \setminus (\max(t) + 1) \rangle \Vdash_{\mathbf{M}} \Phi(\tilde{y})$. For this we construct a sequence $b_0 < b_1 < \dots$ of elements of A and a sequence $B_0 \supseteq B_1 \supseteq \dots$ of subsets of A such that for all $b \in B_{n+1}$, $b_n < b$. Let $B_0 := A$. Given B_n , let s_1, s_2, \dots, s_k enumerate all the subsets of $\{b_i : i < n\}$. Now construct a sequence $B_n^0 \supseteq B_n^1 \supseteq \dots \supseteq B_n^k$ as follows. $B_n^0 := B_n$ and given B_n^{i-1} let $B_n^i \subseteq B_n^{i-1}$ be such that for some \mathbf{M} -name \tilde{x} , $\langle s \cup s_i, B_n^i \rangle \Vdash_{\mathbf{M}} \Phi(\tilde{x})$, if it exists; otherwise let $B_n^i := B_n^{i-1}$. Finally let $b_n := \bigcap B_n^k$, $B_{n+1} := B_n^k \setminus \{b_n\}$ and $B := \{b_n : n \in \omega\}$. Suppose $\langle t, C \rangle \leq \langle s, B \rangle$ and we find an \mathbf{M} -name \tilde{x} such that $\langle t, C \rangle \Vdash_{\mathbf{M}} \Phi(\tilde{x})$. Because there is an $n \in \omega$ such that $s_t := t \setminus s \subseteq \{b_i : i < n\}$ we must have chosen B_n^l so that for some \mathbf{M} -name \tilde{y} , $\langle s \cup s_t, B_n^l \rangle \Vdash_{\mathbf{M}} \Phi(\tilde{y})$. Now $B \setminus (\max(t) + 1) \subseteq B_n^l$, hence $\langle t, B \setminus (\max(t) + 1) \rangle \leq \langle t, B_n^l \rangle \Vdash_{\mathbf{M}} \Phi(\tilde{y})$ and we are done.

If p, q are two \mathbf{M} -conditions, then $p \cap q$ denotes the weakest \mathbf{M} -condition which is stronger than p and q , (if it exists). Let \tilde{x} be an \mathbf{M} -name and p an \mathbf{M} -condition, then $\tilde{x}(p)$ denotes the following name. $\langle \tilde{\sigma}, q \rangle \in \tilde{x}(p)$ if and only if there exists an \mathbf{M} -condition q' such that $\langle \tilde{\sigma}, q' \rangle \in \tilde{x}$ and $q = p \cap q'$. For two \mathbf{M} -names \tilde{x}, \tilde{y} let $\tilde{x} \cup \tilde{y} := \{\langle \tilde{\sigma}, p \rangle : \langle \tilde{\sigma}, p \rangle \in \tilde{x} \vee \langle \tilde{\sigma}, p \rangle \in \tilde{y}\}$. Now we are prepared to prove the corollary. Given $p = \langle s, A \rangle \Vdash_{\mathbf{M}} \exists x \Phi(x)$. Let $B \subseteq A$ be as above. We construct a sequence $b_0 < b_1 < \dots$ of elements of B and subsets $B_0 \supseteq B_1 \supseteq \dots$ of B by induction as follows. Let $B_0 := B$. Given B_n , find $B_{n+1} \subseteq B_n$ so that for all $s' \subseteq \{b_i : i < n\}$ one of the following cases holds:

1. For **all** $b \in B'_{n+1}$ we find an \mathbf{M} -name \tilde{x} (depending on b) such that $\langle s \cup s' \cup \{b\}, B'_{n+1} \setminus (b+1) \rangle \Vdash_{\mathbf{M}} \Phi(\tilde{x})$.
2. For **no** $b \in B'_{n+1}$ we find an \mathbf{M} -name \tilde{x} (which may depend on b) such that $\langle s \cup s' \cup \{b\}, B'_{n+1} \setminus (b+1) \rangle \Vdash_{\mathbf{M}} \Phi(\tilde{x})$.

Because of the choice of B , for each n we find a $B'_{n+1} \subseteq B_n \subseteq B$. Let $b_n := \bigcap B'_{n+1}$, $B_{n+1} := B'_{n+1} \setminus \{b_n\}$ and $A' := \{b_n : n \in \omega\}$. Suppose for $\langle t, C \rangle \leq \langle s, A' \rangle$ we find an \mathbf{M} -name \tilde{x} , such that $\langle t, C \rangle \Vdash_{\mathbf{M}} \Phi(\tilde{x})$. Let $|t|$ be minimal. If $|t| = |s|$ then $t = s$ and we find an \mathbf{M} -name \tilde{y} such that $\langle s, A' \rangle \Vdash_{\mathbf{M}} \Phi(\tilde{y})$. If $|t| > |s|$ then $\max(t) = b_n$ for some n and at stage n , the first case held for some $s' = t \setminus (s \cup \{b_n\})$. Now for each $b_i \in A'$ ($i \geq n$) take an \mathbf{M} -name \tilde{x}_i such that $\langle s \cup s' \cup \{b_i\}, A' \setminus (b_i+1) \rangle \Vdash_{\mathbf{M}} \Phi(\tilde{x}_i)$. Further let $\tilde{y} := \bigcup \{\tilde{x}_i(p_i) : i \geq n \wedge p_i = \langle s \cup s' \cup \{b_i\}, A' \setminus (b_i+1) \rangle\}$. Then we have $\langle s \cup s', A' \setminus (\max(s')+1) \rangle \Vdash_{\mathbf{M}} \Phi(\tilde{y})$, which is a contradiction to the minimality of $|t|$. \dashv

In the next section we start to show the relationship between Mathias-absoluteness and the Ramsey property of projective sets of reals.

4 Σ_2^1 -sets and the Ramsey Property

It is well-known that for $\Sigma_2^1(\mathcal{B})$ and $\Sigma_2^1(\mathcal{L})$ there are characterizations with forcing absoluteness (cf. Theorem 2.3). Such a characterization exists also for $\Sigma_2^1(\mathcal{R})$. Although the proofs for the Baire property and the Lebesgue measurability are similar, the proof for the Ramsey property is different. This is because Mathias forcing does not have the countable chain condition. (But fortunately it has a lot of combinatorial properties.)

THEOREM 4.1 $V \models \Sigma_2^1(\mathcal{R})$ if and only if V is Σ_3^1 -Mathias-absolute.

PROOF: First we prove that Σ_3^1 - \mathbf{M} -absoluteness implies $\Sigma_2^1(\mathcal{R})$. For this let $\Phi(x)$ be a Δ_2^1 -set:

$$\Phi(x) \leftrightarrow \varphi(x) \leftrightarrow \psi(x),$$

where $\varphi(x)$ is a Σ_2^1 -set and $\psi(x)$ is a Π_2^1 -set. Because $\forall x(\varphi(x) \leftrightarrow \psi(x))$ is a Π_3^1 -sentence, by Σ_3^1 - \mathbf{M} -absoluteness we have

$$V^{\mathbf{M}} \models \forall \tilde{x}(\varphi(\tilde{x}) \leftrightarrow \psi(\tilde{x})).$$

By Fact 3.9 1. we know that $V^{\mathbf{M}} \models$ “each Δ_2^1 -set with parameters in V is Ramsey”. Therefore

$$V^{\mathbf{M}} \models \exists \tilde{y}(\forall \tilde{x}_0(\tilde{x}_0 \in [\tilde{y}]^\omega \rightarrow \psi(\tilde{x}_0)) \vee \forall \tilde{x}_1(\tilde{x}_1 \in [\tilde{y}]^\omega \rightarrow \neg\varphi(\tilde{x}_1))).$$

But this is a Σ_3^1 -sentence and because $V \models \varphi(x) \leftrightarrow \psi(x)$, also $V \models$ “ Φ is Ramsey”. Now because $\Phi(x)$ was arbitrary and $\Sigma_2^1(\mathcal{R})$ is equivalent to $\Delta_2^1(\mathcal{R})$ (by Fact 3.8 5.), we have $V \models \Sigma_2^1(\mathcal{R})$.

Now we prove that $\Sigma_2^1(\mathcal{R})$ implies Σ_3^1 - \mathbf{M} -absoluteness. Let $\Psi = \exists x\psi(x)$ be a Σ_3^1 -sentence. If $V \models \Psi$, then by the Shoenfield absoluteness Lemma (see [Je 2] Theorem 98), the Σ_3^1 -sentences are upwards absolute, hence $V^{\mathbf{M}} \models \Psi$. For the other direction assume that $V^{\mathbf{M}} \models \Psi$. Then, because of $V^{\mathbf{M}}$ is full (cf. [Je 2] Lemma 18.6), there is a name \tilde{x} , such that $V^{\mathbf{M}} \models \psi(\tilde{x})$. By Corollary 3.10 there exist reals $r, \bar{x} \in V$ such that $\bar{x} \subseteq r$ and $V \models$ “ $r \Vdash_{\mathbf{M}} \bar{x} = \tilde{x}$ ”. Now, because $V \models \Sigma_2^1(\mathcal{R})$, there is an $s \in [r]^\omega$ such that s is \mathbf{P}_s -generic over $L[\bar{x}][r][D^s]$. Let $m \subseteq s$ be a Mathias real over V , then m is also \mathbf{P}_s -generic over

$L[\bar{x}][r][D^s]$ (by 3.8 2.). $V[m] \models \psi(\bar{x}[m])$, hence $L[\bar{x}][r][D^s][m] \models \psi(\bar{x}[m])$ because ψ is Π_2^1 , $m \subseteq s \subseteq r$ and \bar{x} may be regarded also as a \mathbf{P}_s -name. So there must be a condition $p \in L[\bar{x}][r][D^s]$ such that $L[\bar{x}][r][D^s] \models "p \Vdash_{\mathbf{P}_s} \psi(\bar{x})"$. Let $k = \max(\text{range}(\text{stem}(p)))$, then $s' := s \setminus k$ is \mathbf{P}_s -generic over $L[\bar{x}][r][D^s]$ and there is an $n \in \omega$ such that $s'' := (s' \setminus n) \cup \text{stem}(p)$ satisfies p , (by [JS 1] Definition 1.8 and Lemma 1.12). Hence (again by [JS 1] Lemma 1.12), s'' is \mathbf{P}_s -generic over $L[\bar{x}][r][D^s]$ and because of s'' satisfies p and $p \Vdash_{\mathbf{P}_s} \psi(\bar{x})$ we have $L[\bar{x}][r][D^s][s''] \models \psi(\bar{x}[s''])$ and finally $V \models \exists x \psi(x)$, (by Shoenfield). \dashv

So, we have found a forcing characterization for $\Sigma_2^1(\mathcal{R})$. Such a characterization with Mathias forcing does not exist for higher degrees of Mathias-absoluteness as we will show in the next section.

5 Σ_4^1 -M-absoluteness and the Ramsey Property

THEOREM 5.1 Σ_4^1 -Mathias-absoluteness implies $\Delta_3^1(\mathcal{R})$.

PROOF: Assume that V is Σ_4^1 - \mathbf{M} -absolute. Let $\Phi(x)$ be a Δ_3^1 -set in V with parameters in V : $\Phi(x) \leftrightarrow \varphi(x) \leftrightarrow \psi(x)$ where $\varphi(x)$ is a Σ_3^1 -set and $\psi(x)$ is a Π_3^1 -set. So $V \models \forall x(\varphi(x) \leftrightarrow \psi(x))$ and $\forall x((\varphi(x) \vee \neg\psi(x)) \wedge (\neg\varphi(x) \vee \psi(x)))$ is a Π_4^1 -sentence, hence \mathbf{M} -absolute. Therefore $\Phi(x)$ is still a Δ_3^1 -set in $V^{\mathbf{M}}$.

Assume $V \models "\Phi(x)$ is not Ramsey". Hence $V \models \forall x \exists y_1 y_2 (y_1 \subseteq x \wedge y_2 \subseteq x \wedge \Phi(y_1) \wedge \neg\Phi(y_2))$. Obviously we have $\Phi(y_1)$ iff $\varphi(y_1)$ and $\neg\Phi(y_2)$ iff $\neg\psi(y_2)$ but $\varphi(x)$, $\neg\psi(x)$ are both Σ_3^1 -sets. So $V \models "\Phi(x)$ is not Ramsey" is equivalent to

$$V \models \forall x \exists y_1 y_2 (y_1 \subseteq x \wedge y_2 \subseteq x \wedge \varphi(y_1) \wedge \neg\psi(y_2)) \quad (\equiv: \Theta)$$

where Θ is a Π_4^1 -sentence. Thus by Σ_4^1 - \mathbf{M} -absoluteness we have

$$V^{\mathbf{M}} \models \Theta. \quad (*)$$

Let \mathbf{m} be the canonical name for a Mathias real m over V . Then there is a condition p with empty stem such that $p \Vdash_{\mathbf{M}} \varphi(\mathbf{m})$ or $p \Vdash_{\mathbf{M}} \neg\varphi(\mathbf{m})$, (see Lemmas 3.5 and 3.6 and Fact 3.8 1.). Assume $p \Vdash_{\mathbf{M}} \varphi(\mathbf{m})$, then $p \Vdash_{\mathbf{M}} \exists \tilde{x} \varphi(\tilde{x})$ (otherwise $p \Vdash_{\mathbf{M}} \neg\psi(\mathbf{m})$ and $\neg\psi(\mathbf{m})$ is also Σ_3^1). Because each $y \in [m]^\omega$ is Mathias over V and $\text{stem}(p) = \langle \rangle$ we have $V[y] \models \varphi(y)$. Because $V[y] \subseteq V[m]$ and φ is Σ_3^1 , hence upwards absolute, $V[m]$ is also a model of $\varphi(y)$. So, we get

$$p \Vdash_{\mathbf{M}} \exists \tilde{x} \forall \tilde{y} (\tilde{y} \in [\tilde{x}]^\omega \rightarrow \varphi(\tilde{y})).$$

Now because $V^{\mathbf{M}} \models \forall \tilde{x} (\varphi(\tilde{x}) \leftrightarrow \psi(\tilde{x}))$ we finally have

$$p \Vdash_{\mathbf{M}} \exists \tilde{x} \forall \tilde{y} (\tilde{y} \in [\tilde{x}]^\omega \rightarrow \varphi(\tilde{y}) \wedge \psi(\tilde{y})),$$

but this is a contradiction to (*). \dashv

THEOREM 5.2 $\Delta_3^1(\mathcal{R})$ does not imply Σ_4^1 -Mathias-absoluteness.

PROOF: For this it is enough to find a model V in which all Δ_3^1 -sets are Ramsey, all Δ_2^1 -sets have the property of Baire and ω_1 in this model is the same as ω_1^L .

We have $V \models \Delta_2^1(\mathcal{B})$ if and only if for all reals r in V there is a real in V which is Cohen over $L[r]$. To say this is a Π_4^1 -sentence: For $s \in 2^{<\omega}$ consider $1 \hat{\ } s$ as a binary code for a natural number n ($n > 0$) and let $\#n := s$, ($\#0 := \#1 = \langle \rangle$). We write $n \preceq m$ if $\#m|_{\text{lg}(\#n)} = \#n$. Note that $\#n \subseteq \#m$ is an arithmetical statement. The sentence $\forall r \in [\omega]^\omega \exists c \in [\omega]^\omega \forall x \in [\omega]^\omega (c \text{ is a branch} \wedge (x \in L[r] \wedge x \text{ encodes a dense set} \rightarrow x \cap c \neq \emptyset))$ is a composition of the following sentences.

c is a branch is $\forall nm((n \in c \wedge m \in c) \rightarrow (n \preceq m \vee m \preceq n))$, which is an arithmetical statement.

$x \in L[r]$ is a Σ_2^1 -sentence with parameter r (cf. [Je 2] Theorem 97).

x encode a dense set is $\forall m \exists n(n \in x \wedge m \preceq n)$, which is arithmetic.

Finally $x \cap c \neq \emptyset$ is $\exists t(t \in x \wedge t \in c)$, which is arithmetic, too. \dashv

So, if V is a model with the desired properties and V is Σ_4^1 - \mathbf{M} -absolute, for each real $r \in V[m]$ there is (in $V[m]$) a Cohen real c over $L[r]$. If $r \in V[m]$ is a real and c is a Cohen real over $L[r]$, then $L[r] \cap \omega^\omega$ is a strong measure zero set in $L[r][c]$ (see [Co] Theorem 1.3) and hence we find in $V[m]$ a covering of $L \cap \omega^\omega$ with respect to the real r . So $L \cap \omega^\omega$ is a strong measure zero set belonging to V .

Now if $\omega_1^L = \omega_1^V$ then we get in $V[m]$ a strong measure zero set of cardinality ω_1 with parameter in V , namely $L \cap \omega^\omega$, but this is a contradiction, (cf. [Ba], proof of Theorem 9.7 or cf. [Je 1] Lemma 8.2 and recall that $\mathbf{M} \approx \mathbf{U} * \mathbf{P}_{\bar{D}} \approx \mathbf{U} * \mathbf{P}(\bar{D})$).

It leaves to construct a model V with the desired properties.

In [JS 2] §3 they show, that an ω_1 iteration of Mathias forcing starting from L , yields a model in which every Δ_3^1 -set is Ramsey and ω_1 in this model is the same as ω_1^L . (By the claim of Theorem 5.3 this is already enough).

Now in [Ju 3] Lemma 1.18 it is proved that if we make a suitable ω_1 iteration starting from L , and add alternately Kesef and Cohen reals, we get a model V in which every Δ_3^1 -set is Ramsey, every Δ_2^1 -set has the Baire property and ω_1^V is the same as ω_1^L . \dashv

The next theorem is in fact a consequence of the following: If V is Σ_4^1 - \mathbf{M} -absolute, then ω_1^V is inaccessible in L .

THEOREM 5.3 Σ_4^1 -Mathias-absoluteness implies $\Sigma_3^1(\mathcal{R})$.

PROOF: We first give the following

CLAIM: If V is Σ_4^1 - \mathbf{M} -absolute, then for all reals $r \in V$ we have $\omega_1^{L[r]} < \omega_1^V$, (hence ω_1^V is inaccessible in L).

Now we show that this claim implies that $V \models \Sigma_3^1(\mathcal{R})$.

LEMMA 5.4 If V is Σ_4^1 - \mathbf{M} -absolute and $\forall r \in V(\omega_1^{L[r]} < \omega_1^V)$ then $V \models \Sigma_3^1(\mathcal{R})$.

PROOF OF THE LEMMA: Let $\Phi(x) \equiv \exists y \psi(x, y)$ be a Σ_3^1 -set with parameter $a \in V$. If $V \models \exists z \forall x(x \in [z]^\omega \rightarrow \neg \Phi(x))$, then the set $\Phi(x)$ is Ramsey in V . Therefore let us assume that $V \models \forall z \exists x(x \in [z]^\omega \wedge \Phi(x))$ ($\equiv: \Theta$). Because Θ is a Π_4^1 -sentence with parameter a and by Σ_4^1 - \mathbf{M} -absoluteness we have $V^{\mathbf{M}} \models \Theta$. Now there is a Mathias condition p with empty stem, such that p decides $\Phi(\mathbf{m})$. Because $V^{\mathbf{M}} \models \Theta$, $V^{\mathbf{M}}$ is a model of $\exists \tilde{x}(\tilde{x} \in [\mathbf{m}]^\omega \wedge \Phi(\tilde{x}))$. Further $V^{\mathbf{M}}$ is full and $\Phi(x) \equiv \exists y \psi(x, y)$, hence we find Mathias names \tilde{x}, \tilde{y} such that $V^{\mathbf{M}} \models (\tilde{x} \in [\mathbf{m}]^\omega \wedge \psi(\tilde{x}, \tilde{y}))$.

Consider the statement $V[m] \models \exists y \psi(x, y) \Leftrightarrow V[x] \models \exists z \psi(x, z)$ and further assume that $V \models "q \Vdash_{\mathbf{M}} \psi(\tilde{x}, \tilde{y}) \wedge V[\tilde{x}] \not\models \exists \tilde{z} \psi(\tilde{x}, \tilde{z})"$ (for an \mathbf{M} -condition q). First we have to define the meaning of $q \Vdash_{\mathbf{M}} "V[\tilde{x}] \models \Psi(\tilde{x})"$ where Ψ is an arbitrary formula with at most one free variable: If \tilde{z} is a variable in Ψ for a real and $\tilde{n} \in \tilde{z}$ is a subformula of Ψ , then $q_0 \Vdash_{\mathbf{M}} "V[\tilde{x}] \models \tilde{n} \in \tilde{z}"$ if and only if there exists a Mathias condition $\langle u, U \rangle$ such that

$$\langle u, U \rangle \Vdash_{\mathbf{M}} \tilde{n} \in \tilde{z} \quad \text{and} \quad q_0 \Vdash_{\mathbf{M}} \forall \check{k}(\check{k} \in \check{u} \rightarrow \check{k} \in \tilde{x} \wedge \check{k} \in \tilde{x} \rightarrow (\check{k} \in \check{U} \vee \check{k} \in \check{u})).$$

Let x be the evaluation of \tilde{x} by the Mathias real m . Now because $V^{\mathbf{M}} \models \tilde{x} \subseteq \mathbf{m}$, x is also Mathias over V and $V[x] \models \Psi(x)$ if and only if there exists a Mathias condition $q_0 \in V$

such that $q_0 \in m$ and $q_0 \Vdash_{\mathbf{M}} V[\tilde{x}] \models \Psi(\tilde{x})$. Thus “ $q \Vdash_{\mathbf{M}} V[\tilde{x}] \models \Psi(\tilde{x})$ ” is well defined. Let $r, \bar{x}, \bar{y} \in V$ be such that $r \leq q$ and $V \models “r \Vdash_{\mathbf{M}} \bar{x} = \tilde{x} \wedge \bar{y} = \tilde{y}”$. Further let $r \in s \in V$ be Ramsey over $L[a][r][\bar{x}, \bar{y}]$, then there is a condition $p_0 \in L[a][r][\bar{x}, \bar{y}][D^s]$, $p_0 \leq r$ such that $L[a][r][\bar{x}, \bar{y}][D^s] \models “p_0 \Vdash_{\mathbf{P}_s} \psi(\bar{x}, \bar{y})”$. This is because if $m' \leq s$ is Mathias over V , then m' is \mathbf{P}_s -generic and $L[a][r][\bar{x}, \bar{y}][D^s][m'] \models \psi(\bar{x}[m'], \bar{y}[m'])$ (by Shoenfield). Let s' be \mathbf{P}_s -generic such that $p_0 \in s'$, further let $s' \in m$ be Mathias over V and $x := \bar{x}[m]$ ($= \tilde{x}[m]$). We write \mathbf{P}_s as a two step iteration $Q_1 * \tilde{Q}_2$ and choose g_1 such that g_1 is Q_1 -generic over $L[a][r][\bar{x}, \bar{y}][D^s]$ ($=: N$) and $N[g_1] = N[x]$. Because of $N[x] \subseteq V[x]$, $V[x] \cap [\omega]^\omega \cap N[x]$ is a Σ_2^1 -set in $V[x]$ and $\forall x \exists y \forall z (z \in N[x] \rightarrow \exists n (y_n = z))$ (this is: for all x , $\omega_1^{N[x]}$ is countable) is a Π_4^1 -sentence. Because of $x \subseteq m$ is Mathias over V and V is Σ_4^1 - \mathbf{M} -absolute, it follows that $\omega_1^{V[x]}$ is inaccessible in $N[x]$. Hence, there exists a set $g_2 \in V[x]$ which is $\tilde{Q}_2[x]$ -generic over $N[x]$ such that $N[x][g_2] \models \psi(x, \bar{y}[g_1 * g_2])$. Now $N[x][g_2] \models \exists y \psi(x, y)$ and $N[x][g_2] \subseteq V[x]$ and because Σ_3^1 -formulas are upwards absolute, $V[x] \models \exists y \psi(x, y)$, which is a contradiction to $q \Vdash_{\mathbf{M}} “V[\tilde{x}] \not\models \exists \tilde{z} \psi(\tilde{x}, \tilde{z})”$.

(If m is Mathias over V and $x \in [m]^\omega \cap V[m]$, then we say that V is Σ_n^1 - \mathbf{M} -correct if for every Σ_n^1 -set $\Phi(x)$ with parameters in V : $V[m] \models \Phi(x) \Leftrightarrow V[x] \models \Phi(x)$.)

Let p be a Mathias condition with empty stem which decides $\Phi(\mathbf{m})$, where m is Mathias over V . Thus

$$V \models “p \Vdash_{\mathbf{M}} \exists \tilde{z} \psi(\tilde{z}, \mathbf{m})” \quad \text{or} \quad V \models “p \Vdash_{\mathbf{M}} \neg \Phi(\mathbf{m})”.$$

If the first case holds, let r, \bar{z} be such that: $r \subseteq p$ and if $m \subseteq r$ is Mathias over V , then $V[m] \models \psi(\bar{z}[m], m)$. In V there exists a Ramsey real $s \subseteq r$ over $L[a][r][\bar{z}]$ and because Π_2^1 -sets are absolute (by Shoenfield) in $L[a][r][\bar{z}]$ there exists a \mathbf{P}_s -condition q with empty stem (note that all $t \in [s]^\omega$ are also Ramsey over $L[a][r][\bar{z}]$) such that $L[a][r][\bar{z}][D^s] \models “q \Vdash_{\mathbf{P}_s} \psi(\bar{z}, \mathbf{g})”$ where \mathbf{g} is the canonical name for the \mathbf{P}_s -generic real over $L[a][r][\bar{z}][D^s]$. In V there is a \mathbf{P}_s -generic real s' such that $s' \subseteq q$, hence for all $t \in [s']^\omega$: $L[a][r][\bar{z}][D^s][t] \models \psi(\bar{z}[t], t)$. Again by Shoenfield we get:

$$V \models \psi(\bar{z}[t], t) \text{ and this implies } V \models \exists y \forall x \in [y]^\omega \Phi(x).$$

Therefore the set $\Phi(x)$ is Ramsey in V .

If the second case holds, we get

$$V \models “p \Vdash_{\mathbf{M}} \forall \tilde{x} \in [\mathbf{m}]^\omega \neg \Phi(\tilde{x})”$$

hence $V \models “p \Vdash_{\mathbf{M}} \exists \tilde{y} \forall \tilde{x} \in [\tilde{y}]^\omega \neg \Phi(\tilde{x})”$ which is a Σ_4^1 -sentence (with parameters in V) and says, that $\Phi(x)$ is Ramsey. Therefore by Σ_4^1 - \mathbf{M} -absoluteness the set $\Phi(x)$ has to be Ramsey in V . \dashv

Now we have to show that the claim holds.

PROOF OF THE CLAIM: Assume V is Σ_4^1 - \mathbf{M} -absolute, then by Theorem 4.1 $V \models \Sigma_2^1(\mathcal{R})$, and by the Facts 3.8 5., 3.8 6. and 3.9 2. the following is true in V :

$$\forall u \in [\omega]^\omega \forall r \in L[u] \cap [\omega]^\omega \exists s \in [r]^\omega (s \text{ is Ramsey over } L[u][D^s]).$$

To say this is a Π_4^1 -sentence:

$$\begin{array}{ll} \text{Define } \flat : & [\omega]^2 \quad \longrightarrow \quad \omega \\ & \{n, m\} \quad \longmapsto \quad \frac{1}{2}(\max(\{n, m\})^2 - \max(\{n, m\})) + \min(\{n, m\}). \end{array}$$

Note that \flat is a bijection and arithmetic. With \flat we can consider each $\pi \in [\omega]^\omega$ as a function from $[\omega]^2$ to 2, namely by

$$\pi(\{n, m\}) = 0 \iff \flat\{n, m\} \in \pi.$$

The sentence

$$\forall u \in [\omega]^\omega \forall r \in L[u] \cap [\omega]^\omega \exists s \in [r]^\omega (s \text{ is } \mathbf{P}_s\text{-generic over } L[u][D^s])$$

is a composition of the following sentences.

$r \in L[u]$ is a Σ_2^1 -sentence with parameter u .

$s \in [r]^\omega$ is $\forall i(i \in s \rightarrow i \in r)$, which is arithmetic.

s is \mathbf{P}_s -generic over $L[u][D^s]$, which is again a composition of the following sentences.

$x \in L[u][D^s]$ is a Σ_2^1 -sentence with parameters u and s .

$\pi|_{[s \setminus n]^2}$ is constant is an arithmetical sentence because of \flat is arithmetic.

D^s is an ultrafilter in $L[u][D^s]$ is $\forall x \in L[u][D^s] \cap [\omega]^\omega \exists n(s \setminus n \subseteq x \vee s \setminus n \cap x = \emptyset)$, which is a Π_2^1 -sentence with parameters u and s .

$\forall \pi \in L[u][D^s] \exists n(\pi|_{[s \setminus n]^2} \text{ is constant})$, which is also a Π_2^1 -sentence with the parameters u and s . \dashv

Therefore if V is Σ_4^1 - \mathbf{M} -absolute, in $V^{\mathbf{M}}$ for each real u there exists a real s which dominates the reals of $L[u]$ (cf. Fact 3.9 2.). Let m be Mathias over V . Because \mathbf{M} is flexible (cf. Lemma 3.7), \mathbf{M} adds a dominating family of size ω_1 (see [Br] Theorem 3.1). If there is a real $r \in V$ such that $\omega_1^{L[r]} = \omega_1^V$ and m is Mathias over V , then the reals of $L[r][m]$ dominates the reals of $V[m]$. (Note that the \mathbf{M} -names \check{f}_α ($\alpha < \omega_1$) which are constructed in [Br] Theorem 3.1 can all be defined within $L[r]$.) But this contradicts that in $V[m]$ we have a dominating real over $L[r][m]$. \dashv

This concludes the proof of the Theorem. \dashv

We can prove even more, as we will see in the next section.

6 Higher Degrees of Mathias-absoluteness

COROLLARY 6.1 Σ_5^1 -Mathias-absoluteness implies $\Delta_4^1(\mathcal{R})$.

PROOF: Let $\Phi(x)$ be a Δ_4^1 -set:

$$\Phi(x) \leftrightarrow \varphi(x) \leftrightarrow \neg\psi(x)$$

where $\varphi(x)$ and $\psi(x)$ are Σ_4^1 -sets. By Σ_5^1 - \mathbf{M} -absoluteness, $\Phi(x)$ is still a Δ_4^1 -set in $V^{\mathbf{M}}$. Let p be an \mathbf{M} -condition with empty stem such that

$$V \models "p \Vdash_{\mathbf{M}} \varphi(\mathbf{m})",$$

(if $V \models "p \Vdash_{\mathbf{M}} \neg\varphi(\mathbf{m})"$ then $V \models "p \Vdash_{\mathbf{M}} \psi(\mathbf{m})"$), then there is an \mathbf{M} -name \tilde{y} and (by Corollary 3.11) a $p' \subseteq p$ with empty stem, such that

$$V \models "p' \Vdash_{\mathbf{M}} \varphi_0(\mathbf{m}, \tilde{y})"$$

(where $\varphi(x) \equiv \exists y \varphi(x, y)$ and φ_0 is a Π_3^1 -formula). Let $m \subseteq p'$ be Mathias over V , then

$$V[m] \models \varphi_0(m, \tilde{y}[m]).$$

Now in the proof of Lemma 5.4 in fact we showed, that if m is Mathias over V , $m' \in [m]^\omega \cap V[m]$, $\forall r \in [\omega]^\omega \cap V(\omega_1^{L[r]} < \omega_1^V)$ and $\Phi(x)$ is a Σ_3^1 -set (or a Π_3^1 -set) with parameters in V , then

$$V[m'] \models \Phi(m') \Leftrightarrow V[m] \models \Phi(m').$$

Because of $m' \in [m]^\omega \cap V[m]$, m' is also Mathias over V and the sentence $\forall x \in [m]^\omega (\varphi_0(x, \tilde{y}[m]))$ holds in $V[m]$. Therefore $\exists z \forall x \in [z]^\omega \neg \psi(x)$, which is a Σ_5^1 -sentence with parameters in V , is true in $V[m]$. Hence, $V \models \text{“}\Phi(x) \text{ is Ramsey”}$ and because $\Phi(x)$ was arbitrary we get $V \models \Delta_4^1(\mathcal{R})$. \dashv

To prove the last results, we need two slightly technical lemmas.

LEMMA 6.2 *If $\forall r \in [\omega]^\omega \cap V(\omega_1^{L[r]} < \omega_1^V)$ and $\Phi(\tilde{z})$ is a Σ_3^1 -formula (where \tilde{z} is an \mathbf{M} -name in V for a real), then: for all \mathbf{M} -conditions q in V there is a real a and an \mathbf{M} -condition l in V such that (q is an \mathbf{M} -condition in $L[a]$ and $l \leq q$) and for all reals m : if m is Mathias over V and $l \in m$, then (m is Mathias over $L[a]$ and $\tilde{z}[m] \in L[a]$ and ($L[a][m] \models \Phi(\tilde{z}[m])$ if and only if $V[m] \models \Phi(\tilde{z}[m])$)).*

PROOF: To simplify the notation we assume that the parameters of Φ are in L .

Assume $V \models \text{“}q_0 \Vdash_{\mathbf{M}} \Psi(\tilde{z}, \tilde{x})\text{”}$ where $q_0 \leq q$ and $\Phi(z) \equiv \exists x \Psi(z, x)$. Let $r_0 \leq q_0$ and \tilde{z}, \tilde{x} such that $V \models \text{“}r_0 \Vdash_{\mathbf{M}} \tilde{z} = \tilde{z} \wedge \tilde{x} = \tilde{x}\text{”}$. Let a be a real which encode the reals $r_0, \tilde{z}, \tilde{x}$ and q . In $L[a]$ there must be an \mathbf{M} -condition $q_1 \leq r_0$ such that $L[a] \models \text{“}q_1 \Vdash_{\mathbf{M}} \Psi(\tilde{z}, \tilde{x})\text{”}$ (because of the absoluteness of Π_2^1 -formulas). Let $l \in V$ be Mathias over $L[a]$ such that $q_1 \in l$ and further let m be Mathias over V such that $l \in m$, then $L[a][m] \models \Psi(\tilde{z}[m], \tilde{x}[m])$ and $V[a][m] \models \Psi(\tilde{z}[m], \tilde{x}[m])$.

If $V \models \text{“}q_0 \Vdash_{\mathbf{M}} \neg \Phi(\tilde{z})\text{”}$ for all $q_0 \leq q$ which decides $\Phi(\tilde{z})$, there is an \mathbf{M} -condition q_1 as in the former case, (because Π_3^1 -formulas are downwards absolute). The rest of the proof in this case is the same as above. \dashv

We say $L[a]$ computes well the Σ_3^1 formula $\Phi(\tilde{z})$ (the Π_3^1 formula $\neg \Phi(\tilde{z})$, respectively) with respect to q_1 .

LEMMA 6.3 *If V is Σ_4^1 - \mathbf{M} -absolute, then V is Σ_4^1 - \mathbf{M} -correct.*

PROOF: If not, then there is a Σ_4^1 -formula $\Phi(x)$ and an \mathbf{M} -condition $p \in V$ such that $V \models \text{“}p \Vdash_{\mathbf{M}} \tilde{x} \in [\mathbf{m}]^\omega \wedge \Phi(\tilde{x}) \wedge V[\tilde{x}] \not\models \Phi(\tilde{x})\text{”}$. Because $V \models \text{“}p \Vdash_{\mathbf{M}} \Phi(\tilde{x})\text{”}$ there is an \mathbf{M} -name \tilde{y} such that $V \models \text{“}p \Vdash_{\mathbf{M}} \Psi(\tilde{x}, \tilde{y})\text{”}$ where $\Phi(x) \equiv \exists y \Psi(x, y)$ and $\Psi(x, y)$ is a Π_3^1 -formula.

Let r, \tilde{x}, \tilde{y} be such that $r \leq p$ and $V \models \text{“}r \Vdash_{\mathbf{M}} \tilde{x} = \tilde{x} \wedge \tilde{y} = \tilde{y}\text{”}$. By Lemma 6.2 there is an $a \in V$ and an \mathbf{M} -condition $q \leq r$ such that $L[a]$ computes well $\Psi(\tilde{x}, \tilde{y})$ with respect to q . Let l and m as in the Lemma 6.2 and further let $x := \tilde{x}[m]$. Because m is Mathias over $L[a]$ and $x \in L[a][m]$ we can write the Mathias forcing as a two step iteration $Q_1 * \tilde{Q}_2$ and choose (as in the proof of Lemma 5.4) $g_1, g_2 \in V[x]$ such that g_1 is Q_1 -generic over $L[a]$, g_2 is $\tilde{Q}_2[g_1]$ -generic over $L[a][g_1]$, $g_1 * g_2$ is \mathbf{M} -generic over $L[a]$ with respect to q and $L[a][g_1] = L[a][x]$. With the same arguments as in the proof of Lemma 5.4 we have $L[a][x][g_2] \models \Psi(x, \tilde{y}[g_1 * g_2])$. Now because $L[a]$ computes well the Π_3^1 -formula Ψ and $g_2 \in V[x]$, we finally have $V[x] \models \Phi(x)$. \dashv

THEOREM 6.4 *Σ_6^1 -Mathias-absoluteness implies $\Sigma_4^1(\mathcal{R})$.*

PROOF: Let $\Phi(x)$ be a Σ_4^1 -formula with parameters in V and further let $p \in V$ be an \mathbf{M} -condition which decides $\Phi(\mathbf{m})$.

If $V \models "p \Vdash_{\mathbf{M}} \Phi(\mathbf{m})"$ then by Lemma 6.3 $V \models "p \Vdash_{\mathbf{M}} \exists x \forall y \in [x]^\omega \Phi(y)"$.

If $V \models "p \Vdash_{\mathbf{M}} \neg \Phi(\mathbf{m})"$ then by Lemma 6.3 $V \models "p \Vdash_{\mathbf{M}} \exists x \forall y \in [x]^\omega \neg \Phi(y)"$.

In both cases (by Σ_6^1 - \mathbf{M} -absoluteness) we get that $\Phi(x)$ is Ramsey in V and because $\Phi(x)$ was arbitrary we have $V \models \Sigma_4^1(\mathcal{R})$. \dashv

COROLLARY 6.5 Σ_6^1 -Mathias-absoluteness implies $\Delta_5^1(\mathcal{R})$.

PROOF: Let $\Phi(x)$ be a Δ_5^1 -set:

$$\Phi(x) \leftrightarrow \varphi(x) \leftrightarrow \neg \psi(x)$$

where $\varphi(x)$ and $\psi(x)$ are Σ_5^1 -sets. By Σ_6^1 - \mathbf{M} -absoluteness, $\Phi(x)$ is still a Δ_5^1 -set in $V^{\mathbf{M}}$. Let p be an \mathbf{M} -condition with empty stem such that $V \models "p \Vdash_{\mathbf{M}} \varphi(\mathbf{m})"$, (if $V \models "p \Vdash_{\mathbf{M}} \neg \varphi(\mathbf{m})"$ then $V \models "p \Vdash_{\mathbf{M}} \psi(\mathbf{m})"$), then there is an \mathbf{M} -name \tilde{y} and (by Corollary 3.11) a $p' \subseteq p$ with empty stem, such that

$$V \models "p' \Vdash_{\mathbf{M}} \varphi_0(\mathbf{m}, \tilde{y})"$$

(where $\varphi(x) \equiv \exists y \varphi_0(x, y)$ and φ_0 is a Π_4^1 -formula). Let $m \subseteq p'$ be Mathias over V , then

$$V[m] \models \varphi_0(m, \tilde{y}[m]).$$

Because of Lemma 6.3 and because $m' \in [m]^\omega \cap V[m]$ is Mathias over V , the sentence $\forall m' \in [m]^\omega \varphi_0(m', \tilde{y}[m'])$ which is $V[m] \models \exists z \forall x \in [z]^\omega \varphi(x)$, holds in $V[m]$. Therefore $\exists z \forall x \in [z]^\omega \neg \psi(x)$ which is a Σ_6^1 -sentence with parameters in V is true in $V[m]$.

Hence, $V \models "\Phi(x)$ is Ramsey" and because $\Phi(x)$ was arbitrary we get

$$V \models \Delta_5^1(\mathcal{R}).$$

\dashv

References

- [BJ] T. BARTOSZYŃSKI AND H. JUDAH: "Set Theory: the structure of the real line," (in print).
- [Ba] J. E. BAUMGARTNER: Iterated forcing, in "Surveys in set theory," (A. R. D. Mathias, Ed.), pp. 1–59, Cambridge University Press, London and New York, 1983.
- [Br] J. BRENDLE: Combinatorial properties of classical forcing notions, (to appear).
- [Bn] J. BROWN: The Ramsey sets and related σ -algebras and ideals. *Fund. Math.* **136**(1990), 179–185.
- [Co] P. CORAZZA: Ramsey sets, the Ramsey ideal, and other classes over \mathbf{R} . *J. Symbolic Logic* **57**(1992), 1441–1468.
- [El] E. ELLENTUCK: A new Proof that analytic Sets are Ramsey. *J. Symbolic Logic* **39**(1974), 163–165.
- [ER] P. ERDÖS AND R. RADO: Combinatorial theorems on classifications of subsets of a given set. *Proc. of the London Math. Soc.* **2**(1952), 417–439.
- [Go] M. GOLDSTERN: Tools for your forcing construction, in "Set Theory of the Reals," (H. Judah, Ed.), pp. 305–360, Israel Mathematical Conference Proceedings, Bar-Ilan University, 1993.

- [Je 1] TH. JECH: “Multiple Forcing.” Cambridge University Press, Cambridge 1987.
- [Je 2] TH. JECH: “Set Theory.” Academic Press, London 1978.
- [Ju 1] H. JUDAH: Absoluteness for projective sets, *in* “Logic Colloquium ‘90,” (J. Oikkonen et al., Ed.), pp. 145–154, Springer-Verlag, Berlin, 1993.
- [Ju 2] H. JUDAH: Δ_3^1 -sets of reals, *in* “Set Theory of the Reals,” (H. Judah, Ed.), pp. 361–384, Israel Mathematical Conference Proceedings, Bar-Ilan University, 1993.
- [Ju 3] H. JUDAH: Exact equiconsistency results for Δ_3^1 -sets of reals. *Arch. Math. Logic* **32**(1992), 101–112.
- [JS 1] H. JUDAH AND S. SHELAH: Δ_2^1 -sets of reals. *Ann. of Pure and Applied Logic* **42**(1989), 207–223.
- [JS 2] H. JUDAH AND S. SHELAH: Δ_3^1 -sets of reals. *J. Symbolic Logic* **58**(1993), 72–80.
- [Ku] K. KUNEN: “Set Theory, an Introduction to Independence Proofs.” North Holland, Amsterdam 1983.
- [Ma] A. R. D. MATHIAS: Happy families. *Ann. of Math. Logic* **12**(1977), 59–111.
- [Si] J. SILVER: Every analytic Set is Ramsey. *J. Symbolic Logic* **35**(1970), 60–64.

Lorenz Halbeisen
 Departement Mathematik
 ETH Zentrum
 8092 Zürich
 Switzerland
 E-mail: halbeis@math.ethz.ch

Haim Judah
 Department of Mathematics
 Bar-Ilan University
 52900 Ramat Gan
 Israel
 E-mail: judah@bimacs.cs.biu.ac.il