

# CYCLE INTEGRALS OF MODULAR FUNCTIONS, MARKOV GEODESICS AND A CONJECTURE OF KANEKO

P. BENGOCHEA AND Ö. IMAMOGLU

ABSTRACT. In this paper we study the values of modular functions at the Markov quadratics which are defined in terms of their cycle integrals along the associated closed geodesics. These numbers are shown to satisfy two properties that were conjectured by Kaneko. More precisely we show that the values of a modular function  $f$ , along any branch  $B$  of the Markov tree, converge to the value of  $f$  at the Markov number which is the predecessor of the tip of  $B$ . We also prove an interlacing property for these values.

## 1. INTRODUCTION

A well known theorem of Dirichlet asserts that for any irrational number  $x$ , there are infinitely many rational numbers  $p/q$  satisfying  $|x - \frac{p}{q}| < \frac{1}{q^2}$ . For irrational numbers that are algebraic, thanks to a theorem of Roth [12], the exponent 2 is optimal. The constant factor, on the other hand, can be improved and a classical theorem of Hurwitz asserts that for every irrational number  $x$  there exist infinitely many rational numbers  $p/q$  satisfying

$$\left| x - \frac{p}{q} \right| < \frac{1}{\sqrt{5}q^2}.$$

The constant  $1/\sqrt{5}$  is best possible but if we exclude as  $x$  the numbers that are  $\mathrm{PGL}(2, \mathbb{Z})$ -equivalent to the golden ratio  $(1 + \sqrt{5})/2$ , the constant  $1/\sqrt{5}$  improves to  $1/\sqrt{8}$ . If we also exclude the numbers that are  $\mathrm{PGL}(2, \mathbb{Z})$ -equivalent to  $\sqrt{2}$ , then the constant improves to  $5/\sqrt{221}$ . By proceeding in this way, one obtains the Lagrange spectrum defined by

$$L := \{\nu(x)\}_{x \in \mathbb{R}} \subseteq \left[0, 1/\sqrt{5}\right] \quad \text{with} \quad \nu(x) = \liminf_{q \rightarrow \infty} q \|qx\|,$$

where  $\|x\|$  denotes the distance from a real number  $x$  to a closest integer. The quantity  $\nu(x)$  provides a measure of approximation of  $x$  by the rationals. For almost all  $x \in \mathbb{R}$  we have  $\nu(x) = 0$  and when  $\nu(x) > 0$  we call  $x$  badly approximable. Real quadratic irrationals are badly approximable, the worst ones being the golden ratio and its  $\mathrm{PGL}(2, \mathbb{Z})$ -equivalents, followed by  $\sqrt{2}$  and its  $\mathrm{PGL}(2, \mathbb{Z})$ -equivalents, etc. The Lagrange spectrum is not discrete (cf [5]) but the part of

---

Bengoechea's research is supported by SNF grant 173976.

the spectrum in the subinterval  $(1/3, 1/\sqrt{5}]$  corresponding to classes of worst irrational numbers is, with  $1/3$  as its only accumulation point.  $L \cap (1/3, 1/\sqrt{5}]$  is well understood thanks to the work of Markov (cf [10], [11]) which connects this question of Diophantine approximation to the Diophantine equation

$$(1) \quad x^2 + y^2 + z^2 = 3xyz.$$

The set of Markov triples comprising of the positive integer solutions  $(x, y, z)$  of (1) can be obtained from  $(1, 1, 1)$  by repeated applications of permutations and Vieta involutions  $(x, y, z) \rightarrow (x, y, 3xy - z)$ . The Markov numbers are the greatest coordinates of Markov triples. They form the Markov sequence


$$\{m_i\}_{i=1}^{\infty} = \{1, 2, 5, 13, 29, 34, 89, 169, 194, \dots\}.$$

The Markov number  $m_i$  is associated to a quadratic irrationality


$$\theta_i = \frac{-3m_i + 2k_i + \sqrt{9m_i^2 - 4}}{2m_i},$$

where  $k_i$  is an integer that satisfies  $a_i k_i \equiv b_i \pmod{m_i}$  and  $(a_i, b_i, m_i)$  is a solution to (1) with  $m_i$  maximal. Markov showed that  $\nu(\theta_i) = \sqrt{9 - 4/m_i^2}$ , and  $L \cap (1/3, 1/\sqrt{5}] = \{\nu(\theta_i)\}_{i \geq 1}$ . Moreover, any  $x \in \mathbb{R}$  for which  $\nu(x) \in L \cap (1/3, 1/\sqrt{5}]$  is  $\text{PGL}(2, \mathbb{Z})$ -equivalent to a Markov quadratic  $\theta_i$ .

Markov numbers come with a tree structure, inherit from Vieta involutions, that arranges them as below:



Here  $(a, b, c)$  is a solution to (1). The Markov quadratics inherit the same tree structure which can be given in terms of their continued fractions as



where  $k_n$  means that  $k$  is repeated  $n$  times.

Markov numbers arise in many different contexts: see for example, [1], [9] and references therein and [2], [3], [4] for some recent developments regarding the Markov surfaces.

The main goal of this paper is to study the values of modular functions along the tree associated to the Markov quadratics.

Let  $\Gamma = \text{PSL}(2, \mathbb{Z})$ . For a general quadratic irrationality  $w \in \mathbb{Q}(\sqrt{D})$  and a modular function  $f$  for  $\Gamma$ , the “value” of  $f$  at  $w$  is defined in terms of the integral of  $f$  along the geodesic cycle  $C_w \in \Gamma \backslash \mathcal{H}$  associated to  $w$ . More precisely

$$f(w) := \sqrt{D} \int_{C_w} f(z) ds$$

where  $ds$  is the hyperbolic arc length. We can normalize the number  $f(w)$  by the length of the geodesic  $C_w$  and define

$$f^{nor}(w) := \frac{f(w)}{2 \log \varepsilon_D}$$

where  $\varepsilon_D$  is the fundamental unit.

The values of modular functions at real quadratic irrationalities were introduced in [7] and independently in [8]. In [7] their averages over ideal classes were shown to be coefficients of mock modular forms whereas in [8] Kaneko studied their individual values  $f^{nor}(w)$  (in the case that the modular function is the Klein’s  $j$  invariant), and based on numerical calculations he made several interesting observations and conjectures.

In this paper we prove two of Kaneko’s conjectures which involve the values of modular functions at the Markov quadratics. Our first theorem shows that if  $B$  is any branch of the tree,  $w_n^B$  is the  $n$ -th Markov quadratic on  $B$  and  $w_0^B$  is the predecessor of the tip of  $B$  then the normalized values  $f^{nor}(w_n^B)$ , for any modular function  $f$ , converge to the value  $f^{nor}(w_0^B)$ . More precisely

**Theorem 1.1.** *Let  $f$  be a modular function defined on  $\mathcal{H}$  for  $\Gamma$ . For any branch  $B$  of the Markov tree we have*

$$\lim_{n \rightarrow \infty} f^{nor}(w_n^B) = f^{nor}(w_0^B).$$

Our second theorem proves an eventual monotonicity result which also partially proves the interlacing property of the values for the Markov quadratics that was conjectured by Kaneko;

**Theorem 1.2.** *Let  $f$  be a modular function on  $\mathcal{H}$ , let  $B$  be any branch of the Markov tree and  $r + 1$  be the number of partial quotients of  $w_0^B$ . Then there exists a constant  $N_f$  such that, for all  $n \geq N_f/r$ , the real and imaginary parts of  $f^{nor}(w_{n+1}^B)$  lie between the real and respectively imaginary parts of  $f^{nor}(w_0^B)$  and  $f^{nor}(w_n^B)$ .*

The rest of the paper is organized as follows. In the next section we give the preliminaries about cycle integrals and continued fractions. In section 3, we give the basic properties of the Markov quadratics and the Markov tree. In section 4 and 5 we study the values of modular functions on the Markov tree and prove Theorem 1.1 and Theorem 1.2 respectively.

## 2. PRELIMINARIES

**2.1. Cycle integrals.** Let  $w$  be a real quadratic irrationality and  $\tilde{w}$  be its conjugate.  $w$  and  $\tilde{w}$  are the roots of a quadratic equation

$$ax^2 + bx + c = 0 \quad (a, b, c \in \mathbb{Z}, \quad (a, b, c) = 1)$$

with discriminant  $D = b^2 - 4ac > 0$ . We choose  $a, b, c$  such that  $w = \frac{-b + \sqrt{D}}{2a}$ ,  $\tilde{w} = \frac{-b - \sqrt{D}}{2a}$ . The geodesic  $S_w$  in  $\mathcal{H}$  joining  $w$  and  $\tilde{w}$  is given by the equation

$$a|z|^2 + b \operatorname{Re}(z) + c = 0 \quad (z \in \mathcal{H}).$$

The stabilizer  $\Gamma_w$  of  $w$  in  $\Gamma$  preserves the quadratic form  $Q_w = [a, b, c]$ , and hence  $S_w$ . The group  $\Gamma_w$  is infinite cyclic; it corresponds to the group  $U_D^2$  of units of norm one of  $\mathbb{Q}(\sqrt{D})$  via the isomorphism:

$$(2) \quad \begin{array}{ccc} \Gamma_w & \longrightarrow & U_D^2 \\ \begin{pmatrix} a & b \\ c & d \end{pmatrix} & \mapsto & (a - cw)^2. \end{array}$$

We denote by  $A_w$  the generator of  $\Gamma_w$

$$A_w = \begin{pmatrix} \frac{1}{2}(t - bu) & -cu \\ au & \frac{1}{2}(t + bu) \end{pmatrix},$$

where  $(t, u)$  is the smallest positive solution to Pell's equation  $t^2 - Du^2 = 4$ , and we denote by  $\varepsilon$  the generator of the infinite cyclic part of  $U_D$  whose square corresponds to  $A_w$  by the isomorphism (2).

For any modular function  $f$ , since the group  $\Gamma_w$  preserves the expression  $f(z)Q_w(z, 1)^{-1}dz$ , one can define the cycle integral of  $f$  along  $C_w = S_w/\Gamma_w$ , also viewed as the “value” of  $f$  at  $w$ , by the complex number

$$f(w) := \int_{C_w} \frac{\sqrt{D}f(z)}{Q_w(z, 1)} dz.$$

The factor  $\sqrt{D}$  is introduced here for convenience. The integral defining  $f(w)$  is  $\Gamma$ -invariant and can in fact be taken along any path in  $\mathcal{H}$  from  $z_0$  to  $A_w^{-1}z_0$ , where  $z_0$  is any point in  $\mathcal{H}$ . Note that this gives an orientation on  $S_w$  from  $w$  to  $\tilde{w}$ , which is counterclockwise if  $a > 0$  and clockwise if  $a < 0$ . We normalize the number  $f(w)$  by the length of the geodesic  $C_w$  which is given by

$$\int_{C_w} \frac{\sqrt{D}}{Q_w(z, 1)} dz = 2 \log \varepsilon$$

and we define the normalized value as

$$f^{nor}(w) := \frac{f(w)}{2 \log \varepsilon}.$$

**2.2. The ‘+’ and ‘-’ continued fractions.** Let  $(b_0, b_1, b_2, \dots)$  denote the ‘-’ continued fraction

$$(b_0, b_1, b_2, \dots) = b_0 - \frac{1}{b_1 - \frac{1}{b_2 - \frac{1}{\ddots}}}$$

and  $[a_0, a_1, a_2, \dots]$  be the ‘+’ continued fraction

$$[a_0, a_1, a_2, \dots] = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\ddots}}}$$

Every real number  $w$  has a ‘-’ continued fraction expansion  $w = (b_0, b_1, b_2, \dots)$  with  $b_i \in \mathbb{Z}$  and  $b_i \geq 2$  for  $i \geq 1$  and a ‘+’ continued fraction expansion  $w = [a_0, a_1, a_2, \dots]$  with  $a_i \in \mathbb{Z}$  and  $a_i \geq 1$  for  $i \geq 1$ . The ‘-’ continued fraction expansion of  $w$  is obtained by setting  $w_0 = w$  and inductively  $b_i = [w_i]$ ,  $w_{i+1} = \frac{1}{b_i - w_i} = ST^{-b_i}(w_i)$ , where  $S(x) = -1/x$  and  $T(x) = x + 1$ . The ‘+’ continued fraction expansion is obtained by setting  $a_i = [w_i]$ ,  $w_{i+1} = \frac{1}{w_i - a_i} = \varepsilon T^{-a_i}(w_i)$ , where  $\varepsilon(x) = 1/x$ . Hence the ‘-’ continued fraction is given by transformations of  $\Gamma$  on the real line, whereas the ‘+’ continued fraction corresponds to transformations of  $GL(2, \mathbb{Z})$ . To go from the ‘+’ to the ‘-’ continued fraction expansions,

the general rule is

$$(3) \quad [a_0, a_1, a_2, \dots] \longrightarrow (a_0 + 1, \underbrace{2, \dots, 2}_{a_1-1}, a_2 + 2, \underbrace{2, \dots, 2}_{a_3-1}, a_4 + 2, \dots).$$

It is well known that a real number  $w$  is a quadratic irrationality if and only if its ‘-’ continued fraction expansion (or equivalently, its ‘+’ continued fraction) is eventually periodic:  $w = (b_0, b_1, \dots, b_k, \overline{b_{k+1}, \dots, b_{k+r}})$  where the line over  $b_{k+1}, \dots, b_{k+r}$  denotes the period. We say that  $w$  is *purely periodic* when all the partial quotients repeat. It will be useful for the rest of the paper to remember the following statements:

- (I) two quadratic irrationalities have the same ‘-’ period if and only if they are  $\Gamma$ -equivalent;
- (II)  $w$  has a purely periodic ‘-’ continued fraction expansion if and only if  $0 < \tilde{w} < 1 < w$ , where  $\tilde{w}$  is the conjugate of  $w$ ;
- (III) if  $w = (\overline{b_0, \dots, b_r})$ , then  $\frac{1}{\tilde{w}} = (\overline{b_r, \dots, b_0})$ .

The following lemma gives an upper bound for the distance between two real numbers in terms of the number of first partial quotients for which they coincide.

**Lemma 2.1.** *If the ‘-’ continued fraction expansions of  $u$  and  $v$  coincide in the first  $r + 1$  partial quotients and their ‘+’ continued fraction expansions have only 1’s and 2’s, then*

$$|u - v| \leq 10 \left( \frac{2}{1 + \sqrt{5}} \right)^{2r}.$$

*Proof.* Let  $u$  and  $v$  be as in the statement of the lemma. Then one can see, by applying the rule (3), that also the ‘+’ continued fraction expansions of  $u$  and  $v$  coincide in the first  $r + 1$  partial quotients. Hence, if we set  $a_0, \dots, a_r$  to be those partial quotients, the rational number  $\frac{p}{q} = [a_0, \dots, a_r]$  is a convergent of both  $u$  and  $v$ . Then it is well known that

$$\left| u - \frac{p}{q} \right| \leq \frac{1}{q^2}, \quad \left| v - \frac{p}{q} \right| \leq \frac{1}{q^2}$$

and

$$q \geq \frac{1}{\sqrt{5}} \left( \frac{1 + \sqrt{5}}{2} \right)^r.$$

Therefore,


$$|u - v| \leq \left| u - \frac{p}{q} \right| + \left| v - \frac{p}{q} \right| \leq 10 \left( \frac{2}{1 + \sqrt{5}} \right)^{2r}.$$

□

3. MARKOV TREE

3.1. **Markov’s quadratics.** Let  $\{m_i\}_{i=1}^\infty = \{1, 2, 5, 13, 29, 34, 89, 169, 194, \dots\}$  be the set of Markov numbers. As in the introduction, for each Markov number  $m_i$ , we let  $\theta_i = \frac{-3m_i + 2k_i + \sqrt{9m_i^2 - 4}}{2m_i}$  be the Markov quadratic where  $k_i$  is an integer that satisfies  $a_i k_i \equiv b_i \pmod{m_i}$  and  $(a_i, b_i, m_i)$  is a solution to (1) with  $m_i$  maximal. In Markov’s theory, only  $\text{PGL}(2, \mathbb{Z})$ -equivalence classes are relevant, which implies that the order of  $(a_i, b_i)$  does not matter. Since we need  $\Gamma$ -equivalence to distinguish non-real  $f(\theta_i)$  and its conjugate, here the order of  $(a_i, b_i)$  becomes relevant. We fix it so that  $\text{Im}(f(w)) > 0$ .

The Markov tree  $\mathcal{T}$  associated to the Markov quadratics given in the introduction is in terms of the ‘+’ continued fractions. Since the cycle integrals are  $\Gamma$  and not  $\text{PGL}(2, \mathbb{Z})$  invariant, we will rather work with the ‘-’ continued fraction. By following the rule (3), the Markov tree  $\mathcal{T}$  becomes in the ‘-’ continued fraction:



Note that each branch in the tree  $\mathcal{T}$  comes with a left or right orientation. We call a branch a left (right) branch if starting from its tip on the top and going downwards the branch leans towards left (right). No zig-zag paths are allowed so each branch has a unique orientation. Except for the two singular cases of  $(2, \overline{3})$  and  $(3, \overline{2, 4})$ , each Markov number lies both on a right and a left branch but it is the tip of only a left or a right branch, except for  $(3, \overline{2, 3, 4})$  which is the tip of both the most left and the most right branches.

In the case of ‘+’ continued fractions we consider a *conjunction operation* of two periods as

$$(4) \quad [\overline{s_0, \dots, s_n}] \odot [\overline{t_0, \dots, t_m}] = [\overline{s_0, \dots, s_n, t_0, \dots, t_m}].$$

All Markov quadratics can be constructed by using this operation, starting with  $[\overline{1_2}]$  and  $[\overline{2_2}]$ . Indeed, each Markov quadratic is the result of the conjunction operation of its right and left closest predecessors on the tree.

For the ‘-’ continued fraction, the rule is also the conjunction of periods except for the most left branch, where the  $n$ -th Markov quadratic is  $(3, \overline{2, 3_n, 4})$ . Indeed,

let  $x = [\overline{s_0, \dots, s_n}] = (b_0, \overline{b_1, \dots, b_k})$  and  $y = [\overline{t_0, \dots, t_m}] = (c_0, \overline{c_1, \dots, c_\ell})$ . For any branch different from the right most branch, by applying (3) together with the observation that  $s_n = t_m = 1$  are in odd positions, so they do not contribute in the ‘ $\circ$ ’ expansion, we obtain

$$x \circ y = (b_0, \overline{b_1, \dots, b_{k-1}, t_0 + 2, c_1, \dots, c_{\ell-1}, s_0 + 2}).$$

But  $t_0$  is equal to 1 on the left most branch and 2 on any other branch, and  $s_0 = 2$ . For the right most branch, (3) also gives

$$x \circ y = (b_0, \overline{b_1, \dots, b_{k-1}, 4, c_1, \dots, c_{\ell-1}, s_0 + 2})$$

and  $s_0 = 2$ .

Throughout the paper, we denote by  $w_n^B$  ( $n \geq 1$ ) the  $n$ -th Markov quadratic on a branch  $B$  of the tree and  $w_0^B$  the left (right) predecessor of the tip  $w_1^B$  of  $B$  if  $B$  is a left (right) branch. For example, if  $B = L$  is the left most branch, then  $w_0^L = (2, \overline{3})$ ,  $w_1^L = (3, \overline{2, 3, 4})$ ,  $w_2^L = (3, \overline{2, 3_2, 4})$ ,  $w_3^L = (3, \overline{2, 3_3, 4})$ , etc. If  $B = R$  is the right most branch, then  $w_0^R = (3, \overline{2, 4})$ ,  $w_1^R = (3, \overline{2, 3, 4})$ ,  $w_2^R = (3, \overline{2, 4, 2, 3, 4})$ ,  $w_3^R = (3, \overline{(2, 4)_2, 2, 3, 4})$ , etc.

The  $n$ -th Markov quadratic on a left branch  $B \neq L$  can be written as:

$$(5) \quad w_n^B = (3, \overline{a_1, \dots, a_s, (b_1, \dots, b_r)_n}),$$

where  $w_0^B = (3, \overline{b_1, \dots, b_r})$  and  $a_1, \dots, a_s$  depend only on  $B$ . On a right branch  $B$ , we have

$$w_n^B = (3, \overline{(b_1, \dots, b_r)_{n-1}, a_1, \dots, a_s}),$$

and on the left most branch  $L$  we have

$$w_n^L = (3, \overline{2, 3_n, 4}).$$

**Remark 3.1.** *The left most branch in the Markov tree is also called the Fibonacci branch since the associated Markov numbers on this branch are the odd indexed Fibonacci numbers. Similarly the right most branch is associated with the odd indexed Pell numbers which are defined by the recurrence  $P_0 = 0, P_1 = 1$  and  $P_{n+1} = 2P_n + P_{n-1}$ . (cf. [1] p. 49)*

**3.2. The cycle of quadratics of a Markov number.** For any quadratic irrationality  $w$ , it is known that the hyperbolic element  $A_w$  is conjugate to a word in  $T$  and  $V$ , where

$$T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad V = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}.$$

If  $w = w_n^B$  is a quadratic on  $\mathcal{T}$  ( $n \geq 0$ ), then the associated hyperbolic element  $A_{w_n^B}$  can be written as a word in  $T$  and  $V$ . More specifically,  $A_{w_n^B} = A_0^{-1} \dots A_{\ell_n}^{-1}$ , where  $A_0 = I$  and  $A_i \in \{T^{-1}, V^{-1}\}$  for  $1 \leq i \leq \ell_n$  are given by the finite algorithm:

$$w_{n,0}^B = w_n^B, \quad w_{n,i+1}^B = A_{i+1}(w_{n,i}^B) \quad (i \geq 0),$$



where

$$A_{i+1} = \begin{cases} T^{-1} & \text{if } \lfloor w_{n,i}^B \rfloor \geq 1, \\ V^{-1} & \text{otherwise.} \end{cases}$$

Hence

$$(6) \quad w_{n,i}^B = A_i \dots A_0 w_n^B, \quad i = 0, \dots, \ell_n,$$

and  $\ell_n$  is the length of the word  $A_{w_n^B}$ , or equivalently, the length of the cycle of quadratics  $\{w_{n,i}^B\}_i$  of  $w_n^B$ .

**Example 3.2.** For example, the cycle of  $w_1^L = (3, \overline{2, 3, 4})$  on the most left branch is:

$$\begin{aligned} w_{1,0}^L &= (3, \overline{2, 3, 4}) \\ w_{1,1}^L &= T^{-1}(w_{1,0}^L) = (2, \overline{2, 3, 4}) \\ w_{1,2}^L &= T^{-1}(w_{1,1}^L) = (1, \overline{2, 3, 4}) \\ w_{1,3}^L &= V^{-1}(w_{1,2}^L) = (1, \overline{3, 4, 2}) \\ w_{1,4}^L &= V^{-1}(w_{1,3}^L) = (2, \overline{4, 2, 3}) \\ w_{1,5}^L &= T^{-1}(w_{1,4}^L) = (1, \overline{4, 2, 3}) \\ w_{1,6}^L &= V^{-1}(w_{1,5}^L) = (3, \overline{2, 3, 4}) = w_{1,0}^L. \end{aligned}$$

The length is  $\ell_1 = 6$  and  $A_{w_1^L} = ITTVVTV$ .

#### 4. CONVERGENCE PROPERTY

In this section we study the values of a modular function on the Markov tree. Let  $B$  be any branch of the tree and  $w_n^B$  be the  $n$ -th Markov quadratic on  $B$ . Let  $\rho = e^{\pi i/3}$  and  $z_i = A_0^{-1} \dots A_i^{-1} \rho^2$ . Then

$$\begin{aligned} f(w_n^B) &= -\sqrt{D} \sum_{i=0}^{\ell_n-1} \int_{z_i}^{z_{i+1}} \frac{f(z)}{Q_{w_n^B}(z, 1)} dz \\ &= -\sqrt{D} \sum_{i=0}^{\ell_n-1} \int_{\rho^2}^{A_{i+1}^{-1} \rho^2} \frac{f(z)}{(Q_{w_n^B} | A_0^{-1} \dots A_i^{-1})(z, 1)} dz, \\ &= -\sum_{i=0}^{\ell_n-1} \int_{\rho^2}^{A_{i+1}^{-1} \rho^2} f(z) \left( \frac{1}{z - w_{n,i}^B} - \frac{1}{z - \tilde{w}_{n,i}^B} \right) dz. \end{aligned}$$

Since  $V(\rho^2) = T(\rho^2) = \rho$ , we obtain:

**Lemma 4.1.** *For  $n \geq 0$  we have*

$$(7) \quad f(w_n^B) = \int_{\rho}^{\rho^2} \sum_{i=0}^{\ell_n-1} f(z) \left( \frac{1}{z - w_{n,i}^B} - \frac{1}{z - \tilde{w}_{n,i}^B} \right) dz.$$

Throughout the paper, we denote by  $\mathcal{C}$  the arc of circle joining  $\rho^2$  and  $\rho$ . We denote by  $\varepsilon_n^B$  the image of  $A_{w_n}^B$  under the isomorphism (2), so the length of  $C_{w_n^B}$  equals  $2 \log \varepsilon_n^B$ .

Our first goal is to show that the normalized values  $f^{nor}(w_n^B)$  for any modular function  $f$  along any branch  $B$  converge to the value  $f^{nor}(w_0^B)$ . We call this property ‘convergence property’ and prove it in this section. For simplicity of the notation, from this point on we restrict to a left but not the most left branch. However, the argument in the proof of Theorem 4.3 applies in the same way if  $B$  is a right branch or  $B = L$ . Only the bound  $\delta_1(r, N)$  will be slightly modified. Hence Corollary 4.4 remains true for any branch.

We consider  $w_n^B$  written as in (5). Then

$$\ell_n = n\ell_0 + \sum_{i=1}^s (a_i - 1),$$

where

$$\ell_0 = \sum_{i=1}^r (b_i - 1)$$

is the length of the cycle of  $w_0^B$ . The number of partial quotients in the period of  $w_1^B$  is  $s + r$  and the conjunction operation ensures that this is  $\leq 2r$ . Hence  $s \leq r$  and since  $a_i \leq 4$ , we have

$$(8) \quad \ell_n \leq 3r(n + 1).$$

It is convenient to set

$$a = \sum_{i=1}^s (a_i - 1)$$

and

$$\mathbf{p} = (b_1, \dots, b_r), \quad \mathbf{p}_k = (b_1, \dots, b_r)_k, \quad \mathbf{q}_k = (b_r, \dots, b_1)_k,$$

where the subindex  $k$  means that the continued fraction is repeated  $k$  times. With these notations, the cycle of  $w_n^B$  is of the form:

$$\begin{aligned}
 w_{n,0}^B &= (3, \overline{a_1, \dots, a_s, \mathbf{p}_n}), \\
 w_{n,1}^B &= (2, \overline{a_1, \dots, a_s, \mathbf{p}_n}), \\
 w_{n,2}^B &= (1, \overline{a_1, \dots, a_s, \mathbf{p}_n}), \\
 w_{n,3}^B &= (a_1 - 1, \overline{a_2, \dots, a_s, \mathbf{p}_n}, a_1), \\
 &\quad \vdots \\
 w_{n,a}^B &= (3, \overline{\mathbf{p}_n, a_1, \dots, a_s}), \\
 &\quad \vdots \\
 w_{n,a+\ell_0}^B &= (3, \overline{\mathbf{p}_{n-1}, a_1, \dots, a_s, \mathbf{p}}), \\
 &\quad \vdots \\
 w_{n,a+n\ell_0}^B &= (3, \overline{a_1, \dots, a_s, \mathbf{p}_n}) = w_{n,0}^B.
 \end{aligned}$$

**Remark 4.2.** *One can easily write the continued fraction expansion for the Galois conjugate  $-\tilde{w}_{n,i}^B$  of  $-w_{n,i}^B$  in terms of that of  $w_{n,i}^B$ . Indeed, let  $(d_0, \overline{d_1, \dots, d_m})$  be the continued fraction expansion of  $w_{n,i}^B$ . The quadratic  $ST^{-d_0}(w_{n,i}^B)$  is purely periodic with continued fraction  $(\overline{d_1, \dots, d_m})$  so, by the property (III), its Galois conjugate is  $1/(\overline{d_m, \dots, d_1})$ . Therefore,*

$$\tilde{w}_{n,i}^B = T^{d_0} S(1/(\overline{d_m, \dots, d_1})) = -(d_m - d_0, \overline{d_{m-1}, \dots, d_1, d_m}).$$

**Theorem 4.3.** *Let  $f$  be a modular function,  $B$  be any left branch  $\neq L$  of the Markov tree  $\mathcal{T}$  and  $N \geq 1$ . There exists a complex number  $K = K_{f,B,N}$  such that for all  $n \geq N$ ,*

$$(9) \quad |f(w_n^B) - nf(w_0^B) - K| \leq \delta_1(r, N) \max_{z \in \mathcal{C}} |f(z)|,$$

where

$$(10) \quad \delta_1(r, N) = \frac{80\pi}{3} (2 + r(N + 1)) \left( \frac{2}{1 + \sqrt{5}} \right)^{2(rN-1)}$$

and  $r + 1$  is the number of partial quotients of  $w_0^B$ .

*Proof.* By Lemma 4.1 we have:

$$(11) \quad f(w_n^B) - nf(w_0^B) = \int_{\rho}^{\rho^2} f(z) (S_1(n, N, z) + S_2(n, N, z) + S_3(n, N, z)) dz,$$

where

$$\begin{aligned}
S_1(n, N, z) &= \sum_{i=0}^{a-1} \frac{1}{z - w_{n,i}^B} + \sum_{i=a+(n-N)\ell_0}^{\ell_n-1} \frac{1}{z - w_{n,i}^B} \\
&\quad - \sum_{i=0}^{a+N\ell_0-1} \frac{1}{z - \tilde{w}_{n,i}^B} - N \sum_{i=0}^{\ell_0-1} \left( \frac{1}{z - w_{0,i}^B} - \frac{1}{z - \tilde{w}_{0,i}^B} \right), \\
S_2(n, N, z) &= \sum_{i=a}^{a+(n-N)\ell_0-1} \frac{1}{z - w_{n,i}^B} - (n - N) \sum_{i=0}^{\ell_0-1} \frac{1}{z - w_{0,i}^B}, \\
S_3(n, N, z) &= - \sum_{i=a+N\ell_0}^{a+n\ell_0-1} \frac{1}{z - \tilde{w}_{n,i}^B} + (n - N) \sum_{i=0}^{\ell_0-1} \frac{1}{z - \tilde{w}_{0,i}^B}.
\end{aligned}$$

Moreover, we can also write

$$(12) \quad S_1(n, N, z) = S_1(N, N, z) + (S_1(n, N, z) - S_1(N, N, z)).$$

Define

$$K := \int_{\rho}^{\rho^2} f(z) S_1(N, N, z) dz$$

and

$$c(n, z) := |S_1(n, N, z) - S_1(N, N, z)| + |S_2(n, N, z)| + |S_3(n, N, z)|.$$

Then

$$(13) \quad |f(w_n^B) - nf(w_0^B) - K| \leq \int_{\rho}^{\rho^2} c(n, z) |f(z)| |dz|.$$

Let  $\lambda = \left(\frac{2}{1+\sqrt{5}}\right)^2$ . If we can show that

$$(14) \quad c(n, z) \leq 80(2 + r(N + 1))\lambda^{rN-1}$$

for  $z \in \mathcal{C}$ , then the theorem is proved. Next we show (14).

**Bound for  $|S_2(n, N, z)|$ .** We have that

$$\begin{aligned}
|S_2(n, N, z)| &\leq \sum_{k=0}^{n-N-2} \sum_{i=1}^{\ell_0} \frac{|w_{n,2+a+k\ell_0+i}^B - w_{0,2+i}^B|}{|z - w_{n,2+a+k\ell_0+i}^B| |z - w_{0,2+i}^B|} \\
&\quad + \sum_{i=0}^2 \frac{|w_{n,a+i}^B - w_{0,i}^B|}{|z - w_{n,a+i}^B| |z - w_{0,i}^B|} + \sum_{i=1}^{\ell_0-3} \frac{|w_{n,2+a+(n-N-1)\ell_0+i}^B - w_{0,2+i}^B|}{|z - w_{n,2+a+(n-N-1)\ell_0+i}^B| |z - w_{0,2+i}^B|}.
\end{aligned}$$

The denominators are bounded below by  $\frac{3}{4}$  when  $z \in \mathcal{C}$  and the numerators can be bounded by using Lemma 2.1. For  $i = 0, 1, 2$ ,

$$w_{n,a+i}^B = (3 - i, \overline{\mathbf{p}_n}, a_1, \dots, a_s)$$

and

$$w_{0,i}^B = (3 - i, \overline{\mathbf{p}})$$

coincide at least in the first  $rn + 1$  partial quotients. For each  $0 \leq k \leq n - N - 2$ , we have:

for  $1 \leq i \leq b_1 - 1$ ,

$$(15) \quad w_{n,2+a+k\ell_0+i}^B = (b_1 - i, \overline{b_2, \dots, b_r, \mathbf{p}_{n-1-k}, a_1, \dots, a_s, \mathbf{p}_k, b_1});$$

for the next  $b_2 - 1$  values of  $i$  ( $b_1 \leq i \leq b_1 + b_2 - 2$ ),

$$(16) \quad w_{n,2+a+k\ell_0+i}^B = (b_2 - j, \overline{b_3, \dots, b_r, \mathbf{p}_{n-1-k}, a_1, \dots, a_s, \mathbf{p}_k, b_1, b_2})$$

with  $1 \leq j \leq b_2 - 1$ ; this process goes on until the last  $b_r - 1$  values of  $i$ , where

$$w_{n,2+a+k\ell_0+i}^B = (b_r - j, \overline{\mathbf{p}_{n-1-k}, a_1, \dots, a_s, \mathbf{p}_{k+1}})$$

with  $1 \leq j \leq b_r - 1$ . For  $k = n - N - 1$ , we have the same pattern as before except for the last block of values of  $i$ , where we only have  $b_r - 3$  of them.

Now, for each  $0 \leq k \leq n - N - 1$ , for  $1 \leq i \leq b_1 - 1$ , (15) and

$$w_{0,2+i}^B = (b_1 - i, \overline{b_2, \dots, b_r, b_1})$$

coincide in the first  $rn - rk$  partial quotients. The next  $b_2 - 1$  values of  $i$  (16) and

$$w_{0,2+i}^B = (b_2 - j, \overline{b_3, \dots, b_r, b_1, b_2}) \quad (1 \leq j \leq b_2 - 1)$$

coincide in the first  $rn - rk - 1$  partial quotients, similarly for the next  $b_3 - 1$  values of  $i$ ,  $w_{n,2+a+k\ell_0+i}^B$  and  $w_{0,2+i}^B$  coincide in the first  $rn - rk - 2$  partial quotients, etc. Therefore we have, for  $z \in \mathcal{C}$ ,

$$\begin{aligned} |S_2(n, N, z)| &\leq \frac{40}{3} \left( 3\lambda^{rn} + \sum_{i=1}^r (b_i - 1) \sum_{k=0}^{n-N-1} \lambda^{r(n-k)-i} \right) \\ &\leq \frac{40}{3} \left( 3\lambda^{rn} + 3 \left( \sum_{i=1}^r \lambda^{-i} \right) \left( \sum_{k=N+1}^n \lambda^{rk} \right) \right) \\ &\leq \frac{40}{3} \left( 3\lambda^{rn} + 3 \left( \sum_{i=1}^r \lambda^{r-i} \right) \left( \sum_{k=N}^{n-1} \lambda^{rk} \right) \right) \\ &\leq \frac{40}{3} \left( 3\lambda^{rn} + 3 \left( \sum_{i=0}^{r-1} \lambda^i \right) \left( \sum_{k=N}^{n-1} \lambda^{rk} \right) \right) \\ &\leq 40\lambda^{rN} \left( 1 + \frac{1}{(1-\lambda)(1-\lambda^r)} \right) \\ &\leq 120\lambda^{rN}. \end{aligned}$$

**Bound for  $|S_3(n, N, z)|$ .** In a similar way we bound  $|S_3(n, N, z)|$ . We have that

$$\begin{aligned} |S_3(n, N, z)| &\leq \sum_{k=N}^{n-2} \sum_{i=1}^{\ell_0} \frac{|\tilde{w}_{n,2+a+k\ell_0+i}^B - \tilde{w}_{0,2+i}^B|}{|z - \tilde{w}_{n,2+a+k\ell_0+i}^B| |z - \tilde{w}_{0,2+i}^B|} \\ &+ \sum_{i=0}^2 \frac{|\tilde{w}_{n,a+N\ell_0+i}^B - \tilde{w}_{0,i}^B|}{|z - \tilde{w}_{n,a+N\ell_0+i}^B| |z - \tilde{w}_{0,i}^B|} + \sum_{i=1}^{\ell_0-3} \frac{|\tilde{w}_{n,2+a+(n-1)\ell_0+i}^B - \tilde{w}_{0,2+i}^B|}{|z - \tilde{w}_{n,2+a+(n-1)\ell_0+i}^B| |z - \tilde{w}_{0,2+i}^B|}. \end{aligned}$$

For  $i = 0, 1, 2$ , using Remark 4.2, we have that

$$-\tilde{w}_{n,a+N\ell_0+i}^B = (1 + i, \overline{b_{r-1}, \dots, b_1, \mathbf{q}_{N-1}, a_s, \dots, a_1, q_{n-N}, b_r})$$

and

$$-\tilde{w}_{0,i}^B = (1 + i, \overline{b_{r-1}, \dots, b_1, b_r})$$

coincide in the first  $rN$  partial quotients. For each  $N \leq k \leq n-2$ , we have: for  $1 \leq i \leq b_1 - 1$ ,

$$(17) \quad -\tilde{w}_{n,2+a+k\ell_0+i}^B = (i, \overline{\mathbf{q}_k, a_s, \dots, a_1, \mathbf{q}_{n-k}});$$

for the next  $b_2 - 1$  values of  $i$  ( $b_1 \leq i \leq b_1 + b_2 - 2$ ),

$$(18) \quad -\tilde{w}_{n,2+a+k\ell_0+i}^B = (j, \overline{b_1, \mathbf{q}_k, a_s, \dots, a_1, \mathbf{q}_{n-1-k}, b_r, \dots, b_3, b_2})$$

with  $1 \leq j \leq b_2 - 1$ ; this process goes on until the last  $b_r - 1$  values of  $i$ , where

$$-\tilde{w}_{n,2+a+k\ell_0+i}^B = (j, \overline{b_{r-1}, \dots, b_1, \mathbf{q}_k, a_s, \dots, a_1, \mathbf{q}_{n-1-k}, b_r})$$

with  $1 \leq j \leq b_r - 1$ .

For  $k = n-1$ , we have the same pattern as before except for the last block of values of  $i$ , where we only have  $b_r - 3$  of them. Now, for each  $N \leq k \leq n-1$ , for the first  $b_1 - 1$  values of  $i$  (17) coincide with

$$-\tilde{w}_{0,2+i}^B = (i, \overline{\mathbf{q}})$$

in the first  $rk + 1$  partial quotients. For the next  $b_2 - 1$  values of  $i$  (18) coincide with

$$-\tilde{w}_{0,2+i}^B = (j, \overline{b_1, b_r, \dots, b_2}) \quad (1 \leq j \leq b_2 - 1)$$

in the first  $rk + 2$  partial quotients, for the next  $b_3 - 1$   $i$ -values  $-\tilde{w}_{n,2+a+k\ell_0+i}^B$  and  $-\tilde{w}_{0,2+i}^B$  coincide in the first  $rk + 3$  partial quotients, etc. Therefore we have, for

$z \in \mathcal{C}$ ,

$$\begin{aligned} |S_3(n, N, z)| &\leq \frac{40}{3} \left( 3\lambda^{rN-1} + \sum_{i=1}^r (b_i - 1) \sum_{k=N}^{n-1} \lambda^{rk+i-1} \right) \\ &\leq \frac{40}{3} \left( 3\lambda^{rN-1} + 3 \left( \sum_{i=1}^r \lambda^{i-1} \right) \left( \sum_{k=N}^{n-1} \lambda^{rk} \right) \right) \\ &\leq 40\lambda^{rN-1} \left( 1 + \frac{\lambda}{(1-\lambda)(1-\lambda^r)} \right) \\ &\leq 80\lambda^{rN-1}. \end{aligned}$$

**Bound for  $|S_1(n, N, z) - S_1(N, N, z)|$ .** We have

$$\begin{aligned} |S_1(n, N, z) - S_1(N, N, z)| &\leq \sum_{i=0}^{a-1} \frac{|w_{n,i}^B - w_{N,i}^B|}{|z - w_{n,i}^B||z - w_{N,i}^B|} \\ &+ \sum_{i=a}^{\ell_N-1} \frac{|w_{n,i+(n-N)\ell_0}^B - w_{N,i}^B|}{|z - w_{n,i+(n-N)\ell_0}^B||z - w_{N,i}^B|} + \sum_{i=0}^{a+N\ell_0-1} \frac{|\tilde{w}_{n,i}^B - \tilde{w}_{N,i}^B|}{|z - \tilde{w}_{n,i}^B||z - \tilde{w}_{N,i}^B|}. \end{aligned}$$

Again the denominators are bounded below by  $\frac{3}{4}$  for  $z \in \mathcal{C}$  and we use Lemma 2.1 to bound the numerators. For the first term in the first sum, using

$$(19) \quad w_{n,0}^B = (3, \overline{a_1, \dots, a_s, \mathbf{p}_n})$$

and

$$(20) \quad w_{N,0}^B = (3, \overline{a_1, \dots, a_s, \mathbf{p}_N}),$$

one can see that the successive terms  $w_{n,i}^B$  and  $w_{N,i}^B$  (up to  $i = a - 1$ ) coincide at least in the first  $rN$  partial quotients. This is also true for the second sum, where we have

$$w_{n,a+(n-N)\ell_0}^B = (3, \overline{\mathbf{p}_{n-N}, a_1, \dots, a_s, \mathbf{p}_N})$$

and

$$w_{N,a}^B = (3, \overline{\mathbf{p}_N, a_1, \dots, a_s}),$$

as well as for the third sum, where we can use Remark 4.2 and the continued fractions of (19) and (20). Hence,

$$|S_1(n, N, z) - S_1(N, N, z)| \leq \frac{80}{3} \ell_N \lambda^{rN-1} \stackrel{(8)}{\leq} 80r(N+1)\lambda^{rN-1}.$$

Finally,

$$c(n, z) \leq 80(2 + r(N+1))\lambda^{rN-1}.$$

□

In particular, Theorem 4.3 applied to the function  $f = 1$  gives:

**Corollary 4.4.** *Let  $B$  be any left branch  $\neq L$  of  $\mathcal{T}$  and  $N \geq 1$ . For all  $n \geq N$ , there exists  $K = K_{B,N} \in \mathbb{R}$  such that*

$$(21) \quad |\log \varepsilon_n^B - n \log \varepsilon_0^B - K| \leq \delta_1(r, N)$$

with  $\delta_1(r, N)$  and  $r$  as in (10).

The next corollary proves Theorem 1.1 from the introduction.

**Corollary 4.5.** *Let  $f$  be a modular function. For any left branch  $B \neq L$  of  $\mathcal{T}$ ,*

$$\lim_{n \rightarrow \infty} f^{nor}(w_n^B) = f^{nor}(w_0^B).$$

*Proof.* It follows from Theorem 4.3 and Corollary 4.4 that  $|f(w_n^B) - nf(w_0^B)|$  and  $|\log \varepsilon_n^B - n \log \varepsilon_0^B|$  are bounded above and below by absolute constants (not depending on  $n$ ). Then

$$0 = \lim_{n \rightarrow \infty} \frac{|f(w_n^B) - nf(w_0^B)|}{\log \varepsilon_n^B} = \lim_{n \rightarrow \infty} \left| \frac{f(w_n^B)}{\log \varepsilon_n^B} - \frac{f(w_0^B)}{\log \varepsilon_0^B} \right|.$$

□

## 5. INTERLACING PROPERTY

In this section we prove Theorem 1.2. We restrict again to a left but not the most left branch in what follows, but the argument applies in the same way to any branch, with the bound  $\delta_2(n, r)$  slightly modified. Hence Theorem 1.2 applies in fact to any branch of the Markov tree and it is a consequence of the next theorem whose proof is similar to the proof of Theorem 4.3.

**Theorem 5.1.** *Let  $f$  be a modular function. For every left branch  $B \neq L$  of the Markov tree  $\mathcal{T}$  and for all  $n \geq 1$ ,*

$$(22) \quad |f(w_{n+1}^B) - f(w_n^B) - f(w_0^B)| \leq \delta_2(n, r) \max_{z \in \mathcal{C}} |f(z)|$$

where

$$(23) \quad \delta_2(n, r) = \frac{80\pi}{3}(n+2)r \left( \frac{2}{1+\sqrt{5}} \right)^{2(rn-1)}$$

and  $r+1$  is the number of partial quotients of  $w_0^B$ .

*Proof.* We have

$$(24) \quad f(w_{n+1}^B) - f(w_n^B) = f(w_0^B) + \int_{\rho}^{\rho^2} f(z)R_1(n, z) dz + \int_{\rho}^{\rho^2} f(z)R_2(n, z) dz,$$



where

$$R_1(n, z) = \sum_{i=0}^{a-1} \left( \frac{1}{z - w_{n+1,i}^B} - \frac{1}{z - w_{n,i}^B} \right) + \sum_{i=a}^{\ell_n-1} \left( \frac{1}{z - w_{n+1,\ell_0+i}^B} - \frac{1}{z - w_{n,i}^B} \right) - \sum_{i=0}^{\ell_n-1} \left( \frac{1}{z - \tilde{w}_{n+1,i}^B} - \frac{1}{z - \tilde{w}_{n,i}^B} \right),$$

$$R_2(n, z) = \sum_{i=0}^{\ell_0-1} \left( \frac{1}{z - w_{n+1,a+i}^B} - \frac{1}{z - w_{0,i}^B} - \frac{1}{z - \tilde{w}_{n+1,\ell_n+i}^B} + \frac{1}{z - \tilde{w}_{0,i}^B} \right).$$

Next we give upper bounds for the norms of the two sums above when  $z \in \mathcal{C}$ . We set again  $\lambda = \left( \frac{2}{1+\sqrt{5}} \right)^2$ .

**Bound for  $|R_1(n, z)|$ .** For  $z \in \mathcal{C}$ , we have

$$|R_1(n, z)| \leq \sum_{i=0}^{a-1} \frac{|w_{n+1,i}^B - w_{n,i}^B|}{|z - w_{n+1,i}^B||z - w_{n,i}^B|} + \sum_{i=a}^{\ell_n-1} \frac{|w_{n+1,i+\ell_0}^B - w_{n,i}^B|}{|z - w_{n+1,i+\ell_0}^B||z - w_{n,i}^B|} + \sum_{i=0}^{a+n\ell_0-1} \frac{|\tilde{w}_{n+1,i}^B - \tilde{w}_{n,i}^B|}{|z - \tilde{w}_{n+1,i}^B||z - \tilde{w}_{n,i}^B|} + \sum_{i=a+n\ell_0}^{\ell_n-1} \frac{|\tilde{w}_{n+1,i}^B - \tilde{w}_{n,i}^B|}{|z - \tilde{w}_{n+1,i}^B||z - \tilde{w}_{n,i}^B|}.$$

As before we use the bound of  $\frac{3}{4}$  for the denominators and Lemma 2.1 for the numerators. In the first sum using

$$(25) \quad w_{n+1,0}^B = (3, \overline{a_1, \dots, a_s, \mathbf{p}_{n+1}})$$

and

$$(26) \quad w_{n,0}^B = (3, \overline{a_1, \dots, a_s, \mathbf{p}_n}),$$

one can see that the successive terms  $w_{n+1,i}^B$  and  $w_{n,i}^B$  (up to  $i = a - 1$ ) coincide at least in the first  $rn$  partial quotients. The same is true for the second sum, where

$$w_{n+1,a+\ell_0}^B = (3, \overline{\mathbf{p}_n, a_1, \dots, a_s, \mathbf{p}})$$

and

$$w_{n,a}^B = (3, \overline{\mathbf{p}_n, a_1, \dots, a_s}).$$

For the third and fourth sums, we use once again Remark 4.2 together with (25) and (26), and

$$w_{n+1,a+n\ell_0}^B = (3, \overline{a_1, \dots, a_s, \mathbf{p}_{n+1}})$$

and

$$w_{n,a+n\ell_0}^B = (3, \overline{a_1, \dots, a_s, \mathbf{p}_n})$$

respectively.

Hence

$$|R_1(n, z)| \leq \frac{80}{3} \ell_n \lambda^{rn-1} \stackrel{(8)}{\leq} 80r(n+1)\lambda^{rn-1}.$$

**Bound for  $|R_2(n, z)|$ .** In a similar way we bound this second sum when  $z \in \mathcal{C}$ :

$$|R_2(n, z)| \leq \sum_{i=0}^{\ell_0-1} \frac{|w_{n+1,a+i}^B - w_{0,i}^B|}{|z - w_{n+1,a+i}^B||z - w_{0,i}^B|} + \frac{|\tilde{w}_{n+1,\ell_n+i}^B - \tilde{w}_{0,i}^B|}{|z - \tilde{w}_{n+1,\ell_n+i}^B||z - \tilde{w}_{0,i}^B|}.$$

Again using

$$w_{n+1,a}^B = (3, \overline{\mathbf{p}_{n+1}})$$

and

$$(27) \quad w_{0,0}^B = (3, \overline{\mathbf{p}}),$$

one can see that all the successive terms  $w_{n+1,a+i}^B$  and  $w_{0,i}^B$  in the sum coincide at least in the first  $rn$  partial quotients. For the conjugate terms, one can see from (27) and

$$w_{n+1,\ell_n}^B = (3, \overline{\mathbf{p}, a_1, \dots, a_s, \mathbf{p}_n})$$

that  $-\tilde{w}_{n+1,\ell_n+i}^B$  and  $-\tilde{w}_{0,i}^B$  coincide as well in the first  $rn$  partial quotients. Hence

$$|R_2(n, z)| \leq \frac{80}{3} \ell_0 \lambda^{rn-1} \stackrel{(8)}{\leq} 80r \lambda^{rn-1}.$$

Therefore,

$$\begin{aligned} |f(w_{n+1}) - f(w_n) - f(w_0)| &\leq \int_{\rho}^{\rho^2} |f(z)| (|R_1(n, z)| + |R_2(n, z)|) |dz| \\ &\leq \delta_2(n, r) \max_{z \in \mathcal{C}} |f(z)| \end{aligned}$$

with

$$\delta_2(n, r) = \frac{80\pi}{3} r(n+2) \lambda^{rn-1}.$$

□

Theorem (5.1) applied to the function  $f = 1$  gives:

**Corollary 5.2.** *For every left branch  $B \neq L$  of  $\mathcal{T}$  and for all  $n \geq 1$ ,*

$$|\log \varepsilon_{n+1}^B - \log \varepsilon_n^B - \log \varepsilon_0^B| \leq \delta_2(n, r)$$

with  $\delta_2(n, r)$  and  $r$  as in (23).

We finish this section by giving the proof of Theorem 1.2 in the case that the branch  $B$  is any left branch  $\neq L$ . The proof of the general case goes along the same lines.

**Theorem 5.3.** *Let  $f$  be a modular function,  $B$  be any left branch  $\neq L$  of the Markov tree  $\mathcal{T}$ , and  $r+1$  be the number of partial quotients of  $w_0^B$ . There exists a constant  $N_f$  such that, for all  $n \geq N_f/r$ , the real and imaginary parts of  $f^{\text{nor}}(w_{n+1}^B)$  lie between the real and imaginary parts respectively of  $f^{\text{nor}}(w_0^B)$  and  $f^{\text{nor}}(w_n^B)$ .*

**Remark 5.4.** *By going to a large enough level down on the tree we can have either  $n \gg 0$  or  $r > N_f$ , depending on the branch. Hence from that level on the interlacing property for the values of  $f$  holds on any branch.*

*Proof.* By definition, the inequality

$$\operatorname{Re}(f^{nor}(w_{n+1}^B)) < \operatorname{Re}(f^{nor}(w_n^B))$$

holds if and only if

$$(28) \quad \operatorname{Re}(f(w_{n+1}^B)) \log \varepsilon_n^B < \operatorname{Re}(f(w_n^B)) \log \varepsilon_{n+1}^B.$$

Theorem 5.1 and Corollary 5.2 respectively imply that

$$\operatorname{Re}(f(w_{n+1}^B)) = \operatorname{Re}(f(w_n^B)) + \operatorname{Re}(f(w_0^B)) + \mu(n, r)$$

with  $|\mu(n, r)| \leq \delta_2(n, r) \max_{z \in \mathcal{C}} |f(z)|$  and

$$\log \varepsilon_{n+1}^B = \log \varepsilon_n^B + \log \varepsilon_0^B + \lambda(n, r)$$

with  $|\lambda(n, r)| \leq \delta_2(n, r)$ . Hence (28) is equivalent to

$$(29) \quad (\operatorname{Re}(f(w_0^B)) + \mu(n, r)) \log \varepsilon_n^B < \operatorname{Re}(f(w_n^B)) (\log \varepsilon_0^B + \lambda(n, r)).$$

Now Theorem 4.3 and Corollary 4.4 imply that there exist constants  $N_1(f) \geq 1$ ,  $K_1 = K_{1,B,N_1(f)}$  and  $K_f = K_{f,B,N_1(f)}$  such that, for  $nr \geq N_1(f)$ , the inequality (29) holds if and only if

$$(\operatorname{Re}(f(w_0^B)) + \mu(n, r))(n \log \varepsilon_0^B + K_1) < (n \operatorname{Re}(f(w_0^B)) + K_f)(\log \varepsilon_0^B + \lambda(n, r)),$$

i.e.

$$(30) \quad \operatorname{Re}(f^{nor}(w_0^B)) < \frac{K_f}{K_1} + \frac{\lambda(n, r)(K_f + n \operatorname{Re}(f(w_0^B))) - \mu(n, r)(n \log \varepsilon_0^B + K_1)}{K_1 \log \varepsilon_0^B}.$$

Since  $|\lambda(n, r)| \leq \delta_2(n, r)$ ,  $|\mu(n, r)| \leq \delta_2(n, r) \max_{z \in \mathcal{C}} |f(z)|$ , and  $n \delta_2(n, r) \rightarrow 0$ , as  $nr \rightarrow \infty$ , (30) is equivalent to

$$\operatorname{Re}(f^{nor}(w_0^B)) < \frac{K_f}{K_1}$$

for all  $nr$  greater than some constant  $N_f \geq N_1(f)$ . Since the inequality above does not depend on  $n$ , but only on  $B$  and  $f$ , we have either

$$\operatorname{Re}(f^{nor}(w_{n+1}^B)) < \operatorname{Re}(f^{nor}(w_n^B)),$$

or

$$\operatorname{Re}(f^{nor}(w_{n+1}^B)) > \operatorname{Re}(f^{nor}(w_n^B)),$$

simultaneously for all  $n \geq N_f/r$ . Together with Corollary 4.5, it follows that either

$$\operatorname{Re}(f^{nor}(w_0^B)) < \operatorname{Re}(f^{nor}(w_{n+1}^B)) < \operatorname{Re}(f^{nor}(w_n^B))$$

for all  $n \geq N_f/r$  or

$$\operatorname{Re}(f^{nor}(w_n^B)) < \operatorname{Re}(f^{nor}(w_{n+1}^B)) < \operatorname{Re}(f^{nor}(w_0^B)).$$

The same argument applies to the imaginary parts of  $f^{nor}(w_{n+1}^B)$ ,  $f^{nor}(w_n^B)$  and  $f^{nor}(w_0^B)$ . □

**Acknowledgements.** The authors thank Masanobu Kaneko for his comments on a previous version of the paper.

#### REFERENCES

- [1] M. Aigner, *Markov's Theorem and 100 years of the uniqueness conjecture*, Springer-Verlag 2013.
- [2] J. Bourgain, A. Gamburd, P. Sarnak, *Markoff triples and strong approximation*, *Comptes Rendus Math.* 354 (2016), no. 2, 131–135.
- [3] J. Bourgain, A. Gamburd, P. Sarnak, *Markoff Surfaces and Strong Approximation: 1*, arXiv:math/1607.01530 (2016).
- [4] A. Gosh, P. Sarnak, *Integral points on Markoff type cubic surfaces*, arXiv:math/1706.06712 (2017).
- [5] M. Hall, *On the sum and products of continued fractions*, *Annals of Math.* 48 (1947), 966–993.
- [6] H. Cohn, *Mathematical microcosm of geodesics, free groups and Markoff forms. Classical quantum models and arithmetic problems*, 69–97, *Lecture Notes in Pure and Appl. Math.* **92** Dekker, New York, 1984.
- [7] W. Duke, Ö. Imamoğlu, A. Toth, *Cycle integrals of the  $j$ -function and mock modular forms*, *Ann. of Math. (2)* 173 (2011), no. 2, 947–981.
- [8] M. Kaneko, *Observations on the ‘values’ of the elliptic modular function  $j(\tau)$  at real quadratics*, *Kyushu Journal of Mathematics* 63 (2009), no. 2, 353–364.
- [9] A.V. Malyshev, *Markov and Lagrange Spectra (survey of the literature)*, *Journal of Soviet Mathematics*, 1981, 16:1, 767–788.
- [10] A. Markoff, *Sur les formes quadratiques binaires indéfinies*, *Math. Ann.* 15 (1879), 381–406.
- [11] A. Markoff, *Sur les formes quadratiques binaires indéfinies. II*, *Math. Ann.* 17 (1880), 379–399.
- [12] K.E. Roth, *Rational approximations to algebraic numbers*, *Mathematika*, 2:1–20; corrigendum 168, 1955.

ETH, MATHEMATICS DEPT., CH-8092, ZÜRICH, SWITZERLAND  
*E-mail address:* paloma.bengoechea@math.ethz.ch

ETH, MATHEMATICS DEPT., CH-8092, ZÜRICH, SWITZERLAND  
*E-mail address:* ozlem@math.ethz.ch