

# Kummer Theory for Drinfeld Modules

Richard Pink<sup>1</sup>

February 17, 2012

## Abstract

Let  $\varphi$  be a Drinfeld  $A$ -module of characteristic  $\mathfrak{p}_0$  over a finitely generated field  $K$ . Previous articles determined the image of the absolute Galois group of  $K$  up to commensurability in its action on all prime-to- $\mathfrak{p}_0$  torsion points of  $\varphi$ , or equivalently, on the prime-to- $\mathfrak{p}_0$  adelic Tate module of  $\varphi$ . In this article we consider in addition a finitely generated torsion free  $A$ -submodule  $M$  of  $K$  for the action of  $A$  through  $\varphi$ . We determine the image of the absolute Galois group of  $K$  up to commensurability in its action on the prime-to- $\mathfrak{p}_0$  division hull of  $M$ , or equivalently, on the extended prime-to- $\mathfrak{p}_0$  adelic Tate module associated to  $\varphi$  and  $M$ .

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Extended Tate modules</b>	<b>3</b>
<b>3</b>	<b>Reduction steps</b>	<b>6</b>
<b>4</b>	<b>Previous results on Galois groups</b>	<b>8</b>
<b>5</b>	<b>The primitive case</b>	<b>12</b>
<b>6</b>	<b>The general case</b>	<b>15</b>

---

<sup>\*</sup>Dept. of Mathematics, ETH Zürich, CH-8092 Zürich, Switzerland, pink@math.ethz.ch

# 1 Introduction

Let  $F$  be a finitely generated field of transcendence degree 1 over the prime field  $\mathbb{F}_p$  of characteristic  $p > 0$ . Let  $A$  be the ring of elements of  $F$  which are regular outside a fixed place  $\infty$  of  $F$ . Let  $K$  be another field that is finitely generated over  $\mathbb{F}_p$ , and let  $K^{\text{sep}}$  be a separable closure of  $K$ . Write  $\text{End}(\mathbb{G}_{a,K}) = K[\tau]$  with  $\tau(x) = x^p$ . Let  $\varphi: A \rightarrow K[\tau]$ ,  $a \mapsto \varphi_a$  be a Drinfeld  $A$ -module of rank  $r \geq 1$  and characteristic  $\mathfrak{p}_0$ . Then either  $\mathfrak{p}_0$  is the zero ideal of  $A$  and  $\varphi$  is said to have generic characteristic; or  $\mathfrak{p}_0$  is a maximal ideal of  $A$  and  $\varphi$  is said to have special characteristic.

For brevity we call any maximal ideal of  $A$  a prime of  $A$ . For any prime  $\mathfrak{p} \neq \mathfrak{p}_0$  of  $A$  the  $\mathfrak{p}$ -adic Tate module  $T_{\mathfrak{p}}(\varphi)$  is a free module of rank  $r$  over the completion  $A_{\mathfrak{p}}$ , endowed with a continuous action of the Galois group  $\text{Gal}(K^{\text{sep}}/K)$ . The prime-to- $\mathfrak{p}_0$  adelic Tate module  $T_{\text{ad}}(\varphi) = \prod_{\mathfrak{p} \neq \mathfrak{p}_0} T_{\mathfrak{p}}(\varphi)$  is then a free module of rank  $r$  over  $A_{\text{ad}} = \prod_{\mathfrak{p} \neq \mathfrak{p}_0} A_{\mathfrak{p}}$  carrying a natural action of Galois. This action corresponds to a continuous homomorphism

$$(1.1) \quad \text{Gal}(K^{\text{sep}}/K) \rightarrow \text{Aut}_{A_{\text{ad}}}(T_{\text{ad}}(\varphi)) \cong \text{GL}_r(A_{\text{ad}}).$$

Its image  $\Gamma_{\text{ad}}$  was determined up to commensurability by Pink-Rütsche [7] and Devic-Pink [2]; for special cases see Theorems 1.6 and 4.4 below.

Let  $M \subset K$  be a finitely generated torsion free  $A$ -submodule of rank  $d$  for the action of  $A$  through  $\varphi$ . Then there is an associated prime-to- $\mathfrak{p}_0$  adelic Tate module  $T_{\text{ad}}(\varphi, M)$ , which is a free module of rank  $r+d$  over  $A_{\text{ad}}$  carrying a natural continuous action of  $\text{Gal}(K^{\text{sep}}/K)$ . This module lies in a natural Galois equivariant short exact sequence

$$(1.2) \quad 0 \longrightarrow T_{\text{ad}}(\varphi) \longrightarrow T_{\text{ad}}(\varphi, M) \longrightarrow M \otimes_A A_{\text{ad}} \longrightarrow 0.$$

Define  $\Gamma_{\text{ad},M}$  as the image of the continuous homomorphism

$$(1.3) \quad \text{Gal}(K^{\text{sep}}/K) \rightarrow \text{Aut}_{A_{\text{ad}}}(T_{\text{ad}}(\varphi, M)) \cong \text{GL}_{r+d}(A_{\text{ad}}).$$

Then the restriction to  $T_{\text{ad}}(\varphi)$  induces a surjective homomorphism  $\Gamma_{\text{ad},M} \twoheadrightarrow \Gamma_{\text{ad}}$ , whose kernel we denote by  $\Delta_{\text{ad},M}$ . Since the action on  $M \otimes_A A_{\text{ad}}$  is trivial, there is a natural inclusion

$$(1.4) \quad \Delta_{\text{ad},M} \hookrightarrow \text{Hom}_A(M, T_{\text{ad}}(\varphi)).$$

Any splitting of the sequence (1.2) induces an inclusion into the semidirect product

$$(1.5) \quad \Gamma_{\text{ad},M} \hookrightarrow \Gamma_{\text{ad}} \ltimes \text{Hom}_A(M, T_{\text{ad}}(\varphi)).$$

The aim of this article is to describe these subgroups up to commensurability.

In general the shape of these Galois groups is affected by the endomorphisms of  $\varphi$  over  $K^{\text{sep}}$ , and in special characteristic also by the endomorphisms of the restrictions of  $\varphi$  to all subrings of  $A$ . Any general results therefore involve further definitions and notation. In this introduction we avoid these and mention only a special case; the general case is addressed by Theorems 5.1 and 6.6 and 6.7. Parts (a) and (b) of the following result are due to Pink-Rütsche [7, Thm. 0.1], resp. Devic-Pink [2, Thm. 1.1], and part (c) is a special case of Theorem 5.1 below:

**Theorem 1.6** *Assume that  $\text{End}_{K^{\text{sep}}}(\varphi) = A$ , and in special characteristic also that  $\text{End}_{K^{\text{sep}}}(\varphi|B) = A$  for every integrally closed infinite subring  $B \subset A$ .*

- (a) *If  $\varphi$  has generic characteristic, then  $\Gamma_{\text{ad}}$  is open in  $\text{GL}_r(A_{\text{ad}})$ .*
- (b) *If  $\varphi$  has special characteristic, then  $\Gamma_{\text{ad}}$  is commensurable with  $\overline{\langle a_0 \rangle} \cdot \text{SL}_r(A_{\text{ad}})$  for some central element  $a_0 \in A$  that generates a positive power of  $\mathfrak{p}_0$ .*
- (c) *The inclusions (1.4) and (1.5) are both open.*

The method used to prove Theorem 1.6 (c) and its generalizations is an adaptation of the Kummer theory for semiabelian varieties from Ribet [10] and predecessors. The main ingredients are the above mentioned descriptions of  $\Gamma_{\text{ad}}$  and Poonen's tameness result [8] concerning the structure of  $K$  as an  $A$ -module via  $\varphi$ . A standard procedure would be to first prove corresponding results for  $\mathfrak{p}$ -division points for almost all primes  $\mathfrak{p} \neq \mathfrak{p}_0$  of  $A$ , and for  $\mathfrak{p}$ -power division points for all  $\mathfrak{p} \neq \mathfrak{p}_0$ , and then to combine these individual results by taking products, as in [10], [1], [5], [7], [2], [3]. Instead, we have found a shorter way by doing everything adelically from the start. The core of the argument is the proof of Lemma 5.3. Therein we avoid the explicit use of group cohomology by trivializing an implicit 1-cocycle with the help of a suitable central element of  $\Gamma_{\text{ad}}$ . On first reading the readers may want to restrict their attention to the case of Theorem 1.6, which requires only Section 2, a little from Section 4, and Section 5 with simplifications, avoiding Sections 3 and 6 entirely. Some of this was worked out in the Master's thesis of Häberli [3]. Our results generalize those of Chi-Li [1] and Li [5].

The notation and the assumptions of this introduction remain in force throughout the article.

## 2 Extended Tate modules

Following the usual convention in commutative algebra we let  $A_{(\mathfrak{p}_0)} \subset F$  denote the localization of  $A$  at  $\mathfrak{p}_0$ ; this is equal to  $F$  if and only if  $\varphi$  has generic characteristic. Observe that there is a natural isomorphism of  $A$ -modules

$$(2.1) \quad A_{(\mathfrak{p}_0)}/A \cong \bigoplus_{\mathfrak{p} \neq \mathfrak{p}_0} F_{\mathfrak{p}}/A_{\mathfrak{p}},$$

where the product is extended over all maximal ideals  $\mathfrak{p} \neq \mathfrak{p}_0$  of  $A$  and where  $F_{\mathfrak{p}}$  and  $A_{\mathfrak{p}}$  denote the corresponding completions of  $F$  and  $A$ . This induces a natural isomorphism for the prime-to- $\mathfrak{p}_0$  adelic completion of  $A$ :

$$(2.2) \quad A_{\text{ad}} := \text{End}_A(A_{(\mathfrak{p}_0)}/A) \cong \prod_{\mathfrak{p} \neq \mathfrak{p}_0} A_{\mathfrak{p}}.$$

As a consequence, for any torsion  $A$ -module  $X$  that is isomorphic to  $(A_{(\mathfrak{p}_0)}/A)^{\oplus n}$  for some integer  $n$ , the construction

$$(2.3) \quad T(X) := \text{Hom}_A(A_{(\mathfrak{p}_0)}/A, X)$$

yields a free  $A_{\text{ad}}$ -module of rank  $n$ . Reciprocally  $T(X)$  determines  $X$  completely up to a natural isomorphism  $X \cong T(X) \otimes_{A_{\text{ad}}} (A_{(\mathfrak{p}_0)}/A)$ . Thus any  $A$ -linear group action on  $X$  determines and is determined by the corresponding  $A_{\text{ad}}$ -linear group action on  $T(X)$ . Moreover, with

$$(2.4) \quad T_{\mathfrak{p}}(X) := \text{Hom}_A(F_{\mathfrak{p}}/A_{\mathfrak{p}}, X)$$

the decompositions (2.1) and (2.2) induce a decomposition

$$(2.5) \quad T(X) \cong \prod_{\mathfrak{p} \neq \mathfrak{p}_0} T_{\mathfrak{p}}(X)$$

This will give a concise way of defining the  $\mathfrak{p}$ -adic and adelic Tate modules associated to the given Drinfeld module  $\varphi$ .

We view  $K^{\text{sep}}$  as an  $A$ -module with respect to the action  $A \times K^{\text{sep}} \rightarrow K^{\text{sep}}$ ,  $(a, x) \mapsto \varphi_a(x)$  and are interested in certain submodules. One particular submodule is  $K$ . Let  $M$  be a finitely generated torsion free  $A$ -submodule of rank  $d \geq 0$  contained in  $K$ . Then the *prime-to- $\mathfrak{p}_0$  division hull of  $M$  in  $K^{\text{sep}}$*  is the  $A$ -submodule

$$(2.6) \quad \text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M) := \{x \in K^{\text{sep}} \mid \exists a \in A \setminus \mathfrak{p}_0 : \varphi_a(x) \in M\}.$$

Let  $\text{Div}_K^{(\mathfrak{p}_0)}(M)$  denote the intersection of  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)$  with  $K$ . For later use we recall the following result proved by Poonen [8, Lemma 5] when  $K$  is a global field and  $\varphi$  has generic characteristic, and by Wang [11] in general:

**Theorem 2.7**  $[\text{Div}_K^{(\mathfrak{p}_0)}(M) : M]$  is finite.

As a special case of the above, the prime-to- $\mathfrak{p}_0$  division hull of the zero module  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(\{0\})$  is the module of all prime-to- $\mathfrak{p}_0$  torsion points of  $\varphi$  in  $K^{\text{sep}}$ .

**Proposition 2.8** *There is a natural short exact sequence of  $A$ -modules*

$$0 \longrightarrow \text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(\{0\}) \longrightarrow \text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M) \longrightarrow M \otimes_A A_{(\mathfrak{p}_0)} \longrightarrow 1,$$

where the map on the right hand side is described by  $x \mapsto \varphi_a(x) \otimes \frac{1}{a}$  for any  $a \in A \setminus \mathfrak{p}_0$  satisfying  $\varphi_a(x) \in M$ .

**Proof.** The map is well-defined, because for any  $x \in \text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)$  and any  $a, b \in A \setminus \mathfrak{p}_0$  satisfying  $\varphi_a(x), \varphi_b(x) \in M$  we have

$$\begin{aligned} \varphi_a(x) \otimes \frac{1}{a} &= \varphi_a(x) \otimes \frac{b}{ab} = \varphi_b(\varphi_a(x)) \otimes \frac{1}{ab} \\ &\quad \| \\ \varphi_b(x) \otimes \frac{1}{b} &= \varphi_b(x) \otimes \frac{a}{ab} = \varphi_a(\varphi_b(x)) \otimes \frac{1}{ab}. \end{aligned}$$

The map is clearly  $A$ -linear, and one easily shows that its kernel is  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(\{0\})$ ; hence the sequence is left exact. The exactness on the right results from the fact

that for any  $a \in A \setminus \mathfrak{p}_0$  and any  $m \in M$ , the polynomial equation  $\varphi_a(x) = m$  is separable of positive degree and therefore has a solution in  $K^{\text{sep}}$ .  $\square$

Dividing by  $M$ , the exact sequence from Proposition 2.8 yields a natural short exact sequence of  $A$ -modules

$$(2.9) \quad 0 \longrightarrow \text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(\{0\}) \longrightarrow \text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)/M \longrightarrow M \otimes_A (A_{(\mathfrak{p}_0)}/A) \longrightarrow 0.$$

By the general theory of Drinfeld modules the module on the left is isomorphic to  $(A_{(\mathfrak{p}_0)}/A)^{\oplus r}$ , where  $r$  is the rank of  $\varphi$ . Using the functor  $T$  from (2.3), the *prime-to- $\mathfrak{p}_0$  adelic Tate module of  $\varphi$*  can be described canonically as

$$(2.10) \quad T_{\text{ad}}(\varphi) := T(\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(\{0\}))$$

and is a free  $A_{\text{ad}}$ -module of rank  $r$ . Since  $M$  is a projective  $A$ -module of rank  $d$ , the module on the right of (2.9) is isomorphic to  $(A_{(\mathfrak{p}_0)}/A)^{\oplus d}$ , and together it follows that the module in the middle is isomorphic to  $(A_{(\mathfrak{p}_0)}/A)^{\oplus(r+d)}$ . The *extended prime-to- $\mathfrak{p}_0$  adelic Tate module of  $\varphi$  and  $M$*

$$(2.11) \quad T_{\text{ad}}(\varphi, M) := T(\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)/M)$$

is therefore a free  $A_{\text{ad}}$ -module of rank  $r + d$ . Moreover, the exact sequence (2.9) yields a natural short exact sequence of  $A_{\text{ad}}$ -modules

$$(2.12) \quad 0 \longrightarrow T_{\text{ad}}(\varphi) \longrightarrow T_{\text{ad}}(\varphi, M) \longrightarrow M \otimes_A A_{\text{ad}} \longrightarrow 0.$$

All this decomposes uniquely as  $T_{\text{ad}}(\varphi) = \prod_{\mathfrak{p} \neq \mathfrak{p}_0} T_{\mathfrak{p}}(\varphi)$  etc. as in (2.5).

By construction there is a natural continuous action of the Galois group  $\text{Gal}(K^{\text{sep}}/K)$  on all modules and arrows in Proposition 2.8 and in (2.9). This induces a continuous action on the short exact sequence (2.12), which in turn determines the former two by the following fact:

**Proposition 2.13** *The action of  $\text{Gal}(K^{\text{sep}}/K)$  on  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)$  is completely determined by the action on  $T_{\text{ad}}(\varphi, M)$ .*

**Proof.** For any  $\sigma \in \text{Gal}(K^{\text{sep}}/K)$  the endomorphism  $x \mapsto \sigma(x) - x$  of  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)$  is trivial on  $M$ , because that module is contained in  $K$ . Also, the image of this endomorphism is contained in  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(\{0\})$ , because for any  $a \in A \setminus \mathfrak{p}_0$  with  $\varphi_a(x) \in M$  we have  $\varphi_a(\sigma(x) - x) = \sigma(\varphi_a(x)) - \varphi_a(x) = 0$ . Thus the endomorphism factors through a homomorphism  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)/M \rightarrow \text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(\{0\})$ . But by (2.9) the latter homomorphism is determined completely by the action of  $\sigma$  on  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)/M$ , and thus by the action of  $\sigma$  on  $T_{\text{ad}}(\varphi, M)$ , as desired.  $\square$

Let  $\Gamma_{\text{ad}}$  and  $\Gamma_{\text{ad}, M}$  denote the images of  $\text{Gal}(K^{\text{sep}}/K)$  acting on  $T_{\text{ad}}(\varphi)$  and  $T_{\text{ad}}(\varphi, M)$ , as in (1.1) and (1.3). Restricting to  $T_{\text{ad}}(\varphi)$  induces a surjective homomorphism  $\Gamma_{\text{ad}, M} \twoheadrightarrow \Gamma_{\text{ad}}$ , and we define  $\Delta_{\text{ad}, M}$  by the short exact sequence

$$(2.14) \quad 1 \longrightarrow \Delta_{\text{ad}, M} \longrightarrow \Gamma_{\text{ad}, M} \longrightarrow \Gamma_{\text{ad}} \longrightarrow 1.$$

For any  $m \in M$  take an element  $t \in T_{\text{ad}}(\varphi, M)$  with image  $m \otimes 1$  in  $M \otimes_A A_{\text{ad}}$ . Since any  $\delta \in \Delta_{\text{ad}, M}$  acts trivially on  $T_{\text{ad}}(\varphi)$ , the difference  $\delta(t) - t$  depends only on  $\delta$  and  $m$ . Since  $\delta$  also acts trivially on  $M \otimes_A A_{\text{ad}}$ , the difference lies in  $T_{\text{ad}}(\varphi)$  and therefore defines a map

$$(2.15) \quad \Delta_{\text{ad}, M} \times M \longrightarrow T_{\text{ad}}(\varphi), (\delta, m) \mapsto \langle \delta, m \rangle := \delta(t) - t.$$

By direct calculation this map is additive in  $\delta$  and  $A$ -linear in  $m$ . By the construction of  $\Delta_{\text{ad}, M}$  the adjoint of the pairing (2.15) is therefore a natural inclusion, already mentioned in (1.4):

$$(2.16) \quad \Delta_{\text{ad}, M} \hookrightarrow \text{Hom}_A(M, T_{\text{ad}}(\varphi)).$$

Let  $R := \text{End}_K(\varphi)$  denote the endomorphism ring of  $\varphi$  over  $K$ . This is an  $A$ -order in a finite dimensional division algebra over  $F$ . It acts naturally on  $K$  and  $K^{\text{sep}}$  and therefore on  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(\{0\})$  and  $T_{\text{ad}}(\varphi)$ , turning the latter two into modules over  $R_{\text{ad}} := R \otimes_A A_{\text{ad}}$ . As this action commutes with the action of  $\Gamma_{\text{ad}}$ , it leads to an inclusion

$$(2.17) \quad \Gamma_{\text{ad}} \subset \text{Aut}_{R_{\text{ad}}}(T_{\text{ad}}(\varphi)).$$

The decomposition (2.2) induces a decomposition  $R_{\text{ad}} = \prod_{\mathfrak{p} \neq \mathfrak{p}_0} R_{\mathfrak{p}}$  where  $R_{\mathfrak{p}} := R \otimes_A A_{\mathfrak{p}}$  acts naturally on  $T_{\mathfrak{p}}(\varphi)$ .

If  $M$  is an  $R$ -submodule of  $K$ , then  $R$  and hence  $R_{\text{ad}}$  also act on  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)$  and  $T_{\text{ad}}(\varphi, M)$ , and these actions commute with the action of  $\Gamma_{\text{ad}, M}$ . The inclusion (2.16) then factors through an inclusion

$$(2.18) \quad \Delta_{\text{ad}, M} \hookrightarrow \text{Hom}_R(M, T_{\text{ad}}(\varphi)).$$

Moreover, any  $R$ -equivariant splitting of the sequence (2.12) then induces an embedding into the semidirect product

$$(2.19) \quad \Gamma_{\text{ad}, M} \hookrightarrow \Gamma_{\text{ad}} \ltimes \text{Hom}_R(M, T_{\text{ad}}(\varphi)).$$

### 3 Reduction steps

For use in Section 6 we now discuss the behavior of extended Tate modules and their associated Galois groups under isogenies and under restriction of  $\varphi$  to subrings.

First consider another Drinfeld  $A$ -module  $\varphi'$  and an isogeny  $f: \varphi \rightarrow \varphi'$  defined over  $K$ . Recall that there exists an isogeny  $g: \varphi' \rightarrow \varphi$  such that  $g \circ f = \varphi_a$  for some non-zero  $a \in A$ . From this it follows that  $M' := f(M)$  is a torsion free finitely generated  $A$ -submodule of  $K$  for the action of  $A$  through  $\varphi'$ . Thus  $f$  induces  $A_{\text{ad}}$ -linear maps from the modules in (2.12) to those associated to  $\varphi'$  and  $M'$ . The existence of  $g$  implies that these maps are inclusions of finite index. Together these maps yield a commutative diagram of  $A_{\text{ad}}$ -modules with exact rows

$$(3.1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & T_{\text{ad}}(\varphi) & \longrightarrow & T_{\text{ad}}(\varphi, M) & \longrightarrow & M \otimes_A A_{\text{ad}} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & T_{\text{ad}}(\varphi') & \longrightarrow & T_{\text{ad}}(\varphi', M') & \longrightarrow & M' \otimes_A A_{\text{ad}} \longrightarrow 0. \end{array}$$

By construction all these maps are equivariant under  $\text{Gal}(K^{\text{sep}}/K)$ ; hence the images of Galois in each column are canonically isomorphic. If we denote the analogues of the groups  $\Delta_{\text{ad},M} \subset \Gamma_{\text{ad},M} \twoheadrightarrow \Gamma_{\text{ad}}$  associated to  $\varphi'$  and  $M'$  by  $\Delta_{\text{ad},M'}^{\varphi'} \subset \Gamma_{\text{ad},M'}^{\varphi'} \twoheadrightarrow \Gamma_{\text{ad}}^{\varphi'}$ , this means that we have a natural commutative diagram

$$(3.2) \quad \begin{array}{ccccc} \Gamma_{\text{ad}} & \longleftarrow & \Gamma_{\text{ad},M} & \supset & \Delta_{\text{ad},M} \hookrightarrow \text{Hom}_A(M, T_{\text{ad}}(\varphi)) \\ \parallel & & \parallel & & \parallel \\ \Gamma_{\text{ad}}^{\varphi'} & \longleftarrow & \Gamma_{\text{ad},M'}^{\varphi'} & \supset & \Delta_{\text{ad},M'}^{\varphi'} \hookrightarrow \text{Hom}_A(M', T_{\text{ad}}(\varphi')), \end{array}$$

where the vertical arrow on the right hand side is an inclusion of finite index.

Next let  $B$  be any integrally closed infinite subring of  $A$ . Then  $A$  is a finitely generated projective  $B$ -module of some rank  $s \geq 1$ . The restriction  $\psi := \varphi|B$  is therefore a Drinfeld  $B$ -module of rank  $rs$  over  $K$ , and the given  $A$ -module  $M$  of rank  $d$  becomes a  $B$ -module of rank  $ds$ . Moreover, since the characteristic of  $\varphi$  is by definition the kernel of the derivative map  $a \mapsto d\varphi_a$ , the characteristic of  $\psi$  is simply  $\mathfrak{q}_0 := \mathfrak{p}_0 \cap B$ . In analogy to (2.2) we have

$$(3.3) \quad B_{\text{ad}} := \text{End}_B(B_{(\mathfrak{q}_0)}/B) \cong \prod_{\mathfrak{q} \neq \mathfrak{q}_0} B_{\mathfrak{q}},$$

where the product is extended over all maximal ideals  $\mathfrak{q} \neq \mathfrak{q}_0$  of  $B$ . Thus

$$(3.4) \quad A \otimes_B B_{\text{ad}} \cong \prod_{\mathfrak{p} \nmid \mathfrak{q}_0} A_{\mathfrak{p}}$$

is in a natural way a factor ring of  $A_{\text{ad}}$ . More precisely, it is isomorphic to  $A_{\text{ad}}$  if the characteristic  $\mathfrak{p}_0$  and hence  $\mathfrak{q}_0$  is zero; otherwise it is obtained from  $A_{\text{ad}}$  by removing the finitely many factors  $A_{\mathfrak{p}}$  for all maximal ideals  $\mathfrak{p} \neq \mathfrak{p}_0$  of  $A$  above  $\mathfrak{q}_0$ . In particular we have a natural isomorphism  $A \otimes_B B_{\text{ad}} \cong A_{\text{ad}}$  if and only if  $\mathfrak{p}_0$  is the unique prime ideal of  $A$  above  $\mathfrak{q}_0$ .

**Proposition 3.5** *The exact sequence (2.12) for  $\psi$  and  $M$  is naturally isomorphic to that obtained from the exact sequence (2.12) for  $\varphi$  and  $M$  by tensoring with  $A \otimes_B B_{\text{ad}}$  over  $A_{\text{ad}}$ . In particular we have a commutative diagram with surjective vertical arrows*

$$\begin{array}{ccccccc} 0 & \longrightarrow & T_{\text{ad}}(\varphi) & \longrightarrow & T_{\text{ad}}(\varphi, M) & \longrightarrow & M \otimes_A A_{\text{ad}} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & T_{\text{ad}}(\psi) & \longrightarrow & T_{\text{ad}}(\psi, M) & \longrightarrow & M \otimes_B B_{\text{ad}} \longrightarrow 0. \end{array}$$

If  $\mathfrak{p}_0$  is the only prime ideal of  $A$  above  $\mathfrak{q}_0$ , the vertical arrows are isomorphisms.

**Proof.** According to (2.6) the prime-to- $\mathfrak{p}_0$  division hull of  $M$  with respect to  $\varphi$  and the prime-to- $\mathfrak{q}_0$  division hull of  $M$  with respect to  $\psi$  are

$$\begin{aligned} \text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M) &:= \{x \in K^{\text{sep}} \mid \exists a \in A \setminus \mathfrak{p}_0 : \varphi_a(x) \in M\}, \\ \text{Div}_{K^{\text{sep}}}^{(\mathfrak{q}_0)}(M) &:= \{x \in K^{\text{sep}} \mid \exists b \in B \setminus \mathfrak{q}_0 : \psi_b(x) \in M\}. \end{aligned}$$

Here the latter is automatically contained in the former, because any  $b \in B \setminus \mathfrak{q}_0$  with  $\psi_b(x) \in M$  is by definition an element  $a := b \in A \setminus \mathfrak{p}_0$  with  $\varphi_a(x) \in M$ . Thus  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{q}_0)}(M)/M$  is the subgroup of all elements of  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)/M$  that are annihilated by some element of  $B \setminus \mathfrak{q}_0$ . In other words, it is the subgroup of all prime-to- $\mathfrak{q}_0$  torsion with respect to  $B$ , or again, it is obtained from  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)/M$  by removing the  $\mathfrak{p}$ -torsion for all maximal ideals  $\mathfrak{p} \neq \mathfrak{p}_0$  of  $A$  above  $\mathfrak{q}_0$ . In the same way  $A \otimes_B (B_{(\mathfrak{q}_0)}/B)$  is isomorphic to the submodule of  $A_{(\mathfrak{p}_0)}/A$  obtained by removing the  $\mathfrak{p}$ -torsion for all  $\mathfrak{p} \mid \mathfrak{q}_0$ . The same process applied to the exact sequence (2.9) therefore yields the analogue for  $\psi$  and  $M$ . By definition the exact sequence (2.12) for  $\psi$  and  $M$  is obtained from this by applying the functor

$$X \mapsto \text{Hom}_B(B_{(\mathfrak{q}_0)}/B, X) \cong \text{Hom}_A(A \otimes_B (B_{(\mathfrak{q}_0)}/B), X)$$

analogous to (2.3). The total effect of this is simply to remove the  $\mathfrak{p}$ -primary factors for all  $\mathfrak{p} \mid \mathfrak{q}_0$  from the exact sequence (2.12) for  $\varphi$  and  $M$ , from which everything follows.  $\square$

By construction the diagram in Proposition 3.5 is equivariant under  $\text{Gal}(K^{\text{sep}}/K)$ . It therefore induces a natural commutative diagram of Galois groups

$$(3.6) \quad \begin{array}{ccccccc} \text{Aut}_{A_{\text{ad}}}(T_{\text{ad}}(\varphi)) & \supset & \Gamma_{\text{ad}} & \longleftarrow & \Gamma_{\text{ad},M} & \supset & \Delta_{\text{ad},M} \hookrightarrow \text{Hom}_A(M, T_{\text{ad}}(\varphi)) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \text{Aut}_{B_{\text{ad}}}(T_{\text{ad}}(\psi)) & \supset & \Gamma_{\text{ad}}^\psi & \longleftarrow & \Gamma_{\text{ad},M}^\psi & \supset & \Delta_{\text{ad},M}^\psi \hookrightarrow \text{Hom}_B(M, T_{\text{ad}}(\psi)), \end{array}$$

where the subgroups in the lower row are the analogues for  $\psi$  and  $M$  of those in the upper row. By construction the left two vertical arrows are surjective, and they are isomorphisms if  $\mathfrak{p}_0$  is the only prime ideal of  $A$  above  $\mathfrak{q}_0$ . In that case the rightmost vertical arrow is injective and it follows that the map  $\Delta_{\text{ad},M} \rightarrow \Delta_{\text{ad},M}^\psi$  is an isomorphism as well. In general one can only conclude that  $\Delta_{\text{ad},M}^\psi$  contains the image of  $\Delta_{\text{ad},M}$ . In any case the diagram (3.6) gives a precise way of determining  $\Gamma_{\text{ad},M}^\psi$  from  $\Gamma_{\text{ad},M}$ .

## 4 Previous results on Galois groups

In this section we recall some previous results on the Galois group  $\Gamma_{\text{ad}}$ . Its precise description up to commensurability depends on certain endomorphism rings. The endomorphism ring of a Drinfeld module of generic characteristic is always commutative, but in special characteristic it can be non-commutative. In the latter case it can grow on restricting  $\varphi$  to a subring  $B$  of  $A$ , and this effect can impose additional conditions on  $\Gamma_{\text{ad}}$ . The question of whether the endomorphism ring becomes stationary or grows indefinitely with  $B$  depends on the following property:

**Definition 4.1** *We call a Drinfeld  $A$ -module of special characteristic over  $K$  isotrivial if over  $K^{\text{sep}}$  it is isomorphic to a Drinfeld  $A$ -module defined over a finite field.*

The next definition is slightly ad hoc, but it describes particular kinds of Drinfeld modules to which we can reduce ourselves in all cases, allowing a unified treatment of Kummer theory later on.

**Definition 4.2** *We call the triple  $(A, K, \varphi)$  primitive if the following conditions hold:*

- (a)  $R := \text{End}_K(\varphi)$  is equal to  $\text{End}_{K^{\text{sep}}}(\varphi)$ .
- (b) The center of  $R$  is  $A$ .
- (c)  $R$  is a maximal  $A$ -order in  $R \otimes_A F$ .
- (d) If  $\varphi$  is non-isotrivial of special characteristic, then for every integrally closed infinite subring  $B \subset A$  we have  $\text{End}_{K^{\text{sep}}}(\varphi|B) = R$ .
- (e) If  $\varphi$  is isotrivial of special characteristic, then  $A = \mathbb{F}_p[a_0]$  with  $\varphi_{a_0} = \tau^{[k/\mathbb{F}_p]}$ , where  $k$  denotes the finite field of constants of  $K$ .

**Proposition 4.3** *Let  $A'$  denote the normalization of the center of  $\text{End}_{K^{\text{sep}}}(\varphi)$ .*

- (a) *There exist a Drinfeld  $A'$ -module  $\varphi': A' \rightarrow K^{\text{sep}}[\tau]$  and an isogeny  $f: \varphi \rightarrow \varphi'|A$  over  $K^{\text{sep}}$  such that  $A'$  is the center of  $\text{End}_{K^{\text{sep}}}(\varphi')$ .*
- (b) *The characteristic  $\mathfrak{p}'_0$  of any  $\varphi'$  as in (a) is a prime ideal of  $A'$  above the characteristic  $\mathfrak{p}_0$  of  $\varphi$ .*
- (c) *There exist a finite extension  $K' \subset K^{\text{sep}}$  of  $K$ , a Drinfeld  $A'$ -module  $\varphi': A' \rightarrow K'[\tau]$ , an isogeny  $f: \varphi \rightarrow \varphi'|A$  over  $K'$ , and an integrally closed infinite subring  $B \subset A'$  such that  $A'$  is the center of  $\text{End}_{K^{\text{sep}}}(\varphi')$  and  $(B, K', \varphi'|B)$  is primitive.*
- (d) *The subring  $B$  in (c) is unique unless  $\varphi$  is isotrivial of special characteristic, in which case it is never unique.*
- (e) *For any data as in (c) the characteristic  $\mathfrak{p}'_0$  of  $\varphi'$  is the unique prime ideal of  $A'$  above the characteristic  $\mathfrak{q}_0$  of  $\varphi'|B$ .*

**Proof.** Applying [2, Prop. 4.3] to  $\varphi$  and the center of  $\text{End}_{K^{\text{sep}}}(\varphi)$  yields a Drinfeld  $A$ -module  $\tilde{\varphi}: A \rightarrow K^{\text{sep}}[\tau]$  and an isogeny  $f: \varphi \rightarrow \tilde{\varphi}$  over  $K^{\text{sep}}$  such that  $A'$  is mapped into  $\text{End}_{K^{\text{sep}}}(\tilde{\varphi})$  under the isomorphism  $\text{End}_{K^{\text{sep}}}(\varphi) \otimes_A F \cong \text{End}_{K^{\text{sep}}}(\tilde{\varphi}) \otimes_A F$  induced by  $f$ . Then  $A' \otimes_A F$  is the center of  $\text{End}_{K^{\text{sep}}}(\tilde{\varphi}) \otimes_A F$ , and since  $A'$  is integrally closed, it follows that  $A'$  is the center of  $\text{End}_{K^{\text{sep}}}(\tilde{\varphi})$ . The tautological homomorphism  $A' \hookrightarrow \text{End}_{K^{\text{sep}}}(\tilde{\varphi}) \hookrightarrow K^{\text{sep}}[\tau]$  thus constitutes a Drinfeld  $A'$ -module  $\varphi'$  with  $\varphi'|A \cong \tilde{\varphi}$  and  $\text{End}_{K^{\text{sep}}}(\varphi') = \text{End}_{K^{\text{sep}}}(\tilde{\varphi})$ . In particular the center of  $\text{End}_{K^{\text{sep}}}(\varphi')$  is equal to  $A'$ , proving (a).

For (b) recall that the characteristic of  $\varphi'$  is the kernel of the derivative map  $a' \mapsto d\varphi'_{a'}$ . Calling it  $\mathfrak{p}'_0$ , the characteristic of  $\varphi'|A$  is then  $\mathfrak{p}'_0 \cap A$ . As the characteristic of a Drinfeld module is invariant under isogenies, it follows that  $\mathfrak{p}'_0$  lies above  $\mathfrak{p}_0$ , proving (b).

For the remainder of the proof we take any pair  $\varphi'$  and  $f$  as in (a). We also choose a finite extension  $K' \subset K^{\text{sep}}$  of  $K$  such that  $\varphi'$  and  $f$  are defined over  $K'$  and that  $\text{End}_{K'}(\varphi') = \text{End}_{K^{\text{sep}}}(\varphi')$ .

Suppose first that  $\varphi'$  has generic characteristic. Then  $\text{End}_{K'}(\varphi')$  is commutative and hence equal to  $A'$ , and so the triple  $(A', K', \varphi')$  is already primitive. This proves

(c) with  $B = A'$ . Also, for any integrally closed infinite subring  $B \subset A'$  the ring  $\text{End}_{K^{\text{sep}}}(\varphi'|B)$  is commutative and hence again equal to  $\text{End}_{K^{\text{sep}}}(\varphi') = A'$ . Thus  $(B, K', \varphi'|B)$  being primitive requires that  $B = A'$ , proving (d). Since  $B = A'$ , the assertion of (e) is then trivially true.

Suppose next that  $\varphi'$  is non-isotrivial of special characteristic. Then by [6, Thm. 6.2] there exists a unique integrally closed infinite subring  $B \subset A'$  such that  $B$  is the center of  $\text{End}_{K^{\text{sep}}}(\varphi'|B)$  and that  $\text{End}_{K^{\text{sep}}}(\varphi'|B') \subset \text{End}_{K^{\text{sep}}}(\varphi'|B)$  for every integrally closed infinite subring  $B' \subset A'$ . For use below we note that both properties are invariant under isogenies of  $\varphi'$ , because isogenies induce isomorphisms on the rings  $\text{End}_{K^{\text{sep}}}(\varphi'|B') \otimes_{B'} \text{Quot}(B')$ . By uniqueness it follows that  $B$ , too, is invariant under isogenies of  $\varphi'$ .

After replacing  $K'$  by a finite extension we may assume that  $\text{End}_{K'}(\varphi'|B) = \text{End}_{K^{\text{sep}}}(\varphi'|B)$ . Then the triple  $(B, K', \varphi'|B)$  satisfies the conditions in Definition 4.2 except that  $\text{End}_{K'}(\varphi'|B)$  may not be a maximal order in  $\text{End}_{K'}(\psi') \otimes_B \text{Quot}(B)$ . But applying [2, Prop. 4.3] to  $\varphi'|B$  and  $\text{End}_{K'}(\varphi'|B)$  yields a Drinfeld  $B$ -module  $\psi': B \rightarrow K'[\tau]$  and an isogeny  $g: \varphi'|B \rightarrow \psi'$  over  $K'$  such that  $\text{End}_{K'}(\psi')$  is a maximal order in  $\text{End}_{K'}(\psi') \otimes_B \text{Quot}(B)$  which contains  $\text{End}_{K'}(\varphi'|B)$ . By the preceding remarks we now find that  $(B, K', \psi')$  is primitive. Moreover, the composite homomorphism  $A' \hookrightarrow \text{End}_{K'}(\varphi') \hookrightarrow \text{End}_{K'}(\varphi'|B) \hookrightarrow \text{End}_{K'}(\psi') \hookrightarrow K'[\tau]$  constitutes a Drinfeld  $A'$ -module  $\varphi''$  with  $\varphi''|B \cong \psi'$ . After replacing  $(\varphi', f)$  by  $(\varphi'', g \circ f)$  the data then satisfies all the requirements of (c). Assertion (d) follows from the above stated uniqueness of  $B$ , and (e) follows from [6, Prop. 3.5].

It remains to consider the case where  $\varphi'$  is isotrivial of special characteristic. In this case we may assume that  $\varphi'$  is defined over the constant field  $k'$  of  $K'$ . Any endomorphism of  $\varphi'$  over  $K^{\text{sep}}$  is then defined over a finite extension of  $k'$ , but by assumption also over  $K'$ ; hence it is defined over  $k'$ . In other words we have  $\text{End}_{K^{\text{sep}}}(\varphi') \subset k'[\tau]$ . Since  $\tau^{[k'/\mathbb{F}_p]}$  lies in the center of the  $k'[\tau]$ , it thus corresponds to an element of the center of  $\text{End}_{K^{\text{sep}}}(\varphi')$ . As this center is equal to  $A'$  by assumption, there is therefore an element  $a_0 \in A'$  with  $\varphi'_{a_0} = \tau^{[k'/\mathbb{F}_p]}$ . Set  $B := \mathbb{F}_p[a_0] \subset A'$  which, being isomorphic to a polynomial ring, is an integrally closed infinite subring of  $A'$ . Then  $\text{End}_{K^{\text{sep}}}(\varphi'|B)$  is the commutant of  $\tau^{[k'/\mathbb{F}_p]}$  in  $K^{\text{sep}}[\tau]$  and hence just  $k'[\tau]$ . By a standard construction this is a maximal  $B$ -order in a (cyclic) central division algebra over  $\text{Quot}(B)$ ; hence  $(B, K', \varphi'|B)$  is primitive, proving (c). For (d) observe that replacing  $K'$  and  $k'$  by finite extensions amounts to replacing  $a_0$  by an arbitrary positive power  $a_0^i$ . Thus the ring  $B$  is really not unique in this case, proving (d). Finally, assertion (e) follows from [2, Prop. 6.4 (a)]. This finishes the proof of Proposition 4.3.  $\square$

Assume now that  $(A, K, \varphi)$  is primitive and that  $\varphi$  has rank  $r$ . Then  $R \otimes_A F$  is a central division algebra of dimension  $m^2$  over  $F$  for some factorization  $r = mn$ . Thus for all primes  $\mathfrak{p} \neq \mathfrak{p}_0$  of  $A$ , the ring  $R_{\mathfrak{p}} := R \otimes_A A_{\mathfrak{p}}$  is an  $A_{\mathfrak{p}}$ -order in the central simple algebra  $R \otimes_A F_{\mathfrak{p}}$  of dimension  $m^2$  over  $F_{\mathfrak{p}}$  and is isomorphic to the matrix ring  $\text{Mat}_{m \times m}(A_{\mathfrak{p}})$  for almost all  $\mathfrak{p}$ . Let  $D_{\mathfrak{p}}$  denote the commutant of  $R_{\mathfrak{p}}$  in  $\text{End}_{A_{\mathfrak{p}}}(T_{\mathfrak{p}}(\varphi))$ . This is an  $A_{\mathfrak{p}}$ -order in a central simple algebra of dimension  $n^2$  over  $F_{\mathfrak{p}}$  and is isomorphic to the matrix ring  $\text{Mat}_{n \times n}(A_{\mathfrak{p}})$  for almost all  $\mathfrak{p}$ . Let  $D_{\mathfrak{p}}^1$  denote the

multiplicative group of elements of  $D_{\mathfrak{p}}$  of reduced norm 1. Set

$$D_{\text{ad}} := \prod_{\mathfrak{p} \neq \mathfrak{p}_0} D_{\mathfrak{p}} \subset \text{End}_{A_{\text{ad}}}(T_{\text{ad}}(\varphi))$$

and

$$D_{\text{ad}}^1 := \prod_{\mathfrak{p} \neq \mathfrak{p}_0} D_{\mathfrak{p}}^1 \subset D_{\text{ad}}^\times \subset \text{Aut}_{A_{\text{ad}}}(T_{\text{ad}}(\varphi)).$$

If  $\varphi$  has generic characteristic, we have  $m = 1$  and therefore  $D_{\mathfrak{p}} = \text{End}_{A_{\mathfrak{p}}}(T_{\mathfrak{p}}(\varphi)) \cong \text{Mat}_{r \times r}(A_{\mathfrak{p}})$  for all  $\mathfrak{p}$ .

If  $\varphi$  is non-isotrivial of special characteristic  $\mathfrak{p}_0$ , let  $a_0$  be any element of  $A$  that generates a positive power of  $\mathfrak{p}_0$ . If  $\varphi$  is isotrivial, the element  $a_0$  from Definition 4.2 (d) already has the same property. In both cases we view  $a_0$  as a scalar element of  $D_{\text{ad}}^\times$  via the diagonal embedding  $A \subset A_{\text{ad}} \subset D_{\text{ad}}$ , and let  $\overline{\langle a_0 \rangle}$  denote the pro-cyclic subgroup that is topologically generated by it.

In general the group  $\Gamma_{\text{ad}}$  was described up to commensurability in Pink-Rütsche [7] and Devic-Pink [2]. In the primitive case [7, Thm. 0.1] and [2, Thm. 1.1, Prop. 6.3] imply:

**Theorem 4.4** *Assume that  $(A, K, \varphi)$  is primitive.*

- (a) *If  $\varphi$  has generic characteristic, then  $\Gamma_{\text{ad}}$  is open in  $D_{\text{ad}}^\times$ .*
- (b) *If  $\varphi$  is non-isotrivial of special characteristic, then  $n \geq 2$  and  $\Gamma_{\text{ad}}$  is commensurable with  $\overline{\langle a_0 \rangle} \cdot D_{\text{ad}}^1$ .*
- (c) *If  $\varphi$  is isotrivial of special characteristic, then  $n = 1$  and  $\Gamma_{\text{ad}} = \overline{\langle a_0 \rangle}$  with  $a_0$  from 4.2 (c).*

**Corollary 4.5** *Assume that  $(A, K, \varphi)$  is primitive.*

- (a) *Let  $\Theta_{\text{ad}}$  denote the closure of the  $\mathbb{F}_p$ -subalgebra of  $D_{\text{ad}}$  generated by  $\Gamma_{\text{ad}}$ . Then there exists a non-zero ideal  $\mathfrak{a}$  of  $A$  with  $\mathfrak{a} \not\subset \mathfrak{p}_0$  such that  $\mathfrak{a}D_{\text{ad}} \subset \Theta_{\text{ad}}$ .*
- (b) *There exist a scalar element  $\gamma \in \Gamma_{\text{ad}}$  and a non-zero ideal  $\mathfrak{b}$  of  $A$  with  $\mathfrak{b} \not\subset \mathfrak{p}_0$  such that  $\gamma \equiv 1$  modulo  $\mathfrak{b}A_{\text{ad}}$  but not modulo  $\mathfrak{p}\mathfrak{b}A_{\text{ad}}$  for any prime  $\mathfrak{p} \neq \mathfrak{p}_0$  of  $A$ .*

**Proof.** Any open subgroup of  $D_{\text{ad}}^\times$ , and for  $n \geq 2$  any open subgroup of  $D_{\text{ad}}^1$ , generates an open subring of  $D_{\text{ad}}$ . Thus the assertion (a) follows from Theorem 4.4 unless  $\varphi$  is isotrivial of special characteristic. But in that case we have  $\Theta_{\text{ad}} = \overline{\mathbb{F}_p[a_0]} = A_{\text{ad}} = D_{\text{ad}}$  and (a) follows as well.

In generic characteristic the assertion (b) follows directly from the openness of  $\Gamma_{\text{ad}}$ . In special characteristic some positive power  $a_0^i$  lies in  $\Gamma_{\text{ad}}$ , and so (b) holds with  $\gamma = a_0^i$  and the ideal  $\mathfrak{b} = (a_0^i - 1)$ .  $\square$

## 5 The primitive case

Now we prove the following result, of which Theorem 1.6 (c) is a special case:

**Theorem 5.1** *Assume that  $(A, K, \varphi)$  is primitive. Set  $R := \text{End}_K(\varphi)$  and let  $M$  be a finitely generated torsion free  $R$ -submodule of  $K$ . Then the inclusions  $\Delta_{\text{ad}, M} \subset \text{Hom}_R(M, T_{\text{ad}}(\varphi))$  and  $\Gamma_{\text{ad}, M} \subset \Gamma_{\text{ad}} \ltimes \text{Hom}_R(M, T_{\text{ad}}(\varphi))$  are both open.*

So assume that  $(A, K, \varphi)$  is primitive. Let the subring  $\Theta_{\text{ad}} \subset D_{\text{ad}}$ , the element  $\gamma \in \Gamma_{\text{ad}}$ , and the ideals  $\mathfrak{a}, \mathfrak{b} \subset A$  be as in Corollary 4.5. Since  $M \subset \text{Div}_K^{(\mathfrak{p}_0)}(M)$  has finite index by Theorem 2.7, we can also choose a non-zero ideal  $\mathfrak{c}$  of  $A$  with  $\mathfrak{c} \not\subset \mathfrak{p}_0$  such that  $\mathfrak{c} \cdot \text{Div}_K^{(\mathfrak{p}_0)}(M) \subset M$ . With this data we prove the following more precise version of Theorem 5.1:

**Theorem 5.2** *In the above situation we have  $\mathfrak{abc} \cdot \text{Hom}_R(M, T_{\text{ad}}(\varphi)) \subset \Delta_{\text{ad}, M}$ .*

In the rest of this section we abbreviate  $T_{\text{ad}} := T_{\text{ad}}(\varphi)$  and  $M_{\text{ad}}^* := \text{Hom}_R(M, T_{\text{ad}})$ . Recall that the embedding  $\Delta_{\text{ad}, M} \subset M_{\text{ad}}^*$  is adjoint to the pairing  $\langle \cdot, \cdot \rangle$  from (2.15). The arithmetic part of the proof is a calculation in  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)$  with the following result:

**Lemma 5.3** *For any prime  $\mathfrak{p} \neq \mathfrak{p}_0$  of  $A$  and any element  $m \in M$  satisfying  $\langle \Delta_{\text{ad}, M}, m \rangle \subset \mathfrak{pbc}T_{\text{ad}}$  we have  $m \in \mathfrak{p}M$ .*

**Proof.** The assumption on  $\gamma$ , viewed as an element of  $A_{\text{ad}}$ , means that  $\gamma - 1 \in \mathfrak{b}A_{\text{ad}} \setminus \mathfrak{pb}A_{\text{ad}}$ . By the Chinese remainder theorem we can find an element  $b \in A \setminus \mathfrak{p}_0$  satisfying  $b \equiv \gamma - 1$  modulo  $\mathfrak{pbc}A_{\text{ad}}$ . Then by construction we have  $b \in \mathfrak{b} \setminus \mathfrak{pb}$ , and  $\gamma$  acts on all  $\mathfrak{pbc}$ -torsion points of  $\varphi$  through the action of  $1 + b \in A$ . By the Chinese remainder theorem we can also find elements  $a, c \in A \setminus \mathfrak{p}_0$  with  $a \in \mathfrak{p} \setminus \mathfrak{p}^2$  and  $c \in \mathfrak{c} \setminus \mathfrak{pc}$ . Then the product  $abc$  lies in  $\mathfrak{pbc} \setminus (\mathfrak{p}^2\mathfrak{bc} \cup \mathfrak{p}_0)$ . In particular the order of  $abc$  at  $\mathfrak{p}$  is equal to that of  $\mathfrak{pbc}$ , and so we can also choose an element  $d \in A \setminus \mathfrak{p}$  such that  $d\mathfrak{pbc} \subset (abc)$ .

For better readability we abbreviate the action of any element  $e \in A$  on an element  $x \in K^{\text{sep}}$  by  $ex := \varphi_e(x)$ . Since  $abc \in A \setminus \mathfrak{p}_0$ , we can select an element  $\tilde{m} \in \text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)$  with  $abc\tilde{m} = m$ . Then  $\tilde{m} := d\tilde{m}$  is an element of  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)$  which satisfies  $abc\tilde{m} = dm$ . By construction  $\tilde{m}$  lies in  $K^{\text{sep}}$ , but we shall see that it actually lies in a specific subfield.

Namely, choosing a compatible system of division points of  $\tilde{m}$  we can find an  $A$ -linear map  $\tilde{t}: A_{(\mathfrak{p}_0)} \rightarrow \text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)$  satisfying  $\tilde{t}(\frac{1}{abc}) = \tilde{m}$ . Then  $\tilde{t}(1) = abc\tilde{m} = dm$  lies in  $M$ ; hence  $\tilde{t}$  induces an  $A$ -linear map  $t: A_{(\mathfrak{p}_0)}/A \rightarrow \text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)/M$ . By the construction (2.11) this map is an element of  $T_{\text{ad}}(\varphi, M)$  whose image in  $M \otimes_A A_{\text{ad}}$  is  $dm \otimes 1$ . For any  $\delta \in \Delta_{\text{ad}, M}$  the definition (2.15) of the pairing now says that  $\langle \delta, dm \rangle = \delta(t) - t$ . But the assumption  $\langle \Delta_{\text{ad}, M}, m \rangle \subset \mathfrak{pbc}T_{\text{ad}}$  implies that  $\langle \Delta_{\text{ad}, M}, dm \rangle \subset d\mathfrak{pbc}T_{\text{ad}} \subset abcT_{\text{ad}}$  and therefore  $\delta(t) - t \in abcT_{\text{ad}}$ . Thus  $\delta(t) - t$  is the multiple by  $abc$  of an  $A$ -linear map  $A_{(\mathfrak{p}_0)}/A \rightarrow \text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(\{0\})$ ; hence it is zero on the residue class of  $\frac{1}{abc}$ . By the construction of  $t$  this means that  $\delta(\tilde{m}) - \tilde{m} = 0$ . Varying  $\delta$  we conclude

that  $\tilde{m}$  is fixed by  $\Delta_{\text{ad}, M}$ ; in other words, it lies in the subfield  $K_{\text{ad}} \subset K^{\text{sep}}$  with  $\text{Gal}(K_{\text{ad}}/K) = \Gamma_{\text{ad}}$ .

Now consider any element  $\sigma \in \text{Gal}(K^{\text{sep}}/K)$ . The fact that  $m$  lies in  $K$  implies that

$$abc(\sigma - 1)(\tilde{m}) = (\sigma - 1)(abc\tilde{m}) = (\sigma - 1)(m) = 0.$$

Thus  $(\sigma - 1)(\tilde{m})$  is annihilated by  $abc$  and hence by the ideal  $d\mathfrak{p}\mathfrak{bc} \subset (abc)$ . The element  $(\sigma - 1)(\tilde{m}) = d(\sigma - 1)(\tilde{m})$  is therefore annihilated by the ideal  $\mathfrak{p}\mathfrak{bc}$ . Since  $\gamma$  acts on all  $\mathfrak{p}\mathfrak{bc}$ -torsion points through the action of  $1 + b \in A$ , it follows that

$$(1 + b - \gamma)((\sigma - 1)(\tilde{m})) = 0.$$

On the other hand we have  $\tilde{m} \in K_{\text{ad}}$ , and since  $\gamma$  lies in the center of  $\Gamma_{\text{ad}}$ , its action on  $K_{\text{ad}}$  commutes with the action of  $\sigma$  on  $K_{\text{ad}}$ . Thus the last equation is equivalent to

$$(\sigma - 1)((1 + b - \gamma)(\tilde{m})) = 0.$$

As  $\sigma \in \text{Gal}(K^{\text{sep}}/K)$  was arbitrary, it follows that  $(1 + b - \gamma)(\tilde{m}) \in K$ .

Since  $\tilde{m}$  lies in  $\text{Div}_{K^{\text{sep}}}^{(\mathfrak{p}_0)}(M)$ , we can now deduce that  $(1 + b - \gamma)(\tilde{m}) \in \text{Div}_K^{(\mathfrak{p}_0)}(M)$ . By the choice of  $\mathfrak{c}$  and  $c$  it follows that  $c(1 + b - \gamma)(\tilde{m}) \in M$ . The fact that  $dm = abc\tilde{m}$  thus implies that

$$(1 + b - \gamma)(dm) = (1 + b - \gamma)(abc\tilde{m}) = abc(1 + b - \gamma)(\tilde{m}) \in abM.$$

But  $dm$  is an element of  $K$  and therefore satisfies  $(1 - \gamma)(dm) = 0$ . Thus the last relation shows that actually  $b dm \in abM$  and so  $dm \in aM$ . Since  $a \in \mathfrak{p}$  and  $d \notin \mathfrak{p}$ , this implies that  $m \in \mathfrak{p}M$ , as desired.  $\square$

The rest of the proof of Theorem 5.2 involves rings and modules. It is easier to understand if  $R = A$ , in which case  $D_{\mathfrak{p}} \cong \text{Mat}_{r \times r}(A_{\mathfrak{p}})$  for all  $\mathfrak{p}$ , so the readers may want to restrict themselves to that case on first reading. For general facts on maximal orders in semisimple algebras see Reiner [9].

Using [9, Cor. 11.6], the assumptions 4.2 imply that for any prime  $\mathfrak{p} \neq \mathfrak{p}_0$  of  $A$  the ring  $R_{\mathfrak{p}} := R \otimes_A A_{\mathfrak{p}}$  is a maximal order in a finite dimensional central simple algebra over  $F_{\mathfrak{p}}$ . By [9, Thm. 17.3] we can therefore identify it with the matrix ring  $\text{Mat}_{n_{\mathfrak{p}} \times n_{\mathfrak{p}}}(S_{\mathfrak{p}})$ , where  $S_{\mathfrak{p}}$  is the maximal order in a finite dimensional central division algebra over  $F_{\mathfrak{p}}$ . Here  $n_{\mathfrak{p}} \geq 1$  may vary with  $\mathfrak{p}$ . Let  $L_{\mathfrak{p}} := S_{\mathfrak{p}}^{\oplus n_{\mathfrak{p}}}$  denote the tautological left  $R_{\mathfrak{p}}$ -module. Since  $T_{\mathfrak{p}} := T_{\mathfrak{p}}(\varphi)$  is a non-trivial finitely generated torsion free left  $R_{\mathfrak{p}}$ -module, it is isomorphic to  $L_{\mathfrak{p}}^{\oplus m_{\mathfrak{p}}}$  for some  $m_{\mathfrak{p}} \geq 1$  by [9, Thm. 18.10]. Thus  $D_{\mathfrak{p}} := \text{End}_{R_{\mathfrak{p}}}(T_{\mathfrak{p}})$  is isomorphic to the matrix ring  $\text{Mat}_{m_{\mathfrak{p}} \times m_{\mathfrak{p}}}(S_{\mathfrak{p}}^{\text{opp}})$  over the opposite algebra  $S_{\mathfrak{p}}^{\text{opp}}$ . Let  $N_{\mathfrak{p}} := (S_{\mathfrak{p}}^{\text{opp}})^{\oplus m_{\mathfrak{p}}}$  denote the tautological left  $D_{\mathfrak{p}}$ -module; then as a  $D_{\mathfrak{p}}$ -module  $T_{\mathfrak{p}}$  is isomorphic to  $N_{\mathfrak{p}}^{\oplus n_{\mathfrak{p}}}$ . Moreover, by biduality, using Morita equivalence [9, Thm. 16.14] or direct computation, we have

$$(5.4) \quad R_{\mathfrak{p}} \cong \text{End}_{D_{\mathfrak{p}}}(T_{\mathfrak{p}}).$$

Next, since  $R$  is a maximal order in a division algebra over the Dedekind ring  $A$ , and  $M$  is a finitely generated torsion free  $R$ -module,  $M$  is a projective  $R$ -module by [9,

Cor. 21.5], say of rank  $\ell \geq 0$ . For each  $\mathfrak{p} \neq \mathfrak{p}_0$  we therefore have  $M_{\mathfrak{p}} := M \otimes_A A_{\mathfrak{p}} \cong R_{\mathfrak{p}}^{\oplus \ell}$  as an  $R_{\mathfrak{p}}$ -module. Consequently  $M_{\mathfrak{p}}^* := \text{Hom}_R(M, T_{\mathfrak{p}}) \cong T_{\mathfrak{p}}^{\oplus \ell}$  as a  $D_{\mathfrak{p}}$ -module via the action of  $D_{\mathfrak{p}}$  on  $T_{\mathfrak{p}}$ . Using the biduality (5.4) we obtain a natural isomorphism

$$(5.5) \quad M_{\mathfrak{p}} \cong \text{Hom}_{D_{\mathfrak{p}}}(M_{\mathfrak{p}}^*, T_{\mathfrak{p}}).$$

Taking the product over all  $\mathfrak{p} \neq \mathfrak{p}_0$  yields adelic versions of all this with  $T_{\text{ad}} = \prod T_{\mathfrak{p}}$  and  $R_{\text{ad}} = \prod R_{\mathfrak{p}}$  and  $D_{\text{ad}} = \prod D_{\mathfrak{p}}$  and  $M_{\text{ad}}^* = \prod M_{\mathfrak{p}}^*$ .

Recall that  $\Delta_{\text{ad}, M}$  is a closed additive subgroup of  $M_{\text{ad}}^* = \prod M_{\mathfrak{p}}^*$ . Let  $\Delta_{\mathfrak{p}}$  denote its image under the projection to  $M_{\mathfrak{p}}^*$ .

**Lemma 5.6** *For any  $\mathfrak{p} \neq \mathfrak{p}_0$  and any  $D_{\mathfrak{p}}$ -linear map  $f: M_{\mathfrak{p}}^* \rightarrow N_{\mathfrak{p}}$  satisfying  $f(\Delta_{\mathfrak{p}}) \subset \mathfrak{p}\mathfrak{b}\mathfrak{c}N_{\mathfrak{p}}$ , we have  $f(M_{\mathfrak{p}}^*) \subset \mathfrak{p}N_{\mathfrak{p}}$ .*

**Proof.** Since  $N_{\mathfrak{p}}$  is a  $D_{\mathfrak{p}}$ -module isomorphic to a direct summand of  $T_{\mathfrak{p}}$ , it is equivalent to show that for every  $D_{\mathfrak{p}}$ -linear map  $g: M_{\mathfrak{p}}^* \rightarrow T_{\mathfrak{p}}$  with  $g(\Delta_{\mathfrak{p}}) \subset \mathfrak{p}\mathfrak{b}\mathfrak{c}T_{\mathfrak{p}}$  we have  $g(M_{\mathfrak{p}}^*) \subset \mathfrak{p}T_{\mathfrak{p}}$ . Let  $\langle \cdot, \cdot \rangle: M_{\mathfrak{p}}^* \times M_{\mathfrak{p}} \rightarrow T_{\mathfrak{p}}$  denote the natural  $A_{\mathfrak{p}}$ -bilinear map. Then the biduality (5.5) says that  $g = \langle \underline{\phantom{x}}, m_{\mathfrak{p}} \rangle$  for an element  $m_{\mathfrak{p}} \in M_{\mathfrak{p}}$ . Write  $\mathfrak{p}\mathfrak{b}\mathfrak{c} = \mathfrak{p}^i\mathfrak{d}$  for an integer  $i \geq 1$  and an ideal  $\mathfrak{d}$  of  $A$  that is prime to  $\mathfrak{p}$ . Choose any element  $m \in M$  which is congruent to  $m_{\mathfrak{p}}$  modulo  $\mathfrak{p}^iM_{\mathfrak{p}}$  and congruent to 0 modulo  $\mathfrak{d}M$ . Then the assumption  $\langle \Delta_{\mathfrak{p}}, m_{\mathfrak{p}} \rangle = g(\Delta_{\mathfrak{p}}) \subset \mathfrak{p}\mathfrak{b}\mathfrak{c}T_{\mathfrak{p}}$  implies that  $\langle \Delta_{\text{ad}, M}, m \rangle \subset \mathfrak{p}\mathfrak{b}\mathfrak{c}T_{\text{ad}}$ . By Lemma 5.3 it follows that  $m \in \mathfrak{p}M$ . Consequently  $m_{\mathfrak{p}} \in \mathfrak{p}M_{\mathfrak{p}}$  and therefore  $g(M_{\mathfrak{p}}^*) = \langle \Delta_{\mathfrak{p}}, m_{\mathfrak{p}} \rangle \subset \mathfrak{p}T_{\mathfrak{p}}$ , as desired.  $\square$

Now observe that  $\Delta_{\text{ad}, M} \subset M_{\text{ad}}^*$  is a closed additive subgroup that is invariant under the action of  $\Gamma_{\text{ad}}$ . It is therefore a submodule with respect to the subring  $\Theta_{\text{ad}} := \overline{\mathbb{F}_p[\Gamma_{\text{ad}}]}$  of  $D_{\text{ad}}$  from Corollary 4.5 (a). By 4.5 (a) we therefore have

$$(5.7) \quad \Delta'_{\text{ad}} := \mathfrak{a}D_{\text{ad}}\Delta_{\text{ad}, M} \subset \Delta_{\text{ad}, M}.$$

By construction  $\Delta'_{\text{ad}}$  is a submodule over  $D_{\text{ad}} = \prod D_{\mathfrak{p}}$  and therefore itself a product  $\Delta'_{\text{ad}} = \prod \Delta'_{\mathfrak{p}}$  for  $D_{\mathfrak{p}}$ -submodules  $\Delta'_{\mathfrak{p}} \subset M_{\mathfrak{p}}^*$ .

**Lemma 5.8** *For any  $\mathfrak{p} \neq \mathfrak{p}_0$  and any  $D_{\mathfrak{p}}$ -linear map  $f: M_{\mathfrak{p}}^* \rightarrow N_{\mathfrak{p}}$  satisfying  $f(\Delta'_{\mathfrak{p}}) \subset \mathfrak{p}\mathfrak{a}\mathfrak{b}\mathfrak{c}N_{\mathfrak{p}}$ , we have  $f(M_{\mathfrak{p}}^*) \subset \mathfrak{p}N_{\mathfrak{p}}$ .*

**Proof.** The definition of  $\Delta'_{\mathfrak{p}}$  implies that  $\mathfrak{a}D_{\mathfrak{p}}\Delta_{\mathfrak{p}} \subset \Delta'_{\mathfrak{p}}$ . Thus by assumption we have  $\mathfrak{a}f(\Delta_{\mathfrak{p}}) \subset f(\mathfrak{a}D_{\mathfrak{p}}\Delta_{\mathfrak{p}}) \subset f(\Delta'_{\mathfrak{p}}) \subset \mathfrak{p}\mathfrak{a}\mathfrak{b}\mathfrak{c}N_{\mathfrak{p}}$  and therefore  $f(\Delta_{\mathfrak{p}}) \subset \mathfrak{p}\mathfrak{b}\mathfrak{c}N_{\mathfrak{p}}$ . By Lemma 5.6 this implies that  $f(M_{\mathfrak{p}}^*) \subset \mathfrak{p}N_{\mathfrak{p}}$ .  $\square$

**Lemma 5.9** *For any  $\mathfrak{p} \neq \mathfrak{p}_0$  we have  $\mathfrak{a}\mathfrak{b}\mathfrak{c}M_{\mathfrak{p}}^* \subset \Delta'_{\mathfrak{p}}$ .*

**Proof.** Let  $\mathfrak{m}_{\mathfrak{p}}$  denote the maximal ideal of  $S_{\mathfrak{p}}^{\text{opp}}$ . Then by [9, Thm. 13.2] we have  $\mathfrak{p}S_{\mathfrak{p}}^{\text{opp}} = \mathfrak{m}_{\mathfrak{p}}^e$  for some integer  $e \geq 1$ . The general theory says the following about the structure of the module  $M_{\mathfrak{p}}^*/\Delta'_{\mathfrak{p}}$  over the maximal order  $D_{\mathfrak{p}}$ . On the one hand, by Knebusch [4, Satz 7] the torsion submodule of  $M_{\mathfrak{p}}^*/\Delta'_{\mathfrak{p}}$  is a finite direct sum of indecomposable modules isomorphic to  $N_{\mathfrak{p}}/\mathfrak{m}_{\mathfrak{p}}^{j_{\nu}}N_{\mathfrak{p}}$  for certain integers  $j_{\nu} \geq 1$ . On

the other hand, the factor module of  $M_{\mathfrak{p}}^*/\Delta'_{\mathfrak{p}}$  by its torsion submodule is projective by [9, Cor. 21.5] and hence isomorphic to a direct sum of copies of  $N_{\mathfrak{p}}$ . That the factor module is projective also implies that  $M_{\mathfrak{p}}^*/\Delta'_{\mathfrak{p}}$  is isomorphic to the direct sum of its torsion submodule with the factor module. Together it follows that  $M_{\mathfrak{p}}^*/\Delta'_{\mathfrak{p}}$  is a finite direct sum of modules isomorphic to  $N_{\mathfrak{p}}$  or to  $N_{\mathfrak{p}}/\mathfrak{m}_{\mathfrak{p}}^{j_{\nu}}N_{\mathfrak{p}}$  for certain integers  $j_{\nu} \geq 1$ .

To use this fact, let  $\mathfrak{p}^i$  denote the highest power of  $\mathfrak{p}$  dividing  $\mathfrak{abc}$ . If no summand isomorphic to  $N_{\mathfrak{p}}$  occurs in  $M_{\mathfrak{p}}^*/\Delta'_{\mathfrak{p}}$  and all exponents  $j_{\nu}$  are  $\leq ei$ , then  $M_{\mathfrak{p}}^*/\Delta'_{\mathfrak{p}}$  is annihilated by  $\mathfrak{p}^i S_{\mathfrak{p}}^{\text{opp}} = \mathfrak{m}_{\mathfrak{p}}^{ei}$ . In this case it follows that  $\mathfrak{abc}M_{\mathfrak{p}}^* = \mathfrak{p}^i M_{\mathfrak{p}}^* \subset \Delta'_{\mathfrak{p}}$ , as desired.

Otherwise there exists a surjective  $D_{\mathfrak{p}}$ -linear map  $M_{\mathfrak{p}}^*/\Delta'_{\mathfrak{p}} \rightarrow N_{\mathfrak{p}}/\mathfrak{m}_{\mathfrak{p}}^{ei+1}N_{\mathfrak{p}}$ . Composed with the isomorphism

$$N_{\mathfrak{p}}/\mathfrak{m}_{\mathfrak{p}}^{ei+1}N_{\mathfrak{p}} \cong \mathfrak{m}_{\mathfrak{p}}^{e-1}N_{\mathfrak{p}}/\mathfrak{p}^{i+1}N_{\mathfrak{p}} = \mathfrak{m}_{\mathfrak{p}}^{e-1}N_{\mathfrak{p}}/\mathfrak{pabc}N_{\mathfrak{p}},$$

this yields a  $D_{\mathfrak{p}}$ -linear map  $M_{\mathfrak{p}}^*/\Delta'_{\mathfrak{p}} \rightarrow N_{\mathfrak{p}}/\mathfrak{pabc}N_{\mathfrak{p}}$  whose image is not contained in  $\mathfrak{p}N_{\mathfrak{p}}/\mathfrak{pabc}N_{\mathfrak{p}} = \mathfrak{m}_{\mathfrak{p}}^e N_{\mathfrak{p}}/\mathfrak{pabc}N_{\mathfrak{p}}$ . As  $M_{\mathfrak{p}}^*$  is a projective  $D_{\mathfrak{p}}$ -module, the latter map can be lifted to a  $D_{\mathfrak{p}}$ -linear map  $f: M_{\mathfrak{p}}^* \rightarrow N_{\mathfrak{p}}$ . By construction this map then satisfies  $f(\Delta'_{\mathfrak{p}}) \subset \mathfrak{pabc}N_{\mathfrak{p}}$  and  $f(M_{\mathfrak{p}}^*) \not\subset \mathfrak{p}N_{\mathfrak{p}}$ . But that contradicts Lemma 5.8; hence this case is not possible and the lemma is proved.  $\square$

Taking the product over all  $\mathfrak{p}$ , Lemma 5.9 and the inclusion (5.7) imply that  $\mathfrak{abc}M_{\text{ad}}^* \subset \Delta'_{\text{ad}} \subset \Delta_{\text{ad}, M}$ . This finishes the proof of Theorem 5.2. In particular it proves the first assertion of Theorem 5.1, from which the second assertion directly follows.

## 6 The general case

First we note the following general fact on homomorphisms of modules:

**Proposition 6.1** *Let  $S$  be a unitary ring, not necessarily commutative, and let  $M$  and  $N$  be left  $S$ -modules. Let  $X$  be a subset of  $M$  and  $SX$  the  $S$ -submodule generated by it. Let  $\text{Hom}_{(S)}(X, N)$  denote the set of maps  $\ell: X \rightarrow N$  such that for any finite collection of  $s_i \in S$  and  $x_i \in X$  with  $\sum_i s_i x_i = 0$  in  $M$  we have  $\sum_i s_i \ell(x_i) = 0$  in  $N$ .*

- (a) *The restriction of maps induces a bijection  $\text{Hom}_S(SX, N) \xrightarrow{\sim} \text{Hom}_{(S)}(X, N)$ .*
- (b) *If  $R$  is a unitary subring of  $S$  such that  $X$  is an  $R$ -submodule and the natural map  $S \otimes_R X \rightarrow M$ ,  $\sum_i s_i \otimes x_i \mapsto \sum_i s_i x_i$  is injective, then  $\text{Hom}_{(S)}(X, N) = \text{Hom}_R(X, N)$ .*
- (c) *If  $X$  is an  $S$ -submodule of  $M$ , then  $\text{Hom}_{(S)}(X, N) = \text{Hom}_S(X, N)$ .*

**Proof.** Let  $F := \bigoplus_{x \in X} S \cdot [x]$  be the free left  $S$ -module over the set  $X$  and consider the natural  $S$ -linear map  $F \rightarrow M$ ,  $\sum_i s_i [x_i] \mapsto \sum_i s_i x_i$ . Since  $S$  is unitary, the image of this map is  $SX$ . Let  $T$  denote its kernel. Then giving an  $S$ -linear map  $SX \rightarrow N$  is equivalent to giving an  $S$ -linear map  $F \rightarrow N$  which vanishes on  $T$ . Using the universal property of  $F$  we find that the latter is equivalent to giving an element of  $\text{Hom}_{(S)}(X, N)$ . The total correspondence is given by restriction of maps, proving (a).

In (b) we have  $S \otimes_R X \xrightarrow{\sim} SX$ ; hence the adjunction between tensor product and Hom yields bijections  $\text{Hom}_S(SX, N) \xrightarrow{\sim} \text{Hom}_S(S \otimes_R X, N) \xrightarrow{\sim} \text{Hom}_R(X, N)$ . Their composite is again just restriction of maps; so by (a) the restriction map  $\text{Hom}_{(S)}(X, N) \rightarrow \text{Hom}_R(X, N)$  is also bijective, proving (b).

Finally, (c) is a special case of (a) or (b), according to taste.  $\square$

Now we return to the situation of Section 4. We choose data  $(A', K', \varphi', f, B)$  as in Proposition 4.3 (c), that is: We let  $A'$  denote the normalization of the center of  $\text{End}_{K^{\text{sep}}}(\varphi)$ , take a finite extension  $K' \subset K^{\text{sep}}$  of  $K$ , a Drinfeld  $A'$ -module  $\varphi': A' \rightarrow K'[\tau]$ , an isogeny  $f: \varphi \rightarrow \varphi'|A$  over  $K'$ , and an integrally closed infinite subring  $B \subset A'$  such that  $A'$  is the center of  $\text{End}_{K^{\text{sep}}}(\varphi')$  and  $(B, K', \varphi'|B)$  is primitive. By Proposition 4.3 (e) the characteristic  $\mathfrak{p}'_0$  of  $\varphi'$  is then the only prime ideal of  $A'$  above the characteristic  $\mathfrak{q}_0$  of  $\varphi'|B$ . We will apply the reduction steps from Section 3 to the isogeny  $f$  and to each of the inclusions  $A \subset A' \supset B$ .

Specifically, let us set  $\psi' := \varphi'|B$  and  $S' := \text{End}_{K^{\text{sep}}}(\psi')$ . Then  $M' := A'f(M)$  is a finitely generated  $A'$ -submodule of  $K'$  for the action of  $A'$  through  $\varphi'$ , and so  $N' := S'M'$  is a finitely generated  $B$ -submodule for the action of  $B$  through  $\psi'$ . The modules  $M'$  and  $N'$  may have torsion, but since they are finitely generated, their torsion is annihilated by some non-zero element  $a' \in A'$ . Replacing the isogeny  $f$  by  $\varphi'_{a'} \circ f$  replaces  $M'$  by  $a'M'$  and  $N'$  by  $a'N'$ ; hence we may without loss of generality assume that  $M'$  and  $N'$  are torsion free.

Let  $\Delta_{\text{ad}, M'}^{\varphi'} \subset \Gamma_{\text{ad}, M'}^{\varphi'} \twoheadrightarrow \Gamma_{\text{ad}}^{\varphi'}$  denote the Galois groups as in (2.14) associated to  $(K', \varphi', M')$  in place of  $(K, \varphi, M)$ , and similarly for  $(K', \varphi', N')$ , respectively for  $(K', \psi', N')$ , and so on. Then Proposition 3.5 for the inclusion  $A' \supset B$  yields a natural commutative diagram with exact rows

$$(6.2) \quad \begin{array}{ccccccc} 0 & \longrightarrow & T_{\text{ad}}(\varphi') & \longrightarrow & T_{\text{ad}}(\varphi', N') & \longrightarrow & N' \otimes_{A'} A'_{\text{ad}} \longrightarrow 0 \\ & & \parallel & & \parallel & & \parallel \\ 0 & \longrightarrow & T_{\text{ad}}(\psi') & \longrightarrow & T_{\text{ad}}(\psi', N') & \longrightarrow & N' \otimes_B B_{\text{ad}} \longrightarrow 0. \end{array}$$

The action of  $S'$  on the lower row thus yields a natural action on the upper row. Recall from (2.19) that any  $S'$ -equivariant splitting induces an embedding

$$(6.3) \quad \Gamma_{\text{ad}, N'}^{\psi'} \hookrightarrow \Gamma_{\text{ad}}^{\psi'} \ltimes \text{Hom}_{S'}(N', T_{\text{ad}}(\psi')).$$

Since  $(B, K', \psi')$  is primitive, this embedding is open by Theorem 5.1. The isomorphisms from (3.6) thus yield an open embedding

$$(6.4) \quad \Gamma_{\text{ad}, N'}^{\varphi'} \hookrightarrow \Gamma_{\text{ad}}^{\varphi'} \ltimes \text{Hom}_{S'}(N', T_{\text{ad}}(\varphi')).$$

Since  $N' = S'M'$ , the Galois action on  $T_{\text{ad}}(\varphi', N')$  is completely determined by the action on  $T_{\text{ad}}(\varphi', M')$ ; in other words the restriction induces a natural isomorphism  $\Gamma_{\text{ad}, N'}^{\varphi'} \cong \Gamma_{\text{ad}, M'}^{\varphi'}$ . This together with Proposition 6.1 (a) yields an open embedding

$$(6.5) \quad \Gamma_{\text{ad}, M'}^{\varphi'} \hookrightarrow \Gamma_{\text{ad}}^{\varphi'} \ltimes \text{Hom}_{(S')}(M', T_{\text{ad}}(\varphi')).$$

The next natural step would be the passage from  $(K', \varphi', M')$  to  $(K', \varphi'|A, M')$ . However, this runs into the problem that  $S'$  does not necessarily act on  $T_{\text{ad}}(\varphi'|A)$ , because  $T_{\text{ad}}(\varphi'|A)$  is obtained from  $T_{\text{ad}}(\varphi') \cong \prod_{\mathfrak{p}' \neq \mathfrak{p}_0} T_{\mathfrak{p}'}(\varphi')$  by removing all factors with  $\mathfrak{p}'|\mathfrak{p}_0$ , which are not necessarily preserved by the non-commutative ring  $S'$ . Thus if  $\mathfrak{p}'_0$  is not the only prime above  $\mathfrak{p}_0$ , it would be ugly to precisely describe the image of  $\text{Hom}_{(S')}(M', T_{\text{ad}}(\varphi'))$  in  $\text{Hom}_A(M', T_{\text{ad}}(\varphi'|A))$  in general, though of course it can be done. We therefore restrict ourselves to two special cases, with the following results:

**Theorem 6.6** *Assume that  $A$  is the center of  $\text{End}_{K^{\text{sep}}}(\varphi)$ . With  $A' := A$  let  $(K', \varphi', f, B)$  be as in Proposition 4.3 (c), and set  $S := \text{End}_{K^{\text{sep}}}(\varphi|B)$ . Let  $M$  be a finitely generated torsion free  $A$ -submodule of  $K$ . Then  $\Delta_{\text{ad}, M}$  is commensurable with the subgroup  $\text{Hom}_{(S)}(M, T_{\text{ad}}(\varphi))$  of  $\text{Hom}_A(M, T_{\text{ad}}(\varphi))$ , and  $\Gamma_{\text{ad}, M}$  is commensurable with  $\Gamma_{\text{ad}} \ltimes \text{Hom}_{(S)}(M, T_{\text{ad}}(\varphi))$ .*

**Proof.** With the above notation  $f$  induces an isomorphism  $M \xrightarrow{\sim} M'$  and an embedding of finite index  $T_{\text{ad}}(\varphi) \hookrightarrow T_{\text{ad}}(\varphi')$ . It also induces an isomorphism  $S \otimes_B \text{Quot}(B) \cong S' \otimes_B \text{Quot}(B)$  under which the intersection of  $S$  and  $S'$  has finite index in both. Since  $T_{\text{ad}}(\varphi)$  and  $T_{\text{ad}}(\varphi')$  are torsion free  $A_{\text{ad}}$ -modules, this implies the equalities and an inclusion of finite index in the diagram

$$\begin{array}{ccc} \text{Hom}_{(S)}(M, T_{\text{ad}}(\varphi)) & = & \text{Hom}_{(S \cap S')}(M, T_{\text{ad}}(\varphi)) \\ & & \downarrow f \circ (\quad) \circ f^{-1} \\ \text{Hom}_{(S')}(M', T_{\text{ad}}(\varphi')) & = & \text{Hom}_{(S \cap S')}(M', T_{\text{ad}}(\varphi')). \end{array}$$

With (3.2) and the open embedding (6.5) this shows that the image of  $\text{Gal}(K^{\text{sep}}/K')$  associated to  $(K', \varphi, M)$  is an open subgroup of  $\Gamma_{\text{ad}} \ltimes \text{Hom}_{(S)}(M, T_{\text{ad}}(\varphi))$ . This implies the assertion about  $\Gamma_{\text{ad}, M}$ , from which the assertion about  $\Delta_{\text{ad}, M}$  directly follows.  $\square$

In the other special case we drop all assumptions on endomorphisms, but instead assume something about  $M$ :

**Theorem 6.7** *Let  $(A', K', \varphi', f, B)$  be as in Proposition 4.3 (c), and set  $S := \text{End}_{K^{\text{sep}}}(\varphi'|B)$ . Let  $M$  be a finitely generated torsion free  $A$ -submodule of  $K$  such that the natural map*

$$S \otimes_A M \rightarrow K^{\text{sep}}, \quad \sum_i s_i \otimes m_i \mapsto \sum_i s_i f(m_i)$$

*is injective. Then  $\Delta_{\text{ad}, M}$  is an open subgroup of  $\text{Hom}_A(M, T_{\text{ad}}(\varphi))$ , and  $\Gamma_{\text{ad}, M}$  is an open subgroup of  $\Gamma_{\text{ad}} \ltimes \text{Hom}_A(M, T_{\text{ad}}(\varphi))$ .*

**Proof.** With the above notation the assumption implies that the natural map

$$S' \otimes_A f(M) \rightarrow S' f(M) = N', \quad \sum_i s'_i \otimes m'_i \mapsto \sum_i s'_i m'_i$$

is injective and therefore an isomorphism. Thus by Proposition 6.1 the restriction induces a natural isomorphism

$$\text{Hom}_{S'}(N', T_{\text{ad}}(\varphi')) \xrightarrow{\sim} \text{Hom}_A(f(M), T_{\text{ad}}(\varphi')).$$

The openness of the embedding (6.4), together with the surjectivity in (3.6) for the inclusion  $A \subset A'$ , thus implies the openness of the embedding

$$\Gamma_{\text{ad}, f(M)}^{\varphi'|A} \hookrightarrow \Gamma_{\text{ad}}^{\varphi'|A} \ltimes \text{Hom}_A(f(M), T_{\text{ad}}(\varphi'|A)).$$

With (3.2) it follows that

$$\Gamma_{\text{ad}, M} \hookrightarrow \Gamma_{\text{ad}} \ltimes \text{Hom}_A(M, T_{\text{ad}}(\varphi))$$

is an open embedding. This proves the assertion about  $\Gamma_{\text{ad}, M}$ , from which the assertion about  $\Delta_{\text{ad}, M}$  directly follows.  $\square$

## References

- [1] Chi, W.-Ch., Li, A.: Kummer theory of division points over Drinfeld modules of rank one. *J. Pure Appl. Algebra* **156**, no.2–3 (2001) 171–185.
- [2] Devic, A., Pink, R.: Adelic openness for Drinfeld modules in special characteristic. Preprint (January 2012), 60 p. [arxiv:1103.3398v3 \[math.NT\]](https://arxiv.org/abs/1103.3398v3), to appear in *J. Number Theory*
- [3] Häberli, S.: *Kummer Theory of Drinfeld Modules*. Master's thesis ETH Zürich, May 2011, 39 p., [www.math.ethz.ch/~pink/Theeses/2011-Master-Simon-Haeberli.pdf](http://www.math.ethz.ch/~pink/Theeses/2011-Master-Simon-Haeberli.pdf)
- [4] Knebusch, M.: Elementarteilertheorie über Maximalordnungen. *J. Reine Angew. Math.* **226** (1967) 175–183.
- [5] Li, A.: A note on Kummer theory of division points over singular Drinfeld modules. *Bull. Austral. Math. Soc.* **64**, no.1 (2001) 15–20.
- [6] Pink, R.: The Galois representations associated to a Drinfeld module in special characteristic. II. Openness *J. Number Theory* **116**, no. 2 (2006) 348–372.
- [7] Pink, R., Rütsche, E.: Adelic openness for Drinfeld modules in generic characteristic. *J. Number Theory* **129** no.4 (2009) 882–907.
- [8] Poonen, B.: Local height functions and the Mordell-Weil theorem for Drinfeld modules. *Compositio Math.* **97**, no.3 (1995) 349–368.
- [9] Reiner, I.: *Maximal orders*. London Mathematical Society Monographs, New Series 28. Oxford University Press (2003).
- [10] Ribet, K. A.: Kummer theory on extensions of abelian varieties by tori. *Duke Math. J.* **46**, no.4 (1979) 745–761.
- [11] Wang, J. T.-Y.: The Mordell-Weil theorems for Drinfeld modules over finitely generated function fields. *Manuscripta Math.* **106**, no.3 (2001) 305–314.