

Avoiding Small Subgraphs in Achlioptas Processes

Michael Krivelevich,^{1,*} Po-Shen Loh,^{2,†} Benny Sudakov^{3,‡}

¹School of Mathematical Sciences, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv 69978, Israel;

e-mail: krivelev@post.tau.ac.il

²Department of Mathematics, Princeton University, Princeton, New Jersey 08544;

e-mail: ploh@math.princeton.edu

³Department of Mathematics, UCLA, Los Angeles, California 90095;

e-mail: bsudakov@math.ucla.edu

Received 2 August 2007; accepted 13 October 2008

Published online 10 November 2008 in Wiley InterScience (www.interscience.wiley.com).

DOI 10.1002/rsa.20254

ABSTRACT: For a fixed integer r , consider the following random process. At each round, one is presented with r random edges from the edge set of the complete graph on n vertices, and is asked to choose one of them. The selected edges are collected into a graph, which thus grows at the rate of one edge per round. This is a natural generalization of what is known in the literature as an Achlioptas process (the original version has $r = 2$), which has been studied by many researchers, mainly in the context of delaying or accelerating the appearance of the giant component.

In this article, we investigate the small subgraph problem for Achlioptas processes. That is, given a fixed graph H , we study whether there is an online algorithm that substantially delays or accelerates a typical appearance of H , compared to its threshold of appearance in the random graph $G(n, M)$. It is easy to see that one cannot accelerate the appearance of any fixed graph by more than the constant factor r , so we concentrate on the task of avoiding H . We determine thresholds for the avoidance of all cycles C_r , cliques K_r , and complete bipartite graphs $K_{t,t}$, in every Achlioptas process with parameter $r \geq 2$. © 2008 Wiley Periodicals, Inc. *Random Struct. Alg.*, 34, 165–195, 2009

Keywords: Achlioptas process; small subgraph; threshold; random graph

Correspondence to: M. Krivelevich

*Supported in part by USA-Israel BSF Grants (2002-133, 2006-322); Israel Science Foundation (526/05).

†Supported in part by a Fannie and John Hertz Foundation Graduate Fellowship, an NSF Graduate Research Fellowship, and a Princeton Centennial Fellowship.

‡Supported in part by NSF CAREER Award (DMS-0812005), and USA-Israel BSF Grant (2004-029).

© 2008 Wiley Periodicals, Inc.

1. INTRODUCTION

The standard Erdős-Rényi random graph model $G(n, M)$ can be described as follows. Start with the empty graph on n vertices, and perform M rounds, adding one random edge to the graph at each round. For any monotone increasing graph property (such as containment of K_4 as a subgraph, say), it is natural to ask whether there is some value of M at which the probability of $G(n, M)$ satisfying the property changes rapidly from nearly 0 to nearly 1. More precisely, a function $M^*(n)$ is said to be a threshold for a property \mathcal{P} if for any $M(n) \ll M^*(n)$, the random graph $G(n, M)$ does not satisfy \mathcal{P} **whp**, but for any $M(n) \gg M^*(n)$, the random graph $G(n, M)$ satisfies \mathcal{P} **whp**. Here, **whp** stands for “with high probability,” that is, with probability tending to 1 as $n \rightarrow \infty$, and $f(n) \ll g(n)$ means that $f(n)/g(n) \rightarrow 0$ as $n \rightarrow \infty$. A classical result of Bollobás and Thomason [10] implies that every monotone graph property has a threshold, and much work has been done to determine thresholds for various properties.

Recently, there was much interest in the following natural variant of the classical model. We still begin with the empty graph and perform a series of rounds, but at each round, one is now presented with two independent and uniformly random edges, and is asked to choose one of them to add to the graph. This is known in the literature as an Achlioptas process, after Dimitris Achlioptas, who asked the question of whether there was an online algorithm which could, with high probability, substantially delay the appearance of the giant component (a connected component with $\Omega(n)$ vertices).

The trivial algorithm, which arbitrarily chooses the first edge in each offered pair, essentially produces the random graph $G(n, M)$ after M rounds, so $G(n, M)$ serves as the benchmark against which comparisons are made. A classical result of Erdős and Rényi [11] states that if $M = cn$ for any absolute constant $c > 1/2$, then the random graph $G(n, M)$ contains a giant component **whp**. For the Achlioptas process, Bohman and Frieze [4] presented an algorithm which could run for $0.535n$ rounds, while keeping the size of the largest component only poly-logarithmic in n **whp**. Since then, much work has been done [2, 3, 5–7, 13, 16]. The current best result for this problem is due to Spencer and Wormald [16], who exhibit an algorithm that can run for $0.829n$ rounds while keeping all component sizes bounded by $O(\log n)$ **whp**. In the opposite direction, Bohman, Frieze, and Wormald [5] have shown that no algorithm can succeed **whp** past $0.964n$ rounds. Several variants have also been studied, such as the offline version, a two-player version, and the question of embracing (accelerating the appearance of) the giant component.

While the main focus of the research mentioned above was the giant component, it is natural to study other graph properties in the context of Achlioptas processes. In this article, we study the problem which in the literature is referred to as the small subgraph problem. This was one of the main problems studied in the seminal paper of Erdős and Rényi [11] from 1960, which was the starting point of the theory of random graphs. The original problem, stated for the random graph model $G(n, M)$, was as follows: given a fixed graph H (a triangle or K_4 , say), find the smallest value of M such that the random graph $G(n, M)$ contains H as a (not necessarily induced) subgraph **whp**. The subgraph is called “small” because its size is fixed while n tends to infinity.

It turns out that in this problem, the relevant parameter is the maximum edge density $m(G) = \max\{e(H)/v(H) : H \text{ is a subgraph of } G\}$. In their original paper, Erdős and Rényi found thresholds for all balanced graphs, which are the graphs whose edge density $e(H)/v(H)$ equals the maximum edge density $m(H)$. It was not until 20 years later that Bollobás [9] solved the problem for all graphs, proving that for any H with $m(H) \geq 1$,

the threshold for H appearing in $G(n, M)$ is $M^* = n^{2-\frac{1}{m(H)}}$. For further reading about the small subgraph problem in $G(n, M)$, we direct the interested reader to the monographs by Bollobás [8] and by Janson, Łuczak, and Ruciński [14], each of which contains an entire section discussing this problem.

In this article, we consider the small subgraph problem in the context of Achlioptas processes, and investigate whether one can substantially affect thresholds by introducing this power of choice. Actually, we study a natural generalization of the process, which we call an Achlioptas process with parameter r . In this process, r edges of K_n are presented at each round, and one of them is selected. We will always consider r to be fixed as n tends to infinity (note that $r = 2$ corresponds to the original Achlioptas process).

Let us now state our model precisely. At the i -th round, one is presented with r independent random edges, each distributed uniformly over all $\binom{n}{2} - (i - 1)$ remaining edges that have not yet been chosen for the graph. Note that this eliminates the possibility of choosing the same edge twice, so our final graph is simple. However, we do allow the possibility that edges may be offered more than once, which simplifies our arguments. One may consider models in which all sampling is with replacement (which may create multigraphs), or in which every edge is offered at most once, but our results in this article will still carry over because we always run the process for $o(n^2)$ rounds.

Note that the graph after the k -th round of the Achlioptas process with parameter r is a subgraph of the random graph with rk edges. So, the question of accelerating the appearance of a fixed graph is immediately resolved in the negative. Clearly, the threshold cannot move forward by more than a (constant) factor of r .

So, in this article we concentrate on the avoidance problem. We may pose it as a single player game in which the player loses when he creates a (not necessarily induced) subgraph isomorphic to a certain fixed graph H . The player's objective is to postpone losing for as long as possible. We say that a function $m^*(n)$ is a threshold for avoiding H if: (i) given any function $m(n) \ll m^*(n)$, there exists an online strategy by which the player survives through m rounds **whp**, and (ii) given any function $m(n) \gg m^*(n)$, the player loses by the end of m rounds **whp**, regardless of the choice of such a strategy.

Note, however, that it is not obvious that thresholds necessarily exist. Furthermore, unlike the situation in the small subgraph problem, there are no simple first-moment calculations that suggest what the thresholds should be. As it turned out, a substantial part of the difficulty in obtaining our results was in conjecturing the correct thresholds. We were able to solve the problem for all cycles C_t , cliques K_t , and complete bipartite graphs $K_{t,t}$. Let us now state our main result:

Theorem 1.1.

- (i) For $t \geq 3$, the threshold for avoiding C_t in the Achlioptas process with parameter $r \geq 2$ is $n^{2-\frac{(r-2)t+2}{(r-1)t+1}}$.
- (ii) For $t \geq 4$, the threshold for avoiding K_t in the Achlioptas process with parameter $r \geq 2$ is $n^{2-\theta}$, where θ is defined as follows:

$$s = \lfloor \log_r[(r - 1)t + 1] \rfloor, \quad \theta = \frac{r^s(t - 2) + 2}{r^s \left[\binom{t}{2} - s \right] + \frac{r^s - 1}{r - 1}}.$$

- (iii) For $t \geq 3$, the threshold for avoiding $K_{t,t}$ in the Achlioptas process with parameter $r \geq 2$ is $n^{2-\theta}$, where θ is defined as follows:

$$s = \lfloor \log_r[(r - 1)t + 1] \rfloor, \quad \theta = \frac{r^s(2t - 2) + 2}{r^s(t^2 - s) + \frac{r^s - 1}{r - 1}}.$$

Remark. In all of these cases, we provide deterministic online algorithms that achieve the thresholds **whp**, but show that even randomized algorithms cannot survive beyond the thresholds.

The rest of this article is organized as follows. In the next section, we present some tools from extremal combinatorics and the theory of random graphs, which we will use in our proofs. Then, we present the proof of our theorem, which is divided into several sections. We begin in Section 3 with the case of avoiding K_4 when $r = 2$, which turns out to be the first nontrivial case. We treat this case in detail, because our argument there is the prototype for the general argument that we later use to prove thresholds for K_t , $K_{t,t}$, and C_t .

We extend the argument to almost all other K_t and r in Section 4. The proof requires many inequalities whose somewhat tedious verifications would interfere with the exposition, so their precise statements are recorded in the appendix.¹ This also makes it easier to distill the abstract argument, which we present in Section 5. Next, we apply the abstraction to prove thresholds for avoiding C_t in Section 6 and $K_{t,t}$ in Section 7. We treat the last remaining case of avoiding K_4 in the Achlioptas process with parameter 3 in Section 8. The final section contains some concluding remarks and open problems.

2. PRELIMINARIES

2.1. Notation and Terminology

Throughout our paper, we will omit floor and ceiling signs whenever they are not essential, to improve clarity of presentation. The following (standard) asymptotic notation will be utilized extensively. For two functions $f(n)$ and $g(n)$, we write $f(n) = o(g(n))$ or $g(n) = \omega(f(n))$ if $\lim_{n \rightarrow \infty} f(n)/g(n) = 0$, and $f(n) = O(g(n))$ or $g(n) = \Omega(f(n))$ if there exists a constant M such that $|f(n)| \leq M|g(n)|$ for all sufficiently large n . We also write $f(n) = \Theta(g(n))$ if both $f(n) = O(g(n))$ and $f(n) = \Omega(g(n))$ are satisfied.

Let us introduce the following abbreviations for some phrases that we will use many times in our proof. As mentioned in the introduction, **whp** will stand for “with high probability,” i.e., with probability $1 - o(1)$. It is also convenient for us to introduce the abbreviation **wep**, which stands for “with exponential probability,” i.e., with probability $1 - o(e^{-n^c})$ for some $c > 0$. We will say that a function f is a positive power of n if $f = \Omega(n^c)$ for some $c > 0$. Analogously, we will say that a function f is a negative power of n if $f = O(n^{-c})$ for some $c > 0$.

Next, let us discuss the graph-specific terms that we will use. We often need to consider the graphs at intermediate stages of the Achlioptas process, so G_i will always denote the graph after the i -th round. Our main interest in G_i will be to count copies of subgraphs. Here, we define a copy of a graph H in another graph G to be an injective map from $V(H)$ to $V(G)$ that preserves the edges of H . Note that copies are not necessarily induced subgraphs, and are labeled, i.e., we do not take automorphisms into account when computing the number of copies of H in a graph.

The player’s objective in the Achlioptas process is to avoid creating a copy of a certain fixed graph H , but our analysis needs to consider subgraphs of H as well. It is therefore

¹The proofs of these inequalities are rather technical and not so interesting, so they only appear in the unabridged version of this paper which is on the arXiv at <http://arxiv.org/abs/0708.0443>.

convenient to introduce the notation $H \setminus ke$ to represent any graph which can be obtained by deleting any k edges from H . (When $k = 1$, we will simply write $H \setminus e$.) This enables us to concisely refer to all graphs of the form $H \setminus ke$ in the aggregate. For example, the phrase “the number of copies of $H \setminus ke$ ” should be understood to be the total number of copies of all graphs of the form $H \setminus ke$.

We keep track of the number of copies of these subgraphs by studying how counts are affected by the addition of an edge at a pair of vertices. This motivates the following definition. Let G and H be graphs, let k be an integer, and let a, b be a pair of distinct vertices of G . Let G^+ be the graph obtained from G by adding the edge between a and b if it is not yet present, and let G^- be the graph obtained by deleting that edge if it was present. Note that G is equal to either G^+ or G^- . Then, we say that the pair $\{a, b\}$ completes t copies of $H \setminus ke$ if t is the difference between the number of copies of $H \setminus ke$ in G^+ and the number in G^- .

Sometimes, we need to be specific about which graphs of the form $H \setminus (k + 1)e$ are completed into graphs of the form $H \setminus ke$. Let H_1 and H_2 be graphs on the same vertex set U , with $E(H_1) \subset E(H_2)$, but differing only in exactly one edge. Let $\{u, v\} \subset U$ be the endpoints of that edge. Let G be another graph, and let a, b be a pair of distinct vertices of G . Then, we say that the pair $\{a, b\}$ extends t copies of H_1 into H_2 if t is the number of injective graph homomorphisms $\phi : H_1 \rightarrow G$ that map $\{u, v\}$ to $\{a, b\}$. Note that this definition is insensitive to the presence of an edge between a and b .

2.2. Extremal Combinatorics

In this section, we present two extremal results, which are used in the proofs of the upper bounds in our thresholds (i.e., that no strategy can survive for too many rounds). The following lower bound on the number of paths in a graph was obtained in [12] using a matrix inequality of Blackley and Roy.

Lemma 2.1. *Every graph with n vertices and average degree d contains at least $(1 + o(1))nd^{t-1}$ copies of the t -vertex path P_t . Here, we consider t to be fixed, while d and n tend to infinity.*

Next, we record the following well-known extremal result, which lower bounds the number of copies of the complete bipartite graph $K_{s,t}$ that can appear in any graph with a fixed number of edges. The classical proof (via two applications of convexity) is based on the ideas used by Kővári, Sós, and Turán [15] to bound the Turán number $\text{ex}(n, K_{s,t})$.

Lemma 2.2. *For fixed positive integers $s \leq t$, and any function $p \gg n^{-1/s}$, every graph with n vertices and $\binom{n}{2}p$ edges contains at least $(1 + o(1))n^{s+t}p^{st}$ copies of the complete bipartite graph $K_{s,t}$.*

2.3. Random Graphs

We begin by recalling the Chernoff bound for exponential concentration of a binomial random variable. We use the formulation from [1].

Theorem 2.3. *For any $\epsilon > 0$, there exists $c_\epsilon > 0$ such that the following holds. Let X be any binomial random variable, and let μ be its expectation. Then $\mathbb{P}[|X - \mu| > \epsilon\mu] < 2e^{-c_\epsilon\mu}$.*

Using the Chernoff bound and a standard coupling argument, we prove a result that allows us to relate G_m (the graph after the m -th round of the Achlioptas process) to the more familiar random graph $G(n, p)$.

Lemma 2.4. *Suppose that $n \ll m \ll n^2$. Then we may couple the Achlioptas process with $G(n, p = 4rm/n^2)$ in such a way that **wep**, G_m is a subgraph of $G(n, p)$.*

Proof. In the Achlioptas process, r random edges are presented at each round, independently and uniformly distributed over all potential edges that have not yet been picked for the graph. So, we may couple the first m rounds of the process with the edge-uniform random graph $G(n, rm)$ in such a way that if we consider the graph G_m^+ obtained by taking every edge that was offered (instead of choosing only one per round), G_m^+ is always a subgraph of $G(n, rm)$. Yet G_m is always a subgraph of G_m^+ , so it remains to relate $G(n, rm)$ with $G(n, p = 4rm/n^2)$. This final part is standard and proceeds via coupling with the random graph process; under this coupling, $G(n, rm) \subset G(n, p)$ as long as $\text{Bin}\binom{n}{2}, p \geq rm$, and the Chernoff bound shows that this event occurs **wep**. ■

Our analysis revolves around counting copies of fixed subgraphs in G_m . The previous lemma allows us to apply results from the theory of $G(n, p)$ to assist us in this pursuit. We now record several such theorems, translated in terms of G_m . The following definition is crucial for counting subgraphs in $G(n, p)$.

Definition 2.5. *A graph H is **balanced** if for any subgraph $H' \subset H$, $\frac{e(H')}{v(H')} \leq \frac{e(H)}{v(H)}$.*

Theorem 2.6. *Let H be a fixed balanced graph with v vertices and e edges. Suppose that $n \ll m \ll n^2$, and let $p = 2m/n^2$. Also suppose that $n^v p^e$ is a positive power of n . Then the number of copies of H in G_m is $O(n^v p^e)$ **wep**.*

Proof. By Lemma 2.4, it suffices to count copies of H in $G(n, 2rp)$. The expected number of copies is $(1 + o(1))n^v (2rp)^e = \Theta(n^v p^e)$, which is a positive power of n by assumption. This allows us to apply Corollary 6.3 of [17], which uses Kim-Vu polynomial concentration to prove the following result: for any balanced graph H such that the expected number of copies of H in the random graph is $\mu \gg \log n$, the probability that the actual number of copies exceeds 2μ is $e^{-\Omega(\mu)}$. In our case, μ is a positive power of n , so this implies that **wep**, the number of copies is $O(n^v p^e)$, as desired. ■

The previous result provides a very precise count of the number of copies of a fixed graph in the random graph $G(n, p)$. However, the point of the Achlioptas process was to deviate from $G(n, p)$ by introducing the power of choice. So, our analysis will have to take the potential of choice into account. We keep track of the number of copies of subgraphs by studying how counts are affected by the addition of an edge at a pair of vertices; this motivated the notions of a pair completing t copies of $H \setminus ke$ and of the pair extending t copies of H_1 into H_2 , which we defined at the end of Section 2.1.

This is essentially the problem of counting extensions, which has also been well-studied in $G(n, p)$. We refer the interested reader to Chapter 10 of [1]. As in the case of counting subgraphs in $G(n, p)$, a suitable definition of balanced-ness is required to count extensions.

Definition 2.7.

- (i) *Let H_1 and H_2 be graphs on the same vertex set U , with $E(H_1) \subset E(H_2)$, but differing only on the edge joining the vertices $u, v \in U$. We say that the pair (H_1, H_2) is a*

- balanced extension pair* if for every proper subset $U' \subset U$ that still contains $\{u, v\}$, the induced subgraph $H' = H_1[U']$ has the property that $\frac{e(H')}{v(H')-2} \leq \frac{e(H_1)}{v(H_1)-2}$.
- (ii) $H \setminus ke$ has the **balanced extension property** if every pair (H_1, H_2) with $V(H_1) = V(H_2) = V(H)$, $E(H_1) \subset E(H_2) \subset E(H)$, $e(H_1) = e(H) - k$, and $e(H_2) = e(H) - k + 1$, is a balanced extension pair.

Theorem 2.8. *Suppose that $n \ll m \ll n^2$, and let $p = 2m/n^2$. Let (H_1, H_2) be a balanced extension pair, and let v and e be the numbers of vertices and edges in H_1 , respectively. Finally, let j be an arbitrary integer constant.*

- (i) *Suppose that $n^{v-2}p^e$ is a positive power of n . Then **wep**, every pair of distinct vertices $\{a, b\}$ of G_{jm} extends $O(n^{v-2}p^e)$ copies of H_1 into H_2 .*
- (ii) *Suppose that $n^{v-2}p^e$ is a negative power of n . Then, for any constant $\gamma > 0$, there exists a constant C such that with probability $1 - o(n^{-\gamma})$, every pair of distinct vertices $\{a, b\}$ of G_{jm} extends at most C copies of H_1 into H_2 .*

Proof. By Lemma 2.4, it suffices to consider $G(n, 2rjp)$ instead of G_{jm} in both parts of the theorem. For part (i), the expected number of extensions at a pair in $G(n, 2rjp)$ is $(1 + o(1))n^{v-2}(2rjp)^e = \Theta(n^{v-2}p^e)$, which is a positive power of n by assumption. This allows us to apply Corollary 6.7 of [17], which uses Kim-Vu polynomial concentration to prove the following result: for any balanced extension pair (H_1, H_2) such that the expected number μ of copies of H_1 that a fixed edge extends into H_2 in the random graph is a positive power of n , the probability that the actual number of extensions exceeds 2μ is $e^{-\Omega(\mu)}$. In our case, μ is a positive power of n , so even after taking a union bound over all $O(n^2)$ pairs of vertices, this implies that **wep**, every pair of vertices extends $O(n^{v-2}p^e)$ copies of H_1 into H_2 . This establishes (i).

For part (ii), let us bound the probability that $\{a, b\}$ extends C copies of H_1 into H_2 . Recall that H_1 and H_2 shared the same vertex set U , and differed only on the edge joining $u, v \in U$. Consider any graph F which is formed by the superposition of C distinct copies of H_1 , all with $\{u, v\}$ mapping to the same pair of vertices $\{u', v'\} \in V(F)$. Let $v' = v(F)$ and $e' = e(F)$.

The probability that $\{a, b\}$ has an extension to F (an injective map from $V(F)$ sending $\{u', v'\} \mapsto \{a, b\}$) in $G(n, 2rjp)$ is at most $n^{v'-2}(2rjp)^{e'} = O((np^{e'/(v'-2)})^{v'-2})$. An easy and standard induction, using the fact that (H_1, H_2) is a balanced extension pair, implies that $\frac{e'}{v'-2} \geq \frac{e}{v-2}$. Hence this probability is at most $O((np^{e/(v-2)})^{v'-2}) = O((n^{v-2}p^e)^{\frac{v'-2}{v-2}})$.

We assumed that $n^{v-2}p^e$ was a negative power of n . Also, since the C copies of H_1 in F are distinct, one can trivially bound $C \leq (v' - 2)^{v-2} \Rightarrow v' - 2 \geq C^{\frac{1}{v-2}}$. So, for a sufficiently large constant C , the probability that $\{a, b\}$ has an extension to F is $o(n^{-\gamma-2})$. Taking a union bound over all $O(n^2)$ pairs of vertices, we see that the probability that there exists any pair of vertices with an extension to F is $o(n^{-\gamma})$. Since C is a constant, the number of non-isomorphic ways to form F (a superposition of C distinct copies of H_1 , overlapping on one particular edge) is still a constant. Taking another union bound over all such F , we complete the proof. ■

Corollary 2.9. *Suppose that $n \ll m \ll n^2$, and let $p = 2m/n^2$. Let $H \setminus ke$ have the balanced extension property, and let v and e be the numbers of vertices and edges in $H \setminus ke$. Suppose that $n^{v-2}p^e$ is a negative power of n . Let us consider G_{jm} , where j is an arbitrary*

integer constant. Then, for any constant $\gamma > 0$, there exists a constant C such that with probability $1 - o(n^{-\gamma})$, every pair of distinct vertices $\{a, b\}$ of G_m completes at most C copies of $H \setminus (k - 1)e$.

Proof. Fix a pair $\{a, b\}$. When counting the number of copies of $H \setminus (k - 1)e$ completed by that pair, each copy arises from an extension pair (H_1, H_2) and an extension of H_1 to H_2 at the pair. In fact, this correspondence is bijective. The balanced extension property guarantees that all such pairs are balanced. Since H is a fixed graph, only a constant number of non-isomorphic pairs (H_1, H_2) can arise in this way, so repeated application of Theorem 2.8(ii) completes the proof. ■

3. WARM-UP

The purpose of this section is to illustrate on a concrete example the main ideas and techniques that we will use in our proofs. We investigate the first nontrivial case, which is the problem of avoiding K_4 in the Achlioptas process with parameter 2. This turns out to be the model for the general case.

Theorem. *The threshold for avoiding K_4 in the Achlioptas process with parameter 2 is $n^{28/19}$.*

Proof of lower bound. We need to specify a strategy, and prove that it avoids K_4 for many rounds. At any intermediate stage in the process, consider a pair of points to be 2-dangerous if the addition of an edge between them will create a copy of K_4 . Otherwise, if the addition of the edge will create a copy of $K_4 \setminus e$, call the pair 1-dangerous. Every other pair is considered to be 0-dangerous (not dangerous). The strategy is then to make an arbitrary choice among the incoming edges that are minimally dangerous.

Let m be a function of n that satisfies $m \ll n^{28/19}$. It suffices to show that for any such m , this strategy succeeds **whp**. We also may assume without loss of generality that $m \gg n^{28/19} / \log n$. The precise form of the lower bound on m is not essential; it simplifies the argument by disposing of uninteresting pathological cases when m is too small. As it is easier to work with $G(n, p)$, we will make all of our computations with respect to p , which we define to be $2m/n^2$. Note that $n^{-10/19} / \log n \ll p \ll n^{-10/19}$. The following three claims analyze the performance of our strategy.

- (i) With probability $1 - o(n^{-4})$, G_m has $O(n^4 p^4)$ copies of $K_4 \setminus 2e$ and every pair of vertices completes $O(1)$ copies of $K_4 \setminus e$.
- (ii) With probability $1 - o(n^{-2})$, G_m has $O(n^6 p^9)$ copies of $K_4 \setminus e$.
- (iii) The probability of failure in m rounds is $o(1)$.

For (i), it is easy to verify that $K_4 \setminus 2e$ is a balanced graph, no matter which two edges are deleted. Then the number of copies of $K_4 \setminus 2e$ is roughly what it should be in the random graph $G(n, p)$ —this is made precise by Theorem 2.6, which bounds the number of copies of $K_4 \setminus 2e$ in G_m by $O(n^4 p^4)$ **wep** since $n^4 p^4$ is a positive power of n . It is also easy to verify that $K_4 \setminus 2e$ has the balanced extension property, so since $n^2 p^4$ is a negative power of n , Corollary 2.9 shows that there is some constant C such that with probability $1 - o(n^{-4})$, every pair of vertices in G_m completes at most C copies of $K_4 \setminus e$. This proves (i).

For (ii), fix some $i < m$ and consider the $(i + 1)$ -st round. In this round, the strategy will create one or more copies of $K_4 \setminus e$ only if both incoming edges span pairs that are 1- or 2-dangerous. The number of such pairs is at most $O(1)$ times the number of copies of $K_4 \setminus 2e$. Since $G_i \subset G_m$, claim (i) shows that with probability $1 - o(n^{-4})$, G_i has $O(n^4 p^4)$ copies of $K_4 \setminus 2e$ and every pair of vertices completes $O(1)$ copies of $K_4 \setminus e$. Call this event A_i , and condition on it. Even after conditioning, the incoming edges at the $(i + 1)$ -st round are still independently and uniformly distributed over the $\Omega(n^2)$ unoccupied pairs of G_i , so the probability that we are forced to create a new copy of $K_4 \setminus e$ in this round is $O((\frac{n^4 p^4}{n^2})^2) = O(n^4 p^8)$. Furthermore, each time this occurs, we only create $O(1)$ new copies of $K_4 \setminus e$ because of our conditioning. Therefore, the number of new copies of $K_4 \setminus e$ in the $(i+1)$ -st round is stochastically dominated by $O(1)$ times the Bernoulli random variable with parameter $O(n^4 p^8)$. Letting i run through all m rounds, we see that with probability at least $1 - \sum \mathbb{P}[\neg A_i] \geq 1 - o(n^{-2})$, the number of copies of $K_4 \setminus e$ in G_m is $O(1) \cdot \text{Bin}[m, O(n^4 p^8)]$. Since $m = n^2 p/2$, the expectation of this binomial is a positive power of n , so the Chernoff bound implies that **wep**, it is $O(m \cdot n^4 p^8) = O(n^6 p^9)$. This proves (ii).

For (iii), fix some i and consider the probability that we lose in the $(i + 1)$ -st round. The strategy fails precisely when both of the incoming edges span pairs that are 2-dangerous (completing K_4), and the number of such pairs is at most $O(1)$ times the number of copies of $K_4 \setminus e$. Since $G_i \subset G_m$, claim (ii) shows that with probability $1 - o(n^{-2})$, G_i has $O(n^6 p^9)$ copies of $K_4 \setminus e$. Call this event B_i , and condition on it. Even after conditioning, the incoming edges are still independently and uniformly distributed over the $\Omega(n^2)$ unoccupied pairs of G_i , so the probability that both incoming edges are 2-dangerous is $O((\frac{n^6 p^9}{n^2})^2) = O(n^8 p^{18})$. Therefore, letting i run through all $m = n^2 p/2$ rounds, a union bound shows that the probability that we are forced to complete a copy of K_4 by the end of the m -th round is $\mathbb{P} \leq O(n^2 p \cdot n^8 p^{18}) + \sum \mathbb{P}[\neg B_i] = O(n^{10} p^{19}) + o(1) = o(1)$. ■

Proof of upper bound. Now suppose that $m \gg n^{28/19}$. It suffices to show that we will lose within the first $4m$ rounds **whp**. Again, we may assume without loss of generality that $m \ll n^{28/19} \log n$, and we will work in terms of $G(n, p)$ with $p = 2m/n^2$. Note that $n^{-10/19} \ll p \ll n^{-10/19} \log n$. Let us specify a sequence of graphs such that each graph is obtained from the previous one by adding a single edge: let $H_0 = P_4$ (4-vertex path), $H_1 = C_4$ (4-cycle), $H_2 = K_4 \setminus e$, and $H_3 = K_4$. It is easy to verify that the corresponding pairs (H_0, H_1) , (H_1, H_2) , and (H_2, H_3) are all balanced extension pairs. Our result follows from the following four claims:

- (i) G_m always contains $\Omega(n^4 p^3)$ copies of H_0 . Also, **wep**, every pair of vertices in G_{2m} extends $O(n^2 p^3)$ copies of H_0 into H_1 .
- (ii) G_{2m} contains $\Omega(n^4 p^4)$ copies of H_1 **whp**, and with probability $1 - o(n^{-2})$, every pair of vertices in G_{3m} extends $O(1)$ copies of H_1 into H_2 .
- (iii) G_{3m} contains $\Omega(n^6 p^9)$ copies of H_2 **whp**, and with probability $1 - o(n^{-2})$, every pair of vertices in G_{4m} extends $O(1)$ copies of H_2 into H_3 .
- (iv) The probability of survival through $4m$ rounds is $o(1)$.

Proof of (i). Since the average degree in G_m is precisely $2m/n = np \gg 1$, from Lemma 2.1 we conclude that the number of 4-vertex paths is $\Omega(np^3)$. The second part of this claim follows from Theorem 2.8(i) since (H_0, H_1) is a balanced extension pair and $n^2 p^3$ is a positive power of n . ■

Proof of (ii). The second part of (ii) follows from Theorem 2.8(ii) since (H_1, H_2) is balanced and n^2p^4 is a negative power of n . To prove the first part of (ii), consider the $(i + 1)$ -st round, where $m \leq i < 2m$. Regardless of the choice of strategy, if both incoming edges span pairs that extend $\Omega(n^2p^3)$ copies of H_0 into H_1 , we will be forced to create $\Omega(n^2p^3)$ new copies of H_1 .

By (i), the total number of copies of H_0 in $G_i \supset G_m$ is $\Omega(n^4p^3)$. For a pair of vertices $\{a, b\}$, let $n_{a,b}$ be the number of copies of H_0 that $\{a, b\}$ extends to H_1 . Recall that this definition does not depend on the presence of an edge between a and b . Since $G_i \subset G_{2m}$, claim (i) shows that **wep**, in G_i every $n_{a,b} = O(n^2p^3)$. Call this event A_i , and condition on it.

Let us estimate the average value of $n_{a,b}$ over all pairs. Since H_0 differs from H_1 at exactly one edge, each copy of H_0 has a pair at which it contributes $+1$ to the sum $\sum n_{a,b}$. Therefore, averaging over all $\binom{n}{2}$ pairs of vertices, we obtain that the average number of copies of H_0 that are extended to H_1 at a pair is $\Omega(n^2p^3)$. On the other hand, every pair of vertices in G_i extends $O(n^2p^3)$ copies of H_0 into H_1 . Therefore, at least a constant fraction γ (where $\gamma = \Omega(1)$ can be chosen to be the same for all i) of all $\binom{n}{2}$ pairs have the property of extending $\Omega(n^2p^3)$ copies of H_0 into H_1 . Let P be the set of all such pairs. Regardless of the choice of strategy, if both incoming edges span pairs in P , we will be forced to create $\Omega(n^2p^3)$ copies of H_1 . Since $i = o(n^2) = o(|P|)$ and incoming edges are uniformly distributed over the $\binom{n}{2} - i = (1 - o(1))\binom{n}{2}$ unoccupied pairs, we conclude that the probability that both incoming edges span pairs in P is $q \geq (1 + o(1))\gamma^2 = \Omega(1)$.

Let i run from m to $2m$. Then, up to an error probability of at most $\sum \mathbb{P}[\neg A_i] = o(1)$, the number of copies of H_1 in G_{2m} is at least $\text{Bin}(m, q) \cdot \Omega(n^2p^3)$. By the Chernoff bound, the binomial factor exceeds $mq/2 = \Omega(n^2p)$ **wep**; thus, **whp** G_{2m} has $\Omega(n^2p \cdot n^2p^3) = \Omega(n^4p^4)$ copies of H_1 . ■

Proof of (iii). The second part of (iii) follows from Theorem 2.8(ii) since (H_2, H_3) is balanced and n^2p^5 is a negative power of n . For the first part of (iii), let us consider the $(i + 1)$ -st round, with $2m \leq i < 3m$. Regardless of the choice of strategy, if both incoming edges span pairs that extend copies of H_1 into H_2 , we will create a copy of H_2 . Let P be the set of all such pairs. We need a lower bound on $|P|$. Condition on the event B that G_{2m} contains $\Omega(n^4p^4)$ copies of H_1 , which occurs **whp** by (ii). Also by (ii), with probability $1 - o(n^{-2})$, every pair of vertices in G_i only extends $O(1)$ copies of H_1 into H_2 , since $G_i \subset G_{3m}$. Call this event C_i , and condition on it.

Note that every copy of H_1 contributes a pair to P which extends H_1 into H_2 , namely the pair at which it is missing an edge compared to H_2 . On the other hand, every such pair was only counted $O(1)$ times, since every pair in G_i extends $O(1)$ copies of H_1 into H_2 . This implies that $|P| = \Omega(n^4p^4)$. The incoming edges are uniformly distributed over all unoccupied pairs. If at least half of the pairs in P were occupied, then we would have $\Omega(n^4p^4) \gg n^6p^9$ copies of H_2 , which would already give the conclusion of (iii). Otherwise, the probability that both incoming edges span pairs in P (hence forcing the creation of a new copy of H_2) is $q \geq (1 + o(1))\left(\frac{|P|/2}{n^2/2}\right)^2 = \Omega\left(\left(\frac{n^4p^4}{n^2}\right)^2\right) = \Omega(n^4p^8)$.

Letting i run from $2m$ to $3m$, we see that with error probability at most $\mathbb{P}[\neg B] + \sum \mathbb{P}[\neg C_i] = o(1)$, either we already obtained the conclusion of (iii), or the total number of copies of H_2 is at least $\text{Bin}(m, q)$. The expectation of this binomial is $(n^2p/2)q = \Omega(n^6p^9)$, which is a positive power of n . Hence, by the Chernoff bound, G_{3m} has $\Omega(n^6p^9)$ copies of H_2 **whp**. ■

Proof of (iv). Consider the $(i + 1)$ -st round, where $3m \leq i < 4m$. Regardless of the choice of strategy, if both incoming edges span pairs that complete copies of $H_3 = K_4$, we lose. We can lower bound the number of such pairs by $\Omega(n^6 p^9)$ by conditioning on the following events. Let D be the event that G_{3m} contains $\Omega(n^6 p^9)$ copies of H_2 , which occurs **whp** by (iii). Also by (iii), with probability $1 - o(n^{-2})$, every pair of vertices in G_i extends $O(1)$ copies of H_2 into H_3 ; call this event E_i .

Even after conditioning, incoming edges in the $(i + 1)$ -st round are independently and uniformly distributed over the $\binom{n}{2} - i = \Theta(n^2)$ unoccupied pairs of G_i . Therefore, the probability that both pairs complete K_4 , conditioned on survival through the i -th round, is $p_i = \Omega\left(\left(\frac{n^6 p^9}{n^2}\right)^2\right) = \Omega(n^8 p^{18})$. Letting i run from $3m$ to $4m$, we see that the probability that any strategy can survive for $4m$ rounds is at most

$$\begin{aligned} \mathbb{P} &\leq \mathbb{P}[-D] + \sum \mathbb{P}[-E_i] + \prod (1 - p_i) \leq o(1) + \exp\left\{-\sum p_i\right\} \\ &\leq o(1) + \exp\{-\Omega(n^2 p \cdot n^8 p^{18})\} = o(1) + e^{-\omega(1)} = o(1), \end{aligned}$$

which completes the proof. ■

4. AVOIDING K_t , GENERAL CASE

The previous section proved the threshold for avoiding K_t in the Achlioptas process with parameter r , when $t = 4$ and $r = 2$. The case $t = 3$ will be covered in Section 6, which considers all cycles C_t . In this section, we resolve all other cases, except for the special case $(t, r) = (4, 3)$ which requires more delicate analysis. We postpone this final case to Section 8.

Theorem. *For either $t \geq 5$ and $r \geq 2$, or $t = 4$ and $r \geq 4$, the threshold for avoiding K_t in the Achlioptas process with parameter $r \geq 2$ is $n^{2-\theta}$, where θ is defined as follows:*

$$s = \lfloor \log_r[(r - 1)t + 1] \rfloor, \quad \theta = \frac{r^s(t - 2) + 2}{r^s \left(\binom{t}{2} - s\right) + \frac{r^s - 1}{r - 1}}.$$

Before we begin the proof, let us prove an inequality that we will use in two claims in the lower bound, and the last claim of the upper bound.

Inequality 4.1. *Let $a > 2$, $b > 0$, and $r > 1$, and let s be a positive integer. Define the sequences $\{x_s, x_{s-1}, \dots, x_0\}$ and $\{y_s, y_{s-1}, \dots, y_0\}$ as follows. Set $x_s = a$ and $y_s = b$, and define the rest of the terms recursively by*

$$x_{k-1} = 2 + (x_k - 2)r, \quad y_{k-1} = 1 + y_k r.$$

Then for any $p \gg n^{-x_0/y_0}$, $n^k p^{y_k}$ is a positive power of n for every $k \in \{s, \dots, 1\}$.

Proof. Fix any $k \in \{s, \dots, 1\}$. One can easily solve the recursions for x_k and y_k to find:

$$x_k = r^{s-k}(a - 2) + 2, \quad y_k = r^{s-k}b + \frac{r^{s-k} - 1}{r - 1}.$$

Therefore,

$$\frac{x_k}{y_k} = \frac{r^{s-k}(a-2) + 2}{r^{s-k}b + \frac{r^{s-k}-1}{r-1}} = \frac{r^s(a-2) + 2r^k}{r^s b + \frac{r^s-r^k}{r-1}}.$$

By the original definition via the recursions, x_k and y_k are both positive, so the numerator and denominator of the final fraction above are positive. Yet as k decreases, the numerator decreases and the denominator increases. Therefore, $x_k/y_k > x_0/y_0$. In particular, since we assumed that $p \gg n^{-x_0/y_0}$, we conclude that $n^k p^{y_k}$ is a positive power of n , as desired. ■

Note that if we choose $a = v(K_t) = t$ and $b = e(K_t) - s = \binom{t}{2} - s$, then the above recursions produce x_0 and y_0 such that the fraction x_0/y_0 is equal to our θ . Let us now return to the proof of our thresholds for avoiding K_t .

Proof of Theorem (lower bound). The strategy is a natural extension of the one used to avoid K_4 . At any intermediate stage in the process, for any $1 \leq d \leq s$, consider a pair of points to be d -dangerous if d is the maximal integer such that the addition of an edge between them will create a copy of $K_t \setminus (s-d)e$. If there is no such d , consider the pair to be 0-dangerous. The strategy is then to make an arbitrary choice among the incoming edges that are minimally dangerous.

Let $m \ll n^{2-\theta}$, and let $p = 2m/n^2$. Again, we assume without loss of generality that $m \gg n^{2-\theta} / \log n$. Note that $n^{-\theta} / \log n \ll p \ll n^{-\theta}$. We will analyze the performance of our strategy by proving three successive claims:

- (i) With probability $1 - o(n^{-2s})$, G_m has $O(n^t p^{\binom{t}{2}-s})$ copies of $K_t \setminus se$, and every pair of vertices completes $O(1)$ copies of $K_t \setminus (s-1)e$.
- (ii) For each $k \in \{s, s-1, \dots, 2\}$, and constants x and y such that $(n^2 p)(\frac{n^x p^y}{n^2})^r$ is a positive power of n , statement (a) implies statement (b), which are defined as follows:
 - (a) With probability $1 - o(n^{-2k})$, G_m has $O(n^x p^y)$ copies of $K_t \setminus ke$, and every pair of vertices completes $O(1)$ copies of $K_t \setminus (k-1)e$.
 - (b) With probability $1 - o(n^{-2(k-1)})$, G_m has $O((n^2 p)(\frac{n^x p^y}{n^2})^r)$ copies of $K_t \setminus (k-1)e$, and every pair of vertices completes $O(1)$ copies of $K_t \setminus (k-2)e$.
- (iii) The probability of failure in m rounds is $o(1)$.

Again, we separate the proofs of the claims for clarity. At several points, we require certain inequalities whose rather tedious proofs would interfere with the exposition. The appendix contains the precise formulations of these statements.

Proof of (i). Lemma A.3 verifies that $K_t \setminus se$ is a balanced graph, and the $k = s$ case of Inequality 4.1 shows that $n^t p^{\binom{t}{2}-s}$ is a positive power of n , so Theorem 2.6 implies that the number of copies of $K_t \setminus se$ in G_m is $O(n^t p^{\binom{t}{2}-s})$ **wep**. For the second part of claim (i), Lemma A.4 verifies that $K_t \setminus se$ has the balanced extension property, and Inequality A.8 shows that $n^{t-2} p^{\binom{t}{2}-s}$ is a negative power of n . So, Corollary 2.9 shows that there is some constant C such that with probability $1 - o(n^{-2s})$, every pair of vertices in G_m completes at most C copies of $K_t \setminus (s-1)e$. This finishes claim (i). ■

Proof of (ii). Fix k, x , and y as specified, and let us show that (a) implies (b). First, since every graph of the form $K_t \setminus (k-2)e$ always contains some graph of the form $K_t \setminus (k-1)e$, (a) immediately implies that with probability $1 - o(n^{-2k})$, every pair of vertices completes $O(1)$ copies of $K_t \setminus (k-2)e$; this implies the second part of (b).

It remains to show the first part of (b). Fix some $i < m$ and consider the $(i + 1)$ -st round. In this round, the strategy will create one or more copies of $K_t \setminus (k - 1)e$ only if all r incoming edges span pairs that are at least $(s - k + 1)$ -dangerous (i.e., create copies of $K_t \setminus (k - 1)e$). The number of such pairs is at most $O(1)$ times the number of copies of $K_t \setminus ke$. Since $G_i \subset G_m$, statement (a) implies that with probability $1 - o(n^{-2k})$, G_i has $O(n^x p^y)$ copies of $K_t \setminus ke$ and every pair of vertices completes $O(1)$ copies of $K_t \setminus (k - 1)e$. Call this event A_i , and condition on it. Even after conditioning, incoming edges are still independently and uniformly distributed over the $\Omega(n^2)$ unoccupied pairs of G_i , so the probability that some new copies of $K_t \setminus (k - 1)e$ are created in this round is $O((\frac{n^x p^y}{n^2})^r)$. Also, by our conditioning, the number of newly created copies of $K_t \setminus (k - 1)e$ is still $O(1)$ even when this occurs. Therefore, the number of new copies of $K_t \setminus (k - 1)e$ in the $(i + 1)$ -st round is stochastically dominated by $O(1)$ times the Bernoulli random variable with parameter $O((\frac{n^x p^y}{n^2})^r)$. Letting i run through all m rounds, we see that with probability at least $1 - \sum \mathbb{P}[\neg A_i] \geq 1 - o(n^{-2(k-1)})$, the number of copies of $K_t \setminus (k - 1)e$ in G_m is $O(1) \cdot \text{Bin}[m, O((\frac{n^x p^y}{n^2})^r)]$. Since this binomial has expectation $\frac{n^2 p}{2} \cdot O((\frac{n^x p^y}{n^2})^r)$, which is a positive power of n by the assumption on x and y , a Chernoff bound implies that it is $O((n^2 p)(\frac{n^x p^y}{n^2})^r)$ **wep**. This finishes (ii). ■

Proof of (iii). The idea is to apply claim (i), and then to repeatedly apply claim (ii) until we obtain a high-probability upper bound on the number of copies of $K_t \setminus e$. Then, we complete the proof with essentially the same argument as in claim (iii) of the proof of the lower bound for avoiding K_4 .

To keep track of the exponents of n and p in the successive upper bounds, define the sequences $\{x_s, x_{s-1}, \dots, x_0\}$ and $\{y_s, y_{s-1}, \dots, y_0\}$ as in Inequality 4.1, which then verifies that $n^{x_k} p^{y_k}$ is a positive power of n for every $k \in \{s-1, \dots, 1\}$. Hence we can apply claims (i) and (ii) until we conclude that with probability $1 - o(n^{-2})$, G_m has $O(n^{x_1} p^{y_1})$ copies of $K_t \setminus e$.

Now fix some i and consider the probability that we lose in the $(i + 1)$ -st round. The strategy fails precisely when all r of the incoming edges span pairs that are s -dangerous (completing K_t), and the number of such pairs is at most $O(1)$ times the number of copies of $K_t \setminus e$. Yet since $G_i \subset G_m$, the previous paragraph shows that with probability $1 - o(n^{-2})$, G_i has $O(n^{x_1} p^{y_1})$ copies of $K_t \setminus e$. Call this event B_i , and condition on it. Even after conditioning, incoming edges are still independently and uniformly distributed over the $\Omega(n^2)$ unoccupied pairs of G_i , so the probability that all incoming edges complete K_t is $O((\frac{n^{x_1} p^{y_1}}{n^2})^r)$. Therefore, letting i run through all $m = n^2 p / 2$ rounds, a union bound shows that the probability that we are forced to complete a copy of K_t is $\mathbb{P} \leq O((n^2 p)(\frac{n^{x_1} p^{y_1}}{n^2})^r) + \sum \mathbb{P}[\neg B_i] = O(n^{x_0} p^{y_0}) + o(1)$. This in turn is $o(1)$ because we assumed that $p \ll n^{-\theta}$ with $\theta = x_0 / y_0$. This completes the proof. ■

Proof of Theorem (upper bound). Let $m \gg n^{2-\theta}$, and let $p = 2m/n^2$. We will show that **whp**, any strategy fails within $\Theta(m)$ rounds, which we again break into periods of length m . We may assume that $m \ll n^{2-\theta} \log n$ without loss of generality. Note that $n^{-\theta} \ll p \ll n^{-\theta} \log n$.

As in the proof of the upper bound for avoiding K_4 , we will specify a sequence of graphs such that each graph is obtained from the previous one by adding a single edge. Let $H_1 = K_{\lfloor \frac{t}{2} \rfloor, \lceil \frac{t}{2} \rceil}$ (the largest bipartite subgraph of K_t), and arbitrarily choose the rest of the sequence $\{H_2, H_3, \dots, H_f\}$, where $H_f = K_t$, by adding one missing edge at a time. So, $f = 1 + \binom{t}{2} - \lfloor \frac{t}{2} \rfloor \lceil \frac{t}{2} \rceil$, which is a constant because we assumed t to be fixed. Our result follows from the following five claims:

- (i) G_m contains $\Omega(n^t p^{e(H_1)})$ copies of H_1 **whp**.
- (ii) Let k be a positive integer for which $n^{t-2} p^{e(H_{k-1})}$ is a positive power of n . Then G_{km} contains $\Omega(n^t p^{e(H_k)})$ copies of H_k **whp**.
- (iii) $G_{(f-s)m}$ contains $\Omega(n^t p^{e(H_{f-s})})$ copies of H_{f-s} **whp**. Also, $n^{t-2} p^{e(H_{f-s})}$ is a negative power of n ; hence with probability $1 - o(n^{-2})$, every pair of vertices in $G_{(f-s+1)m}$ extends $O(1)$ copies of H_{f-s} into H_{f-s+1} .
- (iv) For each $k \in \{s, s-1, \dots, 2\}$, and constants x and y such that $n^x p^y \ll n^2$ and $(n^2 p)^{\binom{n^x p^y}{n^2}}$ is a positive power of n , statement (a) implies statement (b), which are defined as follows:
 - (a) $G_{(f-k)m}$ contains $\Omega(n^x p^y)$ copies of H_{f-k} **whp**, and with probability $1 - o(n^{-2})$, every pair of vertices in $G_{(f-k+1)m}$ extends $O(1)$ copies of H_{f-k} into H_{f-k+1} .
 - (b) $G_{(f-k+1)m}$ contains $\Omega((n^2 p)^{\binom{n^x p^y}{n^2}})$ copies of H_{f-k+1} **whp**, and with probability $1 - o(n^{-2})$, every pair of vertices in $G_{(f-k+2)m}$ extends $O(1)$ copies of H_{f-k+1} into H_{f-k+2} .
- (v) The probability of survival through $fm = \Theta(m)$ rounds is $o(1)$.

Proof of (i). We will actually prove that G_m contains $\Omega(n^t p^{e(H_1)})$ copies of H_1 with certainty, not just **whp**. However, the rest of the claims only require a **whp** result in claim (i), so we keep it there for the purpose of generality.

Since we assumed that $p \gg n^{-\theta}$ and Inequality A.6 bounds $-\theta \geq -\lfloor \frac{t}{2} \rfloor^{-1}$, Lemma 2.2 implies that the number of copies of the complete bipartite graph $H_1 = K_{\lfloor \frac{t}{2} \rfloor, \lceil \frac{t}{2} \rceil}$ in any m -edge graph is $\Omega(n^t p^{e(H_1)})$. ■

Proof of (ii). We proceed inductively. The base case of the induction follows from claim (i). Now, suppose k satisfies the property that $n^{t-2} p^{e(H_{k-1})}$ is a positive power of n , and $G_{(k-1)m}$ contains $\Omega(n^t p^{e(H_{k-1})})$ copies of H_{k-1} **whp**. We will show that G_{km} contains $\Omega(n^t p^{e(H_k)})$ copies of H_k **whp**.

Let us begin by conditioning on the high-probability event A from our inductive assumption: that $G_{(k-1)m}$ contains $\Omega(n^t p^{e(H_{k-1})})$ copies of H_{k-1} . Now consider the $(i + 1)$ -st round, where $(k - 1)m \leq i < km$. Since $G_i \supset G_{(k-1)m}$, the total number of copies of H_{k-1} in G_i is $\Omega(n^t p^{e(H_{k-1})})$ by our conditioning.

Lemma A.5 verifies that (H_{k-1}, H_k) is a balanced extension pair, and we assumed that $n^{t-2} p^{e(H_{k-1})}$ was a positive power of n , so Theorem 2.8(i) establishes that **whp**, every pair of vertices in G_{km} extends $O(n^{t-2} p^{e(H_{k-1})})$ copies of H_{k-1} into H_k . Since $G_i \subset G_{km}$, the same bound holds for G_i **whp**; call that event B_i , and condition on it.

For a pair of vertices $\{a, b\}$, let $n_{a,b}$ be the number of copies of H_{k-1} that the pair $\{a, b\}$ extends into H_k . Recall that this definition does not depend on the presence of an edge between a and b . Let us estimate the average value of $n_{a,b}$ over all pairs. Since H_{k-1} differs from H_k at exactly one edge, each copy of H_{k-1} has a pair at which it contributes $+1$ to the sum $\sum n_{a,b}$. Therefore, averaging over all $\binom{n}{2}$ pairs of vertices, we obtain that the average number of copies of H_{k-1} that are extended to H_k at a pair is $\Omega(n^{t-2} p^{e(H_{k-1})})$.

On the other hand, every pair of vertices in G_i extends $O(n^{t-2} p^{e(H_{k-1})})$ copies of H_{k-1} into H_k . Therefore, at least a constant fraction $\gamma = \Omega(1)$ of all $\binom{n}{2}$ pairs have the property of extending $\Omega(n^{t-2} p^{e(H_{k-1})})$ copies of H_{k-1} into H_k . Let P be the set of all such pairs. Regardless of the choice of strategy, if all r incoming edges span pairs in P , we will be forced to create $\Omega(n^{t-2} p^{e(H_{k-1})})$ copies of H_k . Since $i = o(n^2) = o(|P|)$ and incoming edges are uniformly distributed over the $\binom{n}{2} - i = (1 + o(1))\binom{n}{2}$ unoccupied pairs, we conclude that the probability that all incoming edges span pairs in P is $q \geq (1 + o(1))\gamma^r = \Omega(1)$.

Let i run from $(k - 1)m$ to km . Then, up to an error probability of at most $\mathbb{P}[\neg A] + \sum \mathbb{P}[\neg B_i] = o(1)$, the number of copies of H_k in G_{km} is at least $\text{Bin}(m, q) \cdot \Omega(n^{t-2}p^{e(H_{k-1})})$. By the Chernoff bound, the binomial factor exceeds $mq/2 = \Omega(n^2p)$ **wep**; thus, **whp** G_{km} has $\Omega(n^2p \cdot n^{t-2}p^{e(H_{k-1})}) = \Omega(n^t p^{e(H_k)})$ copies of H_k . ■

Proof of (iii). The first part follows directly from claim (ii), because Inequality A.7 verifies that $n^{t-2}p^{e(H_{f-s}-1)}$ is a positive power of n . For the second part, (H_{f-s}, H_{f-s+1}) is a balanced extension pair by Lemma A.5, and $n^{t-2}p^{e(H_{f-s})}$ is a negative power of n by Inequality A.8. Therefore, Theorem 2.8(ii) shows that there is some constant C such that with probability $1 - o(n^{-2})$, every pair of vertices in $G_{(f-s+1)m}$ extends at most C copies of H_{f-s} into H_{f-s+1} . This finishes claim (iii). ■

Proof of (iv). Fix k, x , and y as specified in the statement, and assume statement (a). Let us begin by establishing the second part of (b). Lemma A.5 verifies that (H_{f-k+1}, H_{f-k+2}) is a balanced extension pair, and Inequality A.8 shows that $n^{t-2}p^{e(H_{f-k+1})}$ is a negative power of n for $k \leq s$. Therefore, Theorem 2.8(ii) shows that there is some constant C such that with probability $1 - o(n^{-2})$, every pair of vertices in $G_{(f-k+2)m}$ extends at most C copies of H_{f-k+1} into H_{f-k+2} . This finishes the second part of (b).

It remains to prove the first part of (b). Consider the $(i + 1)$ -st round, with $(f - k)m \leq i < (f - k + 1)m$. Regardless of the choice of strategy, if all r incoming edges span pairs that extend copies of H_{f-k} into H_{f-k+1} , we will create a copy of H_{f-k+1} . Let P be the set of all such pairs. We need a lower bound on $|P|$.

Condition on the high-probability event C of (a) that $G_{(f-k)m}$ contains $\Omega(n^x p^y)$ copies of H_{f-k} . Since $G_i \subset G_{(f-k+1)m}$, (a) implies that with probability $1 - o(n^{-2})$, every pair of vertices in G_i extends $O(1)$ copies of H_{f-k} into H_{f-k+1} . Call this event D_i , and condition on it.

Note that every copy of H_{f-k} contributes a pair to P which extends H_{f-k} into H_{f-k+1} , namely the pair at which it is missing an edge compared to H_{f-k+1} . On the other hand, every such pair was counted at most a constant number of times, since every pair in G_i extends $O(1)$ copies of H_{f-k} into H_{f-k+1} . This implies that $|P| = \Omega(n^x p^y)$. The incoming edges are uniformly distributed over all unoccupied pairs. If at least half of the pairs in P were occupied, then we would have $\Omega(n^x p^y)$ copies of H_{f-k+1} . Yet this would already give us the conclusion of (b) since:

$$n^x p^y \gg (n^2 p) \left(\frac{n^x p^y}{n^2} \right) \gg (n^2 p) \left(\frac{n^x p^y}{n^2} \right)^r.$$

(The first inequality is because $p \ll 1$, and the second inequality follows from the assumption that $n^x p^y \ll n^2$.) Otherwise, if less than half of the pairs in P are occupied, then the probability that all incoming edges span pairs in P (hence forcing the creation of a copy of H_{f-k+1}) is $q \geq (1 + o(1)) \left(\frac{|P|/2}{n^2/2} \right)^r = \Omega\left(\left(\frac{n^x p^y}{n^2}\right)^r\right)$.

Letting i run from $(f - k)m$ to $(f - k + 1)m$, we see that with error probability at most $\mathbb{P}[\neg C] + \sum \mathbb{P}[\neg D_i] = o(1)$, either we already obtained the conclusion of (b), or the total number of copies of H_{f-k+1} is at least $\text{Bin}(m, q)$. The expectation of the binomial is $\left(\frac{n^2 p}{2}\right)q = \Omega\left((n^2 p) \left(\frac{n^x p^y}{n^2}\right)^r\right)$, which is a positive power of n by assumption. Hence, by the Chernoff bound, $G_{(f-k+1)m}$ has $\Omega\left((n^2 p) \left(\frac{n^x p^y}{n^2}\right)^r\right)$ copies of H_{f-k+1} **whp**. ■

Proof of (v). The result of claim (iii) plugs in directly to claim (iv), which we may iterate until it gives us a lower bound on the number of copies of $H_{f-1} = K_t \setminus e$ and an upper bound on the number of copies of H_{f-1} that any pair extends into $H_f = K_t$.

To keep track of exponents in the successive lower bounds, define the sequences $\{x_s, x_{s-1}, \dots, x_0\}$ and $\{y_s, y_{s-1}, \dots, y_0\}$ exactly as in Inequality 4.1. To verify that we can indeed iterate claim (iv), we must show that for all $k \in \{s, s-1, \dots, 2\}$, we have that $n^{x_k} p^{y_k} \ll n^2$, and $n^{x_{k-1}} p^{y_{k-1}}$ is a positive power of n . The first statement follows from an easy induction: claim (iii) establishes it for $k = s$, and if $n^{x_k} p^{y_k} \ll n^2$, then $\frac{n^{x_k} p^{y_k}}{n^2} \ll 1$, so combined with $p \ll 1$, we see that $n^{x_{k-1}} p^{y_{k-1}} = (n^2 p) \left(\frac{n^{x_k} p^{y_k}}{n^2}\right)^r \ll n^2$. The second statement is verified by Inequality 4.1. Therefore, we arrive at the result that $G_{(f-1)m}$ contains $\Omega(n^{x_1} p^{y_1})$ copies of $H_{f-1} = K_t \setminus e$ **whp**. Call this event E , and condition on it. We also find that with probability $1 - o(n^{-2})$, every pair of vertices in G_{fm} extends $O(1)$ copies of H_{f-1} into H_f (i.e., completes $O(1)$ copies of K_t). The same probability bound also holds in G_i for any $i \leq fm$, because $G_i \subset G_{fm}$; let F_i be the corresponding event.

Now consider the $(i+1)$ -st round, where $(f-1)m \leq i < fm$. Regardless of the choice of strategy, if all r incoming edges span pairs that complete copies of K_t , we will lose. We can bound the number of such pairs by $\Omega(n^{x_1} p^{y_1})$ by conditioning on the above events E and F_i . Even after conditioning, the incoming edges in this round still independent and uniformly distributed over the $\binom{n}{2} - i = \Theta(n^2)$ unoccupied pairs of G_i . Therefore, the probability that all r pairs complete K_t , conditioned on survival through the i -th round, is $p_i = \Omega\left(\left(\frac{n^{x_1} p^{y_1}}{n^2}\right)^r\right)$. Letting i run from $(f-1)m$ to fm , we see that the probability that any strategy can survive for fm rounds is at most

$$\begin{aligned} \mathbb{P} &\leq \mathbb{P}[\neg E] + \sum \mathbb{P}[\neg F_i] + \prod (1 - p_i) \leq o(1) + \exp\left\{-\sum p_i\right\} \\ &\leq o(1) + \exp\left\{-\Omega\left((n^2 p) \left(\frac{n^{x_1} p^{y_1}}{n^2}\right)^r\right)\right\} = o(1) + \exp\{-\Omega(n^{x_0} p^{y_0})\}. \end{aligned}$$

This in turn is $o(1)$ because we assumed that $p \gg n^{-\theta}$ with $\theta = x_0/y_0$. This completes the proof. ■

5. ABSTRACTION INTO GENERAL ARGUMENT

Note that we structured our exposition of the previous section in the following manner. The arguments did not directly use properties of the specific graph that we were avoiding (K_t). Rather, they were linked to lemmas and inequalities that proved certain properties (e.g., balanced-ness, etc.) about K_t . Let us now isolate these necessary “ingredients” that one can plug in to our general machinery to prove thresholds.

For the rest of this section, let H be the fixed graph which we wish to avoid. Our arguments allow one to prove the threshold for avoiding H in the Achlioptas process with parameter r simply by specifying several parameters, and then proving some lemmas and inequalities that do not need to refer to the Achlioptas process at all. We first describe the parameters.

- s : this was the number of levels of danger considered by the avoidance strategy in the proof of the lower bound. At any intermediate stage in the process, for any $1 \leq d \leq s$, we considered a pair of points to be d -dangerous if d was the maximal integer such that the addition of an edge between them created a copy of $H \setminus (s-d)e$. If there was no

such d , we considered the pair to be 0-dangerous. Recall that the strategy was then to make an arbitrary choice among the incoming edges that were minimally dangerous.

- A sequence of graphs $\{H_1, \dots, H_f\}$ sharing the same vertex set, with each successive graph containing exactly one more edge: this was used in the upper bound argument to iteratively prove lower bounds on the number of copies of H_i , proceeding from $i = 1$ to $i = f$.

The correct choice of s then determined θ , the negative exponent in the threshold (in terms of p) for avoidance:

$$\theta = \frac{r^s(v(H) - 2) + 2}{r^s(e(H) - s) + \frac{r^s - 1}{r - 1}}.$$

Assuming that the parameters were suitably chosen, one then only needed to establish the following lemmas and inequalities in order to prove that the threshold for avoiding H in the Achlioptas process with parameter r is $n^{2-\theta}$.

For proof of lower bound. Here, $n^{-\theta} / \log n \ll p \ll n^{-\theta}$.

1. $H \setminus se$ is a balanced graph. This allowed us to prove in claim (i) that **wep**, G_m has $O(n^{v(H)} p^{e(H)-s})$ copies of $H \setminus se$. For $H = K_t$, this was provided by Lemma A.3.
2. $H \setminus se$ has the balanced extension property, and $n^{v(H)-2} p^{e(H)-s}$ is a negative power of n . This allowed us to prove in claim (i) that with probability $1 - o(n^{-2s})$, every pair of vertices in G_m completes $O(1)$ copies of $H \setminus (s - 1)e$. For $H = K_t$, these were provided by Lemma A.4 and Inequality A.8.

For proof of upper bound. Here, $n^{-\theta} \ll p \ll n^{-\theta} \log n$.

1. G_m contains $\Omega(n^{v(H_1)} p^{e(H_1)})$ copies of H_1 **whp**. This was claim (i), and for $H = K_t$, it was provided by the extremal estimate on the number of $K_{s,t}$ (Lemma 2.2), along with Inequality A.6, which assured that p was large enough to apply the extremal result.
2. Each consecutive pair (H_k, H_{k+1}) is a balanced extension pair. This was used throughout the proof of the upper bound, and for $H = K_t$, it was provided by Lemma A.5.
3. $n^{v(H)-2} p^{e(H)-s-1}$ is a positive power of n . This was used in claim (iii) to show that we could iterate the argument of claim (ii) enough times to conclude that $G_{(f-s)m}$ contained $\Omega(n^{v(H_{f-s})} p^{e(H_{f-s})})$ copies of H_{f-s} **whp**. For $H = K_t$, this was provided by Inequality A.7.
4. $n^{v(H)-2} p^{e(H)-s}$ is a negative power of n . This was used in claim (iii) to transition to the next inductive process, which relied on the copies of H_{f-s} not being too concentrated on any pair of vertices. **Note:** this statement was already required above for the lower bound, so we do not need to check it again.

6. AVOIDING CYCLES

Now we show by example how to use our machinery to prove avoidance thresholds. We start with an easy application which completely solves the problem for cycles C_t . In light of the previous section, we only need to provide the required parameters, lemmas, and inequalities. We will specify these in the same order that they were presented in the previous section. This will prove the following theorem.

Theorem. For $t \geq 3$, the threshold for avoiding C_t in the Achlioptas process with parameter $r \geq 2$ is $n^{2 - \frac{r(t-2)+2}{r(t-1)+1}}$.

Proof. We use the parameter $s = 1$, and the sequence of graphs $H_1 = C_t \setminus e, H_2 = C_t$. This gives the threshold $n^{2-\theta}$, where $\theta = \frac{r^s(v(C_t)-2)+2}{r^s(e(C_t)-s)+\frac{r^s-1}{r-1}} = \frac{r(t-2)+2}{r(t-1)+1}$, which matches the claimed result. Now we need to provide the required lemmas and inequalities. For the reader's convenience, we have reproduced the italicized statements from Section 5.

For proof of lower bound. Here, $n^{-\theta} / \log n \ll p \ll n^{-\theta}$.

1. $C_t \setminus e$ is a balanced graph. This is obvious.
2. $C_t \setminus e$ has the balanced extension property, and $n^{v(C_t)-2} p^{e(C_t)-1} = n^{t-2} p^{t-1}$ is a negative power of n . The first part is obvious. For the second, since $p \ll n^{-\frac{r(t-2)+2}{r(t-1)+1}}$, we must establish that $(t-2) - (t-1)\frac{r(t-2)+2}{r(t-1)+1} < 0$. Routine algebra shows that the left hand side equals $-\frac{t}{r(t-1)+1}$, which is certainly negative when $t \geq 3, r \geq 2$.

For proof of upper bound. Here, $n^{-\theta} \ll p \ll n^{-\theta} \log n$.

1. G_m contains $\Omega(n^{v(H_1)} p^{e(H_1)})$ copies of H_1 *whp*. The average degree of G_m is precisely np by the definition of $p = 2m/n^2$. We show in item #3 below that np is a positive power of n , so it tends to infinity with n . Thus, we may apply Lemma 2.1, an extremal result counting the number of paths, to conclude that G_m contains at least $(1+o(1))n(np)^{t-1}$ copies of the t -vertex path H_1 , as desired.
2. (H_1, H_2) is a balanced extension pair. This is easy to see.
3. $n^{v(C_t)-2} p^{e(C_t)-1-1} = (np)^{t-2}$ is a positive power of n . It suffices to show that np is a positive power of n . Since $p \gg n^{-\frac{r(t-2)+2}{r(t-1)+1}}$, this amounts to proving that $1 - \frac{r(t-2)+2}{r(t-1)+1} > 0$. Routine algebra shows that the left hand side equals $\frac{r-1}{r(t-1)+1}$, which is certainly positive when $t \geq 3, r \geq 2$.

As we have provided all of the necessary ingredients to apply our machinery, we are done. ■

7. AVOIDING $K_{t,t}$

Now we show a more complex application of our machinery, which completely solves the problem for $K_{t,t}$. This will prove the following theorem.

Theorem. Suppose that $t \geq 3$ and $r \geq 2$ are fixed integers. The threshold for avoiding $K_{t,t}$ in the Achlioptas process with parameter r is $n^{2-\theta}$, where θ is defined as follows:

$$s = \lceil \log_r \lceil (r-1)t + 1 \rceil \rceil, \quad \theta = \frac{r^s(2t-2)+2}{r^s(t^2-s)+\frac{r^s-1}{r-1}}.$$

7.1. Parameters

The value of s is already specified in the statement of the theorem, so we proceed to give the sequence of graphs $\{H_1, \dots, H_f\}$. The sequences are quite different depending on the parity of t , so we describe them separately.

Case 1: t is even. Let H_1 be the 4-partite graph with parts V_1, V_2, V_3, V_4 , each of size $t/2$, and edges such that (V_1, V_2) , (V_1, V_4) , and (V_3, V_2) are complete bipartite graphs. Let $\{H_2, \dots, H_{1+(t/2)}\}$ be obtained by successively adding single edges until $H_{1+(t/2)}$ has a perfect matching between V_3 and V_4 . Then, arbitrarily choose the rest of the sequence $\{H_{2+(t/2)}, \dots, H_f\}$ by adding one edge at a time, until the final term is the complete bipartite graph $K_{t,t}$ with bipartition $(V_1 \cup V_3, V_2 \cup V_4)$. Note that $f = 1 + t^2/4$.

Case 2: t is odd. Let H_1 be a 6-partite graph with parts $\{V_i\}_1^6$ such that V_3 and V_4 are singletons, and the other four parts each have size $\lfloor t/2 \rfloor$. The edges are as follows: the two pairs (V_1, V_2) and (V_5, V_6) are each complete bipartite graphs, the vertex in V_3 is adjacent to all of $V_2 \cup V_4 \cup V_6$, and the vertex in V_4 is adjacent to all of $V_1 \cup V_3 \cup V_5$. There are no more edges.

Let $\{H_2, \dots, H_{1+\lfloor t/2 \rfloor}\}$ be obtained by successively adding single edges until $H_{1+\lfloor t/2 \rfloor}$ has a perfect matching between V_1 and V_6 . To create the next $\lfloor t/2 \rfloor$ graphs in the sequence, we put down a matching between V_5 and V_2 , one edge at a time. Finally, arbitrarily choose the rest of the sequence $\{H_{2+2\lfloor t/2 \rfloor}, \dots, H_f\}$ by adding one edge at a time, until the final term is the complete bipartite graph $K_{t,t}$ with bipartition $(V_1 \cup V_3 \cup V_5, V_2 \cup V_4 \cup V_6)$. Note that $f = 1 + 2\lfloor t/2 \rfloor^2$.

7.2. Lemmas and Inequalities

Next, we provide the required lemmas and inequalities. For the reader's convenience, we have reproduced the italicized statements from Section 5.

For proof of lower bound. Here, $n^{-\theta} / \log n \ll p \ll n^{-\theta}$.

1. $K_{t,t} \setminus se$ is a balanced graph. This is now provided by Lemma B.1. Actually, the graph is not balanced when $t = 3$ and $r = 2$, but in that particular case, Lemma B.1 additionally proves that the number of copies of $K_{t,t} \setminus se$ in G_m is still $O(n^{v(H)} p^{e(H)-s})$ **wep**, which is all we really need.
2. $K_{t,t} \setminus se$ has the balanced extension property, and $n^{v(K_{t,t})-2} p^{e(K_{t,t})-s}$ is a negative power of n . These are now provided by Lemma B.2 and Inequality B.8.

For proof of upper bound. Here, $n^{-\theta} \ll p \ll n^{-\theta} \log n$.

1. G_m contains $\Omega(n^{v(H_1)} p^{e(H_1)})$ copies of H_1 **whp**. This time, we use Inequality B.6 to show that $-\theta > -2/t$. Since we assume that $p \gg n^{-\theta}$ for the upper bound argument, this provides the condition required to apply either Lemma 7.1 if t is even, or Lemma 7.2 if t is odd. Both lemmas (presented below) lead to the required final statement.
2. Each consecutive pair (H_k, H_{k+1}) is a balanced extension pair. This is now provided by Lemma B.4 if t is even, and by Lemma B.5 if t is odd.
3. $n^{v(K_{t,t})-2} p^{e(K_{t,t})-s-1}$ is a positive power of n . This is provided by Inequality B.7.

7.3. Proofs of Supporting Lemmas

We conclude this section by proving the two lemmas that provide the first component of the proof of the upper bound. We start with the lemma that is used when t is even.

Lemma 7.1. *For any fixed positive integers k and l , consider the following 4-partite graph, which we call H . Let the parts be V_1, V_2, V_3, V_4 , with $|V_1| = |V_2| = k$ and $|V_3| = |V_4| = l$, and place edges such that (V_1, V_2) , (V_1, V_4) , and (V_2, V_3) are complete bipartite graphs. There are no more edges. Then, there exists a constant c_k such that for any $p \gg n^{-1/k}$, every graph with n vertices and $\binom{n}{2}p$ edges contains at least $(c_k + o(1))n^{2k+2l}p^{k^2+2kl}$ copies of H .*

Proof. Let us fix an ambient graph G with n vertices and $\binom{n}{2}p$ edges. By Lemma 2.2, the number of copies of $K_{k,k}$ in G is at least $(1 + o(1))n^{2k}p^{k^2}$. Recall that the k -codegree of a set U of k distinct vertices is the number of vertices that are adjacent to all of U . Let us say that a copy of $K_{k,k}$ is *deficient* if either of the sides of its bipartition has k -codegree less than $\frac{1}{2}np^k$ in G . We claim that at most $\frac{1}{2} + o(1)$ of the copies of $K_{k,k}$ are deficient.

To see this, note that if an ordered k -tuple of distinct vertices has k -codegree less than $\frac{1}{2}np^k$, then it can extend to at most $(\frac{1}{2}np^k)^k$ copies of $K_{k,k}$. The number of such k -tuples is at most n^k ; therefore, the number of deficient copies of $K_{k,k}$ is at most $n^k (\frac{1}{2}np^k)^k \leq \frac{1}{2}n^{2k}p^{k^2}$, as claimed.

Yet each nondeficient copy of $K_{k,k}$ extends to at least

$$\binom{\frac{1}{2}np^k - 2k}{l} l! \cdot \binom{\frac{1}{2}np^k - 2k - l}{l} l!$$

copies of H . This is because we may consider the copy of $K_{k,k}$ to be $V_1 \cup V_2$, we choose V_3 from the common neighborhood of V_2 excluding the $2k$ vertices in $V_1 \cup V_2$, and finally we choose V_4 from the common neighborhood of V_1 excluding the $2k+l$ vertices in $V_1 \cup V_2 \cup V_3$. Since we assumed that $p \gg n^{-1/k}$, the binomial coefficients are asymptotically monomials of degree l , so we conclude that each nondeficient copy of $K_{k,k}$ extends to $\Omega((np^k)^l \cdot (np^k)^l) = \Omega(n^{2l}p^{2kl})$ copies of H . Since there are always at least $(\frac{1}{2} + o(1))n^{2k}p^{k^2}$ non-deficient copies of $K_{k,k}$, we conclude that the number of copies of H is always $\Omega(n^{2k+2l}p^{k^2+2kl})$, as claimed. ■

Using Lemma 7.1 as a building block, we now prove the lemma that provides the first component of the upper bound when t is odd. Actually, we prove a result for G_{2m} instead of G_m , but this does not matter for the purpose of the general argument.

Lemma 7.2. *Let k be a positive integer. Let H be a 6-partite graph with parts $\{V_i\}_1^6$ such that V_3 and V_4 are singletons, and the other four parts each have size k . Let the edges of H be as follows: the two pairs (V_1, V_2) and (V_5, V_6) are each complete bipartite graphs, the vertex in V_3 is adjacent to all of $V_2 \cup V_4 \cup V_6$, and the vertex in V_4 is adjacent to all of $V_1 \cup V_3 \cup V_5$. There are no more edges.*

Consider G_{2m} , the graph after the $2m$ -th round of the Achlioptas process with parameter $r \geq 2$. Let $p = 2m/n^2$, and suppose that $p \gg n^{-\theta}$ with $-\theta > -1/(k + \frac{1}{2})$. Then G_{2m} contains $\Omega(n^{v(H)}p^{e(H)})$ copies of H whp.

Proof. Let H_1 be the subgraph of H induced by $V_1 \cup V_2 \cup V_3 \cup V_4$, and let H_0 be the subgraph of H_1 with the edge between V_3 and V_4 deleted. Observe that we can find a copy of H in a graph by first looking for a pair of vertices for the site of the edge between V_3 and V_4 , and then looking for two disjoint copies of H_0 that are extended into H_1 by that pair.

Consider the $(i + 1)$ -st turn, for some $m \leq i < 2m$. By Lemma 7.1, G_m (and hence $G_i \supset G_m$) always contains $\Omega(n^{2k+2}p^{k^2+2k})$ copies of H_0 . Lemma B.3 verifies that (H_0, H_1) is a balanced extension pair, and $n^{2k}p^{k^2+2k}$ is a positive power of n because we assumed that

$p \gg n^{-\theta}$ with $-\theta > -1/(k + \frac{1}{2})$ and $k \geq 1$. Thus, Theorem 2.8(i) establishes that **wep**, every pair of vertices in $G_i \subset G_{2m}$ extends $O(n^{2k}p^{k^2+2k})$ copies of H_0 into H_1 . Call this event A_i , and condition on it.

For a pair of vertices $\{a, b\}$, let $n_{a,b}$ be the number of copies of H_0 that the pair $\{a, b\}$ extends into H_1 . Recall that this definition does not depend on the presence of an edge between a and b . Let us estimate the average value of $n_{a,b}$ over all pairs. Since H_0 differs from H_1 at exactly one edge, each copy of H_0 has a pair at which it contributes $+1$ to the sum $\sum n_{a,b}$. Therefore, averaging over all $\binom{n}{2}$ pairs of vertices, we obtain that the average number of copies of H_0 that are extended to H_1 at any pair is $\Omega(n^{2k+2}p^{k^2+2k})$.

On the other hand, by our conditioning, every pair of vertices in G_i extends $O(n^{2k}p^{k^2+2k})$ copies of H_0 into H_1 . Therefore, at least a constant fraction $\gamma = \Omega(1)$ of all $\binom{n}{2}$ pairs have the property of extending $\Omega(n^{2k}p^{k^2+2k})$ copies of H_0 into H_1 . Let P be the set of all such pairs. Regardless of the choice of strategy, if all r incoming edges span pairs in P , we will be forced to choose a pair in P . This will create $\Omega((n^{2k}p^{k^2+2k})^2) = \Omega(n^{4k}p^{2k^2+4k})$ pairs of copies of H_0 that are extended to H_1 by the chosen pair. Such a pair of copies of H_0 would become a new copy of H after the edge is added, if the pair of copies were disjoint. If the pair of copies of H_0 is not disjoint, then let us say that they create a degenerate copy of H . For now, let us count degenerate copies of H along with the true copies of H . Later, we will show that the degenerate copies are vastly outnumbered by true copies of H .

Since $i = o(n^2) = o(|P|)$ and incoming edges are uniformly distributed over the $\binom{n}{2} - i = (1 + o(1))\binom{n}{2}$ unoccupied pairs, we conclude that the probability that all incoming edges span pairs in P is $q \geq (1 + o(1))\gamma^r = \Omega(1)$. Let i run from m to $2m$. Then **wep**, the number of (possibly degenerate) copies of H in G_{2m} is at least $\text{Bin}(m, q) \cdot \Omega(n^{4k}p^{2k^2+4k})$. By the Chernoff bound, the binomial factor exceeds $mq/2 = \Omega(n^2p)$ **wep**, so we conclude that G_{2m} has $\Omega(n^2p \cdot n^{4k}p^{2k^2+4k}) = \Omega(n^{v(H)}p^{e(H)})$ (possibly degenerate) copies of H **whp**.

To finish the proof of this lemma, we must show that the number of degenerate copies of H in G_{2m} is $o(n^{v(H)}p^{e(H)})$ **whp**. For convenience, we will work with $G(n, p)$ instead of G_{2m} because Lemma 2.4 shows that we may couple G_{2m} with $G(n, 4rp)$, and the constant $4r$ disappears under the “ $o(\cdot)$ ” notation. Note that the underlying graph of a degenerate copy of H is a superposition of two copies of $K_{k+1, k+1}$, overlapping on at least 3 vertices. So, let us consider any such superposition, and call the underlying graph F . Let $v' = v(F)$ and $e' = e(F)$. The copies overlap on at least 3 vertices, so $v' < v(H)$. It is easy to check that since $K_{k+1, k+1}$ is a balanced graph, $\frac{e'}{v'} \geq \frac{e(H)}{v(H)}$. So, the expected number of copies of F in $G(n, p)$ is:

$$\mathbb{E} \leq n^{v'} p^{e'} = (np^{e'/v'})^{v'} \leq (np^{e(H)/v(H)})^{v'} = (n^{v(H)} p^{e(H)})^{v'/v(H)}.$$

Now, we assumed that $p \gg n^{-1/(k+\frac{1}{2})}$, so $n^{v(H)}p^{e(H)} \gg 1$ because $v(H) = 4k + 2$ and $e(H) = 2k^2 + 4k + 1$. Furthermore, $v' < v(H)$, so Markov’s inequality implies that **whp**, $G(n, p)$ has $o(n^{v(H)}p^{e(H)})$ copies of F . Since each copy of F can account for at most a constant number (depending only on k) of degenerate copies of H , and there is only a constant number of non-isomorphic superpositions F , we conclude that **whp**, $G(n, p)$ has $o(n^{v(H)}p^{e(H)})$ degenerate copies of H . This completes the proof of the lemma. ■

8. AVOIDING K_4 IN THE ACHLIOPTAS PROCESS WITH PARAMETER 3

To apply the machinery of Section 5, one needs to prove that certain quantities are positive or negative powers of n . In our study of avoiding cycles, cliques, and complete bipartite graphs, the only case in which we encounter a key exponent that is not separated from zero is when we are avoiding K_4 in the Achlioptas process with parameter 3.

However, the separation of the exponent from zero was merely a convenience which allowed us to bound maxima of families of random variables (e.g., the maximum codegree in a graph) **whp**. When we do not have this condition, we may instead bound the entire distribution of the family.

Lemma 8.1. *Let $n^{-1/2} \ll p \ll n^{-1/2} \log n$. Then $G(n, p)$ satisfies the following property **whp**: all codegrees are at most $np^2 \log n$, and for every integer $4 \leq k \leq \log n$, the number of pairs with codegree at least kn^2 is at most n^2/k^3 .*

This result, which we prove at the end of this section, allows us to prove our final threshold.

Theorem. *The threshold for avoiding K_4 in the Achlioptas process with parameter 3 is $n^{3/2}$.*

Proof of lower bound. A shortsighted strategy works in this instance: arbitrarily select any one of the incoming edges that does not create a copy of K_4 . Let $m \ll n^{3/2}$, and let $p = 2m/n^2$. Again, we assume without loss of generality that $m \gg n^{3/2}/\log n$. Note that $n^{-1/2}/\log n \ll p \ll n^{-1/2}$. We will analyze the performance of our strategy by proving two successive claims:

- (i) G_m has $O(n^4 p^5)$ copies of $K_4 \setminus e$ **wep**.
- (ii) The probability of failure in m rounds is $o(1)$.

The interested reader may check that if we followed the recipe for avoiding K_r in Section 4, we would start by counting copies of $K_4 \setminus 2e$ instead of $K_4 \setminus e$. This is essentially the only change in the lower bound argument, but we provide the details below for completeness.

For (i), $K_4 \setminus e$ is balanced and $n^4 p^5$ is a positive power of n , so Theorem 2.6 implies that the number of copies of $K_4 \setminus e$ in G_m is $O(n^4 p^5)$ **wep**.

For (ii), consider the probability that the strategy fails at the $(i + 1)$ -st round for some $i < m$, i.e., that all 3 incoming edges span pairs that complete copies of K_4 . The number of such pairs is upper bounded by the number of copies of $K_4 \setminus e$. Since $G_i \subset G_m$, claim (i) implies that G_i has $O(n^4 p^5)$ copies of $K_4 \setminus e$ **wep**. Call this event A_i , and condition on it. Then, the chance that all 3 incoming edges complete K_4 is $O((\frac{n^4 p^5}{n^2})^3) = O(n^6 p^{15})$. Letting i run through all $m = n^2 p/2$ rounds, a union bound shows that the probability that we are forced to complete a copy of K_4 by the m -th round is $\mathbb{P} \leq O(n^2 p \cdot n^6 p^{15}) + \sum \mathbb{P}[\neg A_i] = O(n^8 p^{16}) + o(1) = o(1)$, as desired. \blacksquare

Proof of upper bound. Let $m \gg n^{3/2}$, and let $p = 2m/n^2$. We will show that **whp**, any strategy fails within $3m$ rounds, which we break into periods of length m . Again, we may assume that $m \ll n^{3/2} \log n$ without loss of generality. Note that $n^{-1/2} \ll p \ll n^{-1/2} \log n$. Our result follows from the following three claims:

- (i) G_m contains $\Omega(n^2)$ pairs of vertices with codegree at least 2 **whp**.
- (ii) G_{2m} contains $\Omega(n^2p)$ copies of $K_4 \setminus e$ **whp**, and with probability $1 - o(n^{-2})$, every pair of vertices in G_{3m} extends $O(1)$ copies of $K_4 \setminus e$ into K_4 .
- (iii) The probability of survival through $3m$ rounds is $o(1)$.

Proof of (i). In the random graph, the expected codegree is roughly $np^2 \gg 2$, but since we do not know how far p exceeds $n^{-1/2}$, we need a slightly more careful argument. Let S be the sum of the codegrees $\sum_{\{u,v\}} d(u, v)$ over all unordered pairs $\{u, v\}$, and let us decompose $S = S_1 + S_2 + S_3$, where S_1 is the contribution from summands with $d(u, v) \in \{0, 1\}$, S_2 is the contribution from summands with $2 \leq d(u, v) \leq 4np^2$, and S_3 is the remainder. We aim to show that $S_2 = \Omega(n^3p^2)$, which will imply the result.

By double-counting, $S = \sum_v \binom{d(v)}{2}$, where $d(v)$ is the degree of vertex v . By convexity, this is always at least $n \binom{d}{2}$, where d is the average degree. Since G_m has exactly m edges, $d = 2m/n = np \gg 1$. Therefore, $S \geq (0.5 + o(1))n(np)^2$.

On the other hand, Lemma 8.1 shows that **whp**, G_m has the property that all codegrees are at most $np^2 \log n$, and for every integer $4 \leq k \leq \log n$, the number of pairs with codegree at least knp^2 is at most n^2/k^3 . Conditioning on this, we may then bound S_3 , the sum of codegrees which exceed $4np^2$, by:

$$\begin{aligned} S_3 &\leq \sum_{k=4}^{\log n} (k+1)np^2 \cdot \frac{n^2}{k^3} \\ &\leq \frac{5}{4} \sum_{k=4}^{\log n} \frac{n^3p^2}{k^2} \\ &\leq n^3p^2 \cdot \frac{5}{4} \left(\frac{\pi^2}{6} - \frac{1}{1^2} - \frac{1}{2^2} - \frac{1}{3^2} \right) \\ &\leq 0.4n^3p^2. \end{aligned}$$

Also, S_1 , the sum of codegrees which are in $\{0, 1\}$, is trivially at most $\binom{n}{2} \ll n^3p^2$ since we assumed $p \gg n^{-1/2}$. So, S_2 , the sum of codegrees between 2 and $4np^2$, is at least $S_2 = S - S_1 - S_3 \geq 0.05n^3p^2$. Therefore, **whp** the number of pairs with codegree at least 2 is at least $0.05n^3p^2 / (4np^2) = \Omega(n^2)$, as claimed. ■

Proof of (ii). The second part follows from Theorem 2.8(ii) because $(K_4 \setminus e, K_4)$ is a balanced extension pair and n^2p^5 is a negative power of n . Let us now concentrate on the first part. Conditioning on the high probability event in claim (i), we may now assume that in G_m , the proportion of pairs with codegree at least 2 is some $\gamma = \Omega(1)$. Consider the $(i+1)$ -st round, where $m \leq i < 2m$. Regardless of the choice of strategy, if all three incoming edges span pairs that each have codegree at least 2, then we will be forced to create a new copy of $K_4 \setminus e$. Incoming edges are uniformly distributed over unoccupied pairs, and the number of occupied pairs in G_i is exactly $i = o(n^2)$. So, since $G_i \supset G_m$, the probability that all three incoming edges span pairs with codegree at least 2 is $q \geq (1 + o(1))\gamma^3 = \Omega(1)$.

Let i run from m to $2m$. Then, the number of copies of $K_4 \setminus e$ in G_{2m} is at least $\text{Bin}(m, q)$. By the Chernoff bound, this exceeds $mq/2 = \Omega(n^2p)$ **wep**, so we are done. ■

Proof of (iii). Consider the $(i+1)$ -st round, where $2m \leq i < 3m$. Regardless of the choice of strategy, if all three incoming edges span pairs that complete copies of K_4 , we will lose. We can lower bound the number of such pairs by $\Omega(n^2p)$ by conditioning on the following

events. Let A be the event that G_{2m} contains $\Omega(n^2p)$ copies of $K_4 \setminus e$, which occurs **whp** by (ii). Also by (ii), with probability $1 - o(n^{-2})$, every pair of vertices in $G_i \subset G_{3m}$ extends $O(1)$ copies of $K_4 \setminus e$ into K_4 ; call this event B_i .

Even after conditioning, the incoming edges in this round are still independently and uniformly distributed over the $\binom{n}{2} - i = \Theta(n^2)$ unoccupied pairs of G_i . Therefore, the probability that both pairs complete K_4 , conditioned on survival through the i -th round, is $p_i = \Omega\left(\left(\frac{n^2p}{n^2}\right)^3\right) = \Omega(p^3)$. Letting i run from $2m$ to $3m$, we see that the probability that any strategy can survive for $3m$ rounds is at most

$$\begin{aligned} \mathbb{P} &\leq \mathbb{P}[\neg A] + \sum \mathbb{P}[\neg B_i] + \prod (1 - p_i) \leq o(1) + \exp\left\{-\sum p_i\right\} \\ &\leq o(1) + \exp\{-\Omega(n^2p \cdot p^3)\} = o(1) + e^{-\omega(1)} = o(1), \end{aligned}$$

which completes the proof. ■

It remains to establish Lemma 8.1, which we used to control the distribution of codegrees in claim (i) of the upper bound.

Proof of Lemma 8.1. Each codegree is distributed as $\text{Bin}(n - 2, p^2)$, so a union bound shows that the probability that some codegree exceeds $np^2 \log n$ is at most

$$\mathbb{P} \leq n^2 \cdot \binom{n}{np^2 \log n} (p^2)^{np^2 \log n} \leq n^2 \cdot \left(\frac{enp^2}{np^2 \log n}\right)^{np^2 \log n} = o(1).$$

Next, fix any $4 \leq k \leq \log n$, and let X be the number of pairs with codegree at least $kn p^2$. Consider an arbitrary vertex v , and let X_v be the number of vertices $u \neq v$ such that the codegree of $\{v, u\}$ is at least $kn p^2$. Note that $X = \frac{1}{2} \sum X_v$.

Since $d(v)$ is binomially distributed $\text{Bin}(n - 1, p)$ and np is a positive power of n , the degree $d(v)$ is at most $1.1np$ **whp** by Chernoff. Condition on this, and condition further on a neighborhood $N(v)$ of size $d(v)$. For each $w \notin N(v) \cup \{v\}$, define the indicator random variable I_w to be 1 if and only if the codegree of $\{v, w\}$ is at least $kn p^2$, or equivalently, if w has at least $kn p^2$ neighbors in $N(v)$. Note that because we already fixed $N(v)$, these I_w are independent since they are determined by disjoint sets of edges. Yet $k \geq 4$ and $np^2 \gg 1$, so each I_w has probability

$$q = \mathbb{P}[I_w] \leq \binom{1.1np}{kn p^2} p^{kn p^2} \leq \left(\frac{1.1enp^2}{kn p^2}\right)^{kn p^2} \leq \left(\frac{3}{k}\right)^{kn p^2} \ll \frac{1}{k^3}.$$

Therefore, X_v is stochastically dominated by $d(v) + \text{Bin}(n - 1 - d(v), q)$. Since $k \leq \log n$, a Chernoff bound implies that **whp**, $X_v \leq 1.1np + 2nq = o(n/k^3)$, which gives $X = \frac{1}{2} \sum X_v = o(n^2/k^3)$. The result follows by taking a union bound over all v and $4 \leq k \leq \log n$. ■

9. CONCLUDING REMARKS

- Although our theorems treat specific graphs (cycles, cliques, and complete bipartite graphs), we conjecture that the thresholds for avoiding general graphs H follow from the natural generalization of the recipe that we used.

To apply our machinery from Section 5, the first thing that we needed to specify was the parameter s . This was the number of levels of danger considered by the avoidance

strategy in the proof of the lower bound. The correct choice of s then determined θ , the negative exponent in the threshold (in terms of p) for avoidance:

$$\theta(H, r, s) = \frac{r^s(v(H) - 2) + 2}{r^s(e(H) - s) + \frac{r^s - 1}{r - 1}}.$$

Furthermore, it is clear that the threshold for avoiding any fixed subgraph $H' \subset H$ is a lower bound for the threshold for avoiding H itself. This is because any strategy that avoids H' will certainly avoid H as well.

In light of this, we conjecture that the threshold for avoiding H in the Achlioptas process with parameter r is $n^{2-\theta^*}$, where θ^* is the minimum value of $\theta(H', r, s)$ when s runs over all nonnegative integers and H' runs over all subgraphs of H .

- Just as in the case of analyzing the Achlioptas process for giant component avoidance [6], one can also consider the offline version of the fixed subgraph avoidance problem. In this offline version, all random r -tuples of edges arriving during the process are accessible to an algorithm, and it can make its choices at each round, relying on the perfect knowledge of the past and the future. The question is still how long the algorithm can typically avoid the appearance of a copy of a fixed graph H . We expect that in most of the cases there will be a sizable difference between the online and the offline thresholds. Here is a sketch of the illustrative case of $H = K_3$, $r = 2$. For this case we can prove that if $m = o(n^{4/3})$, then one can **whp** avoid a copy of K_3 during the first m rounds in the offline version. This should be compared to the threshold of $m = n^{6/5}$ for the online version, given by Theorem 1.1. The argument proceeds as follows. Set $p = 2m/n^2$. The offline model in this case can be approximated quite accurately by generating a random graph G according to the distribution $G(n, 2m)$, and then splitting the edges of G randomly into m pairs: $(e_1, f_1), \dots, (e_m, f_m)$. Denote the above random matching of $E(G)$ by π . We use the following strategy, while processing the pairs (e_i, f_i) : in each pair (e_i, f_i) choose an arbitrary edge not participating in any triangle in G , otherwise pick an arbitrary edge. It is obvious that using this strategy we can only lose (i.e. create a triangle) if G contains a triangle with edges x_1, x_2, x_3 such that their respective pairings in π also belong to triangles in G . The number of triangles in G is **whp** of order $n^3 p^3$, and therefore the probability of having a triangle whose three edges are paired in π with edges from triangles is at most of order

$$n^3 p^3 \left(\frac{n^3 p^3}{n^2 p} \right)^3 = n^6 p^9 = o(1).$$

It would be very interesting to obtain tight results for the offline small subgraph avoidance version of the Achlioptas process for a wide variety of graphs H and parameter r .

- The appearances of the giant component and of a fixed graph are just two instances that have been addressed so far in the context of the Achlioptas process. Naturally, one can consider other graph theoretic properties as well in this context. We hope to return to questions of this type in the future.

APPENDIX A: SUPPORTING RESULTS FOR AVOIDING K_t

In this section, we collect the supporting lemmas and inequalities that are used to prove thresholds for avoiding K_t . Following a suggestion of the referee to shorten this paper, we do not provide complete proofs of all of these results. Rather, we stop once each statement has been reduced to an inequality in several variables. At that point, the remaining analysis is not so interesting, because such statements can of course in theory be verified (although efficient proofs of non-polynomial inequalities in up to eight variables are not necessarily routine). The interested reader can find the complete proofs on the arXiv at <http://arxiv.org/abs/0708.0443>.

Throughout the appendix, we will set $s = \lfloor \log_r[(r-1)t + 1] \rfloor$. We begin by proving some basic facts about s .

Lemma A.1. *For fixed $t \geq 3$, the parameter s is decreasing in r in the range $r \geq 2$.*

Proof. This follows by routine calculus, as it is not difficult to show that $\frac{\partial f}{\partial r} < 0$. ■

Lemma A.2. *If $t \geq 4$ and $r \geq 2$, then $s \leq t/2$. Furthermore, if $t \geq 5$ and $r \geq 2$, or if $t = 4$ and $r \geq 4$, then $s < t/2$.*

Proof. By Lemma A.1, if $r \geq 2$, then $s \leq \lfloor \log_2(t+1) \rfloor$, and one may verify that this is in turn $\leq t/2$ for all $t \geq 4$, and $< t/2$ for $t \geq 5$. For the other range, when $r \geq 4$, Lemma A.1 gives $s \leq \lfloor \log_4(3t+1) \rfloor$, which is less than $t/2$ at $t = 4$. This finishes the lemma. ■

1.1. Balanced Graphs and Extensions

Lemma A.3. *For any $t \geq 4$ and $r \geq 2$, $K_t \setminus se$ is a balanced graph.*

Proof. We must show that the edge density (number of edges divided by number of vertices) of $K_t \setminus se$ is at least as large as the edge density of any of its proper induced subgraphs. The edge density of $K_t \setminus se$ is exactly $[\binom{t}{2} - s]/t$. Lemma A.2 established that $s \leq t/2$, so the edge density is at least $[\binom{t}{2} - \frac{t}{2}]/t = \binom{t-1}{2}/(t-1)$. Yet the final quantity is precisely the edge density of K_{t-1} , which is an upper bound on the edge density of any proper induced subgraph of any t -vertex graph, so we are done. ■

Lemma A.4. *For any $t \geq 4$ and $r \geq 2$, $K_t \setminus se$ has the balanced extension property.*

Proof. Fix any graph G of the form $K_t \setminus se$, and let u, v be any two nonadjacent vertices of G . We must show that the function $e(H)/(v(H)-2)$ is maximal at $H = G$, where H is allowed to range over all induced subgraphs of G that contain $\{u, v\}$. For any graph H with n vertices that is missing at least one edge (e.g., the edge between $\{u, v\}$), $e(H)/(v(H)-2) \leq [\binom{n}{2} - 1]/(n-2) = (n+1)/2$. For any proper induced subgraph $H \subset G$, we then have $e(H)/(v(H)-2) \leq t/2$.

Yet $e(G)/(v(G)-2) = [\binom{t}{2} - s]/(t-2)$, and Lemma A.2 established that $s \leq t/2$. Using this bound for s , we see that $e(G)/(v(G)-2) \geq [\binom{t}{2} - \frac{t}{2}]/(t-2) = t/2$, which matches our upper bound for $e(H)/(v(H)-2)$, so we are done. ■

Lemma A.5. *Suppose that $t \geq 4$. Let $H_1 = K_{\lfloor \frac{t}{2} \rfloor, \lceil \frac{t}{2} \rceil}$, and arbitrarily choose the rest of the sequence $\{H_2, H_3, \dots, H_f\}$, where $H_f = K_t$, by adding one edge at a time. Then every consecutive pair (H_k, H_{k+1}) is a balanced extension pair.*

Proof. Consider a consecutive pair (H_k, H_{k+1}) . By the construction, H_k contains a complete bipartite subgraph that was H_1 ; let $V_1 \cup V_2$ be the corresponding partition of the vertex set. Let u and v be the endpoints of the edge on which H_k and H_{k+1} differ. Without loss of generality, suppose that $u, v \in V_1$. (They must lie in the same part because H_k already contains all edges between V_1 and V_2 .) Now, consider any subsets $U_1 \subset V_1$ and $U_2 \subset V_2$ such that $u, v \in U_1$ and $U_1 \cup U_2 \neq V_1 \cup V_2$. Let H'_k be the subgraph of H_k induced by $U_1 \cup U_2$. It suffices to show that $e(H_k)/(v(H_k) - 2) \geq e(H'_k)/(v(H'_k) - 2)$.

Let us denote $u_1 = |U_1|$, $u_2 = |U_2|$, and let e_1 and e_2 be the respective numbers of edges of H_k spanned by U_1 and by U_2 . Since the number of edges between V_1 and V_2 is $\lfloor \frac{t}{2} \rfloor \lceil \frac{t}{2} \rceil = \lfloor \frac{t^2}{4} \rfloor$, the number of edges in H_k is at least $e_1 + e_2 + \lfloor \frac{t^2}{4} \rfloor$. On the other hand, the number of edges in H'_k is precisely $e_1 + e_2 + u_1 u_2$. Thus, the result follows from the inequality below (proved in the full version):

$$\frac{e_1 + e_2 + \lfloor \frac{t^2}{4} \rfloor}{t - 2} \geq \frac{e_1 + e_2 + u_1 u_2}{u_1 + u_2 - 2}.$$

■

A.2. Inequalities

For the reader's convenience, we reproduce the definitions of the parameters s and θ :

$$s = \lfloor \log_r[(r - 1)t + 1] \rfloor, \quad \theta = \frac{r^s(t - 2) + 2}{r^s \binom{t}{2} - s} + \frac{r^s - 1}{r - 1}.$$

The following inequalities are proved in the full version of this paper.

Inequality A.6. *Suppose that either $t \geq 5$ and $r \geq 2$, or $t = 4$ and $r \geq 4$. Then $-\theta \geq -\lfloor \frac{t}{2} \rfloor^{-1}$.*

Inequality A.7. *For any $t \geq 4$ and $r \geq 2$, if $p \gg n^{-\theta}$, then $n^{t-2} p^{\binom{t}{2} - s - 1}$ is a positive power of n .*

Inequality A.8. *Suppose that $t \geq 5$ and $r \geq 2$, or $t = 4$ and $r \geq 4$. If $p \ll n^{-\theta}$, then $n^{t-2} p^{\binom{t}{2} - s}$ is a negative power of n .*

APPENDIX B: SUPPORTING RESULTS FOR AVOIDING $K_{t,t}$

Coincidentally, the definition of the parameter s is exactly the same for avoiding K_t and avoiding $K_{t,t}$, so we can still use Lemmas A.1 and A.2 (which prove properties of s) in this section. The specification of θ will be different, however. For the reader's convenience, we reproduce the definitions here.

$$s = \lfloor \log_r[(r - 1)t + 1] \rfloor, \quad \theta = \frac{r^s(2t - 2) + 2}{r^s(t^2 - s) + \frac{r^s - 1}{r - 1}}.$$

B.1. Balanced Graphs

Lemma B.1. *For any $t \geq 3$ and $r \geq 2$, $K_{t,t} \setminus se$ is a balanced graph, except in the case when $t = 3$, $r = 2$, and the graph is $K_{2,3}$ with a pendant edge. In that final case, if $p \gg n^{-18/31} / \log n$, the number of copies of that graph in G_m is still $O(n^6 p^7)$ **wep**.*

Proof. We must show that the edge density (number of edges divided by number of vertices) of $K_{t,t} \setminus se$ is at least the edge density of any proper induced subgraph. The edge density of the complete bipartite graph $K_{a,b}$ is $ab/(a+b)$, which is increasing in both a and b , so the edge density of any proper induced subgraph of $K_{t,t} \setminus se$ is at most $t(t-1)/(2t-1)$. On the other hand, the edge density of $K_{t,t} \setminus se$ is precisely $(t^2 - s)/(2t)$, so we must show that

$$\frac{t(t-1)}{2t-1} \leq \frac{t^2 - s}{2t}.$$

Clearing the denominators, this is equivalent to

$$2t^3 - 2t^2 \leq 2t^3 - t^2 - s(2t - 1).$$

Rearranging terms, this is equivalent to

$$s \leq \frac{t^2}{2t - 1}.$$

Now if $t \geq 4$, Lemma A.2 bounds $s \leq t/2$, which finishes the inequality.

The only remaining case is $t = 3$. However, Lemma A.1 established that the dependence of $s = \lfloor \log_r[(r-1)t + 1] \rfloor$ on r was decreasing, so $s = 1$ for $r \geq 3$, and $s = 2$ for $r = 2$. One may manually verify that of all of the graphs of the form $K_{3,3} \setminus e$ and $K_{3,3} \setminus 2e$, the only one which is not balanced is the deletion from $K_{3,3}$ of two edges incident to the same vertex, which is $K_{2,3}$ with a pendant edge, as claimed. Since that graph, which we denote $K_{2,3} + e$, arises only when $s = 2$, this happens only when $r = 2$.

Now let us bound the number of copies of that graph in $G(n, p)$, when $p \gg n^{-\theta} / \log n$. In the case $t = 3, r = 2$, we have $\theta = -\frac{18}{31}$, and so $n^5 p^6$, roughly the expected number of copies of $K_{2,3}$ in the random graph, is a positive power of n . So, since $K_{2,3}$ is balanced, Theorem 2.6 bounds the number of copies of $K_{2,3}$ in G_m by $O(n^5 p^6)$ **wep**. Also, np is a positive power of n , so we may bound all degrees by $2np$ **wep**. If both situations hold, we may conclude that the number of copies of $K_{2,3} + e$ is $O(n^5 p^6 \cdot np) = O(n^6 p^7)$, as desired. ■

Lemma B.2. *For any $t \geq 3$ and $r \geq 2$, $K_{t,t} \setminus se$ has the balanced extension property.*

Proof. Fix any graph G of the form $K_{t,t} \setminus se$, and let u, v be any two nonadjacent vertices of G . We must show that the function $e(H)/(v(H) - 2)$ is maximized at $H = G$, where H is allowed to range over all proper induced subgraphs of G that contain $\{u, v\}$. Any such H is still bipartite with respect to G 's bipartition; suppose that it has a vertices on one side and b on the other. Since we assumed that H is missing at least the edge joining $\{u, v\}$, we must have $e(H)/(v(H) - 2) \leq (ab - 1)/(a + b - 2)$. This is increasing in both a and b , so its maximum over proper induced subgraphs H is $\lfloor t(t-1) - 1 \rfloor / (2t - 3)$. Thus, the result follows from the inequality below (proved in the full version):

$$\frac{t^2 - s}{2t - 2} \geq \frac{t(t-1) - 1}{2t - 3}. \quad \blacksquare$$

Lemma B.3. *For any fixed positive integer k , consider the following 4-partite graph, which we call H_1 . Let the parts be V_1, V_2, V_3, V_4 , with $|V_1| = |V_2| = k$ and $|V_3| = |V_4| = 1$, and place edges such that (V_1, V_2) , (V_1, V_4) , and (V_3, V_2) are complete bipartite. There are no more edges. Let H_2 be obtained from H_1 by adding the edge between V_3 and V_4 . Then (H_1, H_2) is a balanced extension pair.*

Proof. Consider any subsets $U_1 \subset V_1$ and $U_2 \subset V_2$, and let H'_1 be the subgraph of H_1 induced by $U_1 \cup U_2 \cup V_3 \cup V_4$. We must show that $e(H'_1)/(v(H'_1) - 2) \leq e(H_1)/(v(H_1) - 2)$. Let $a = |U_1|$ and $b = |U_2|$. Then, $\frac{e(H'_1)}{v(H'_1)-2} = \frac{ab+a+b}{a+b} = \frac{ab}{a+b} + 1$, which is increasing in both a and b . Therefore, $\frac{e(H'_1)}{v(H'_1)-2} \leq \frac{k^2+k+k}{k+k} = \frac{e(H_1)}{v(H_1)-2}$, and we are done. ■

Lemma B.4. *Suppose that t is even and at least 4. Let H_1 be the 4-partite graph with parts V_1, V_2, V_3, V_4 , each of size $t/2$, and edges such that (V_1, V_2) , (V_1, V_4) , and (V_3, V_2) are complete bipartite. Let $\{H_2, \dots, H_{1+(t/2)}\}$ be obtained by successively adding single edges until $H_{1+(t/2)}$ has a perfect matching between V_3 and V_4 . Then, arbitrarily choose the rest of the sequence $\{H_{2+(t/2)}, \dots, H_f\}$ by adding one edge at a time, until the final term is the complete bipartite graph $K_{t,t}$ with bipartition $(V_1 \cup V_3, V_2 \cup V_4)$. Then every consecutive pair (H_k, H_{k+1}) is a balanced extension pair.*

The proof breaks into two cases, since there are two stages of edge addition. To give a flavor of the argument, we show how to reduce one of the cases to an inequality in several variables.

Proof of Lemma B.4 for $k \leq t/2$. Consider a consecutive pair (H_k, H_{k+1}) . By the construction, H_k has the following structure. The vertex set is partitioned into $V_1 \cup V_2 \cup V_3 \cup V_4$, with all parts of size $t/2$. The pairs (V_1, V_2) , (V_1, V_4) , and (V_3, V_2) are complete bipartite graphs, and there is a $(k - 1)$ -edge matching between V_3 and V_4 . There are no other edges. Also, there is a pair of vertices $u \in V_3, v \in V_4$, not involved in the $(k - 1)$ -edge matching, at which the addition of an edge creates H_{k+1} . Now consider any family of subsets $U_i \subset V_i$ such that $u \in U_3$ and $v \in U_4$. Let H'_k be the subgraph of H_k induced by $\cup U_i$. We must show that $e(H'_k)/(v(H'_k) - 2) \leq e(H_k)/(v(H_k) - 2)$.

For brevity, let $a = |U_1|, b = |U_2|, c = |U_3|$, and $d = |U_4|$. Since the edges between U_3 and U_4 form a matching of at most $k - 1$ edges which does not involve $u \in U_3$ or $v \in U_4$, there can be at most $\min\{c - 1, d - 1, k - 1\} = \min\{c, d, k\} - 1$ edges there. Therefore,

$$\frac{e(H'_k)}{v(H'_k) - 2} \leq \frac{ab + ad + cb + (\min\{c, d, k\} - 1)}{a + b + c + d - 2}.$$

The result follows by showing that the right hand side is at most $\frac{\frac{3}{4}t^2+(k-1)}{2t-2} = \frac{e(H_k)}{v(H_k)-2}$, which is done in the full version of this article. ■

Lemma B.5. *Suppose that t is odd and at least 3. Let H_1 be a 6-partite graph with parts $\{V_i\}_1^6$ such that V_3 and V_4 are singletons, and the other four parts each have size $\lfloor t/2 \rfloor$. Let there be edges be such that the two pairs (V_1, V_2) and (V_5, V_6) are each complete bipartite graphs, let the vertex in V_3 be adjacent to all of $V_2 \cup V_4 \cup V_6$, and let the vertex in V_4 be adjacent to all of $V_1 \cup V_3 \cup V_5$. There are no more edges.*

Let $\{H_2, \dots, H_{1+\lfloor t/2 \rfloor}\}$ be obtained by successively adding single edges until $H_{1+\lfloor t/2 \rfloor}$ has a perfect matching between V_1 and V_6 . To create the next $\lfloor t/2 \rfloor$ graphs in the sequence, we put down a matching between V_5 and V_2 , one edge at a time. Finally, arbitrarily choose the rest of the sequence $\{H_{2+2\lfloor t/2 \rfloor}, \dots, H_f\}$ by adding one edge at a time, until the final term is the complete bipartite graph $K_{t,t}$ with bipartition $(V_1 \cup V_3 \cup V_5, V_2 \cup V_4 \cup V_6)$.

Then every consecutive pair (H_k, H_{k+1}) is a balanced extension pair.

The proof breaks into three cases, since there are three stages of edge addition. To give a flavor of the argument, we show how to reduce one of the cases to an inequality in several variables.

Proof of Lemma B.5 for $k > 2\lfloor t/2 \rfloor$. Consider a consecutive pair (H_k, H_{k+1}) . By construction, H_k has the following structure. The vertex set is partitioned into $\{V_i\}_1^6$, with $|V_3| = |V_4| = 1$ and all other $|V_i| = \lfloor t/2 \rfloor$. The pairs (V_1, V_2) and (V_5, V_6) are each complete bipartite graphs, the vertex in V_3 is adjacent to all of $V_2 \cup V_4 \cup V_6$, the vertex in V_4 is adjacent to all of $V_1 \cup V_3 \cup V_5$, there is a perfect $\lfloor t/2 \rfloor$ -edge matching between V_1 and V_6 , and another perfect matching between V_5 and V_2 . There may be some more edges as well between V_1 and V_6 or between V_5 and V_2 , but not all such edges are present: without loss of generality, let us suppose that there are two vertices $u \in V_1$ and $v \in V_6$ such that there is no edge between u and v . There are no more edges in the entire graph. Also, H_{k+1} is obtained from H_k by adding the edge joining u and v . Now, consider any family of subsets $U_i \subset V_i$ such that $u \in U_1$ and $v \in U_6$. Let H'_k be the subgraph of H_k induced by $\cup U_i$. We must show that $e(H'_k)/(v(H'_k) - 2) \leq e(H_k)/(v(H_k) - 2)$.

For brevity, let $a = |U_1|$, $b = |U_2|$, $c = |U_3|$, $d = |U_4|$, $e = |U_5|$, and $f = |U_6|$. Let E be the number of edges in H_k between U_1 and U_6 or between U_5 and U_2 . Then

$$\frac{e(H'_k)}{v(H'_k) - 2} = \frac{ab + ef + c(b + f) + (a + e)d + cd + E}{a + b + c + d + e + f - 2}. \tag{1}$$

Next, recall that H_k contained a perfect $\lfloor t/2 \rfloor$ -edge matching between V_1 and V_6 , and between V_5 and V_2 . The maximum number of edges of these matchings that are included in E (i.e., go between U_1 and U_6 , or between U_5 and U_2) is $\min\{a, f\} + \min\{b, e\} \leq (a + f + b + e)/2$. Therefore, the number of edges in H_k between V_1 and V_6 or between V_5 and V_2 is at least $E + 2\lfloor \frac{t}{2} \rfloor - \frac{a+b+e+f}{2}$. The rest of the edges in H_k are easy to count: (V_1, V_2) and (V_5, V_6) are complete bipartite subgraphs $K_{\lfloor t/2 \rfloor, \lfloor t/2 \rfloor}$, the vertex in V_3 is adjacent to all of $V_2 \cup V_4 \cup V_6$, and the vertex in V_4 is adjacent to all of $V_1 \cup V_3 \cup V_5$. Therefore,

$$\frac{e(H_k)}{v(H_k) - 2} \geq \frac{2\lfloor \frac{t}{2} \rfloor^2 + [4\lfloor \frac{t}{2} \rfloor + 1] + [E + 2\lfloor \frac{t}{2} \rfloor - \frac{a+b+e+f}{2}]}{2t - 2}. \tag{2}$$

The result follows by proving that the right hand side of (1) is at most the right hand side of (2). The full version of this article contains the details. ■

B.2. Inequalities

The following inequalities are proved in the full version of this article.

Inequality B.6. *Suppose that $t \geq 3$ and $r \geq 2$. Then $-\theta > -\frac{2}{r}$.*

Inequality B.7. For any $t \geq 3$ and $r \geq 2$, if $p \gg n^{-\theta}$, then $n^{2t-2}p^{t^2-s-1}$ is a positive power of n .

Inequality B.8. For any $t \geq 3$ and $r \geq 2$, if $p \ll n^{-\theta}$, then $n^{2t-2}p^{t^2-s}$ is a negative power of n .

ACKNOWLEDGMENT

The authors would like to thank the referee for suggestions that improved the exposition of this paper.

REFERENCES

- [1] N. Alon and J. Spencer, The probabilistic method, 2nd edition, Wiley, New York, 2000.
- [2] A. Beveridge, T. Bohman, A. Frieze, and O. Pikhurko, Product rule wins a competitive game, Proc Am Math Soc 135 (2007), 3061–3071.
- [3] A. Beveridge, T. Bohman, A. Frieze, and O. Pikhurko, Memoryless rules for Achlioptas processes, SIAM J Discrete Math, (in press).
- [4] T. Bohman and A. Frieze, Avoiding a giant component, Random Struct Algorithms 19 (2001), 75–85.
- [5] T. Bohman, A. Frieze, and N. Wormald, Avoidance of a giant component in half the edge set of a random graph, Random Struct Algorithms 25 (2004), 432–449.
- [6] T. Bohman and J. Kim, A phase transition for avoiding a giant component, Random Struct Algorithms 28 (2006), 195–214.
- [7] T. Bohman and D. Kravitz, Creating a giant component, Combinatorics Probab Comput 15 (2006), 489–511.
- [8] B. Bollobás, Random graphs, 2nd edition, Cambridge studies in advanced mathematics, Vol. 73, Cambridge University Press, Cambridge, 2001.
- [9] B. Bollobás, Random graphs, In Combinatorics, (Swansea, 1981), London Math. Soc. Lecture Note Ser. Vol. 52, Cambridge University Press, Cambridge, 1981, pp. 80–102.
- [10] B. Bollobás and A. Thomason, Threshold functions, Combinatorica 7 (1987), 35–38.
- [11] P. Erdős and A. Rényi, On the evolution of random graphs, Publ Math Inst Hungar Acad Sci 5 (1960), 17–61.
- [12] P. Erdős and M. Simonovits, Compactness results in extremal graph theory, Combinatorica 2 (1982), 275–288.
- [13] A. Flaxman, D. Garmanik, and G. Sorkin, Embracing the giant component, Random Struct Algorithms 27 (2005), 277–289.
- [14] S. Janson, T. Łuczak, and A. Ruciński, Random graphs, Wiley, New York, 2000.
- [15] T. Kövári, V. Sós, and P. Turán, On a problem of K. Zarankiewicz, Colloquium Math 3 (1954), 50–57.
- [16] J. Spencer and N. Wormald, Birth control for giants, Combinatorica 27 (2007), 587–628.
- [17] V. Vu, Concentration of non-Lipschitz functions and applications, Random Struct Algorithms 20 (2002), 262–316.