

A BOUND FOR THE COPS AND ROBBERS PROBLEM*

ALEX SCOTT[†] AND BENNY SUDAKOV[‡]

Abstract. In this short paper we study the game of cops and robbers, which is played on the vertices of some fixed graph G . Cops and a robber are allowed to move along the edges of G , and the goal of cops is to capture the robber. The cop number $c(G)$ of G is the minimum number of cops required to win the game. Meyniel conjectured a long time ago that $O(\sqrt{n})$ cops are enough for any connected G on n vertices. Improving several previous results, we prove that the cop number of an n -vertex graph is at most $n2^{-(1+o(1))\sqrt{\log n}}$. A similar result independently and slightly before us was also obtained by Lu and Peng.

Key words. pursuit-evasion games, graph searching, cop number, games on graphs

AMS subject classifications. Primary, 05C57; Secondary, 91A24, 91A43

DOI. 10.1137/100812963

1. Introduction. Let G be a simple, undirected, connected graph on n vertices. The game of *Cops and Robbers*, which was introduced almost thirty years by Nowakowski and Winkler [12] and by Quilliot [13], is played on the vertices of G as follows. There are two players: a set of $k \geq 1$ cops and one robber. The game begins by the cops occupying some set of k vertices of G and then the robber also chooses a vertex. Afterward they move alternatively, first cops and then the robber, along the edges of graph G . At every step each cop or robber is allowed to move to any neighboring vertex or do nothing and stay where they are. Multiple cops are allowed to occupy the same vertex. The cops win if at some time, there is a cop at the same vertex as the robber; otherwise, the robber wins. The minimum number of cops for which there is a winning strategy, no matter how robber plays, is called *the cop number of G* and is denoted by $c(G)$. Note that this number is clearly at most n and also that the initial position of cops does not matter, since G is connected.

The cop number was introduced by Aigner and Fromme [1] who proved, for example, that if G is planar, then $c(G) \leq 3$. They also observed that if G has girth at least 5 (i.e., no cycles of length ≤ 4), then its cop number is at least the minimum degree of G . In particular, this together with the well-known construction of dense graphs of girth 5 shows that there are n -vertex graphs which require at least $\Omega(\sqrt{n})$ cops. A quarter century ago, Meyniel conjectured that this is tight and that $O(\sqrt{n})$ cops are always sufficient. The first nontrivial upper bound for this problem was obtained by Frankl [8], who proved that $c(G) \leq O(\frac{n \log \log n}{\log n})$. This was later improved by Chiniforooshan [7] to $O(\frac{n}{\log n})$. All logarithms in this paper are binary.

The game of cops and robbers was also studied by Andreae [4], Berarducci and Intrigila [5], and Alspach [3] and for random graph by Bollobás, Kun, and Leader [6] and by Łuczak and Prałat [11]. The aim of this short paper is to prove the following new bound for the cops and robber problem.

*Received by the editors October 27, 2010; accepted for publication (in revised form) July 1, 2011; published electronically September 20, 2011.

<http://www.siam.org/journals/sidma/25-3/81296.html>

[†]Mathematical Institute, 24-29 St Giles', Oxford, OX1 3LB, UK (scott@maths.ox.ac.uk).

[‡]Department of Mathematics, University of California, Los Angeles, Los Angeles, CA 90095 (bsudakov@math.ucla.edu). This author's research was supported in part by NSF grant DMS-1101185, NSF CAREER award DMS-0812005, and by USA–Israeli BSF grant 2008071.

THEOREM 1.1. *The cop number of any connected n -vertex graph is at most $n2^{-(1+o(1))\sqrt{\log n}}$.*

2. Proof of the main result. The proof of the main theorem has two ingredients. The first is a result of Aigner and Fromme which says that one cop can control the shortest path between two vertices. Let P be the shortest path in G between two vertices u and v . Then the following lemma was proved in [1].

LEMMA 2.1. *One cop can move along the vertices of P such that, after a finite number of steps, if the robber ever visits P , then he will be caught in the next step.*

This result can be used for graphs with large diameter, since we can delete a long path from such a graph and use induction. The case where G has relatively small diameter will be treated using the following key lemma, which we think has independent interest.

LEMMA 2.2. *Let G be a graph on $n \geq 2^{30}$ vertices with diameter $D \leq 2\sqrt{\log n} / \log^3 n$. Then $c(G) \leq n(\log n)^3 2^{-\sqrt{\log n}}$.*

Proof. Throughout the proof, we assume that n is large enough for our estimates to hold. Let $t = \sqrt{\log n} - 3 \log \log n$. Consider random subsets $\mathcal{C}_1, \dots, \mathcal{C}_{t+1}$ of G . For every vertex of G we put it into each \mathcal{C}_j with probability $(\log n)^2 2^{-\sqrt{\log n}}$ (so a single vertex may be in many \mathcal{C}_i). Since $|\mathcal{C}_j|$ is binomially distributed with expectation $\mu = n(\log n)^2 2^{-\sqrt{\log n}}$, by the standard Chernoff-type estimates (see, e.g., Appendix A in [2]), we have that the probability that \mathcal{C}_j has more than $2\mu = 2n(\log n)^2 2^{-\sqrt{\log n}}$ vertices is at most $e^{-\mu/3} < n^{-2}$. Then, with probability at least $1 - \frac{\log n}{n^2} > 0.9$, the sum of the sizes of these sets is at most $n(\log n)^3 2^{-\sqrt{\log n}}$, which will give our final bound.

For every subset A of G and integer i let $B(A, i)$ be the ball of radius i around A , i.e., all the vertices of G which can be reached from some vertex in A by a path of length at most i . We need the following simple claim.

Claim 2.3. The following statement holds with probability 0.9: for every $A \subset V(G)$ such that $|A| \leq n2^{-\sqrt{\log n}}$, every $i \leq t$ such that $|B(A, 2^i)| \geq 2\sqrt{\log n}|A|$, and every j ,

$$|B(A, 2^i) \cap \mathcal{C}_j| \geq |A|.$$

Proof. Let $|A| = a$. Note that for any fixed A, i, j the number of points from \mathcal{C}_j in $B(A, 2^i)$ is binomially distributed with expectation at least $a \log^2 n$. Thus by Chernoff-type estimates, the probability that it is smaller than a is at most $e^{-a \log^2 n/3}$. The number of sets of size a is $\binom{n}{a}$ and the number of pairs of indices i, j is at most $\log n$, and the result follows since $\log n \sum_a \binom{n}{a} e^{-a \log^2 n/3} < 0.1$. \square

Thus we can choose subsets $\mathcal{C}_1, \dots, \mathcal{C}_{t+1}$ that satisfy the assertion of this claim and such that the sum of their sizes is at most $n(\log n)^3 2^{-\sqrt{\log n}}$. We place at each vertex u of G one cop for each set \mathcal{C}_j that contains u . We will show that these cops can always catch the robber.

Suppose that the robber is located at vertex v of the graph G . Since the cops move first, note that the degree of this vertex is at most $2\sqrt{\log n}$. Otherwise, by the claim, we already have a cop in its neighborhood who will catch the robber in the first move. Consider the following sequence of sets A_i, D_i for $i = 1, \dots, t + 1$ defined recursively.

Let A_1 be the largest subset of $B(v, 1)$ such that $|B(A_1, 1)| < 2\sqrt{\log n}|A_1|$, and let $D_1 = B(v, 1) - A_1$. Note that for any $X \subset D_1$ we have that $|B(X, 1)| \geq 2\sqrt{\log n}|X|$ since

otherwise we could increase A_1 by adding X to it. Thus by Claim 2.3 we have that $B(X, 1)$ contains more than $|X|$ points from \mathcal{C}_1 . Consider an auxiliary bipartite graph with vertex classes D_1 and \mathcal{C}_1 , where we join a vertex u in D_1 to w in \mathcal{C}_1 if w is within distance at most one from u . Note that, by the above discussion, it follows from Hall's theorem that this bipartite graph has a complete matching from D_1 to \mathcal{C}_1 . Therefore for every vertex $u \in D_1$ we can choose a corresponding vertex w in \mathcal{C}_1 , which is within distance one from u and such that all w are distinct. We can therefore ask the cops in \mathcal{C}_1 to occupy D_1 in the first round.

Now suppose we have already defined A_i, D_i . Consider the ball $B(A_i, 2^{i-1})$. Let A_{i+1} be the largest subset of $B(A_i, 2^{i-1})$ such that $|B(A_{i+1}, 2^i)| < 2^{\sqrt{\log n}}|A_{i+1}|$, and let $D_{i+1} = B(A_i, 2^{i-1}) - A_{i+1}$. Again for any subset $X \subset D_{i+1}$ we have that $|B(X, 2^i)| \geq 2^{\sqrt{\log n}}|X|$. Thus by Claim 2.3 we have that $B(X, 2^i)$ contains more than $|X|$ points from \mathcal{C}_{i+1} . As above, we consider an auxiliary bipartite graph with parts D_{i+1} and \mathcal{C}_{i+1} such that a vertex u in D_{i+1} is joined to w in \mathcal{C}_{i+1} if w is within distance at most 2^i from u . The bipartite graph has a complete matching from D_{i+1} to \mathcal{C}_{i+1} , and so for every vertex $u \in D_{i+1}$ we can choose a corresponding vertex w in \mathcal{C}_{i+1} , which is within distance at most 2^i from u and such that all w are distinct. We will ask the cops in \mathcal{C}_{i+1} to occupy D_{i+1} by the end of the first 2^i rounds (by rounds we mean here that cops will move first, then the robber moves, and so on until both the cops and robber have each made 2^i moves).

Next we claim that there is an index $s \leq t + 1$ such that A_s is empty. Indeed note that by construction we have $|A_{i+1}| \leq |B(A_i, 2^{i-1})| \leq 2^{\sqrt{\log n}}|A_i|$. Therefore if A_{t+1} is not empty, we have that

$$|A_{t+1}| \leq (2^{\sqrt{\log n}})^{t+1} \leq n2^{-2\sqrt{\log n}}.$$

Moreover, we also have that $B(A_{t+1}, 2^t)$ has size at most $2^{\sqrt{\log n}}|A_{t+1}| \ll n$, and this contradicts the assumption that the diameter of G is at most $2^{\sqrt{\log n}}/\log^3 n$.

Let $s \leq t + 1$ be such that A_s is empty. We have shown that we can move cops from the sets \mathcal{C}_i , $1 \leq i \leq s$, such that after 2^{i-1} rounds they will occupy the set D_i . Also we have that $B(A_{s-1}, 2^{s-2}) = D_s$. Now we claim that after 2^{s-1} rounds we have already caught the robber.

After one round the robber has made one step, so he is in $B(v, 1)$. Note that D_1 is already occupied by cops after the first round, so in order to survive the robber must be in $A_1 = B(v, 1) - D_1$. Suppose by induction that after 2^{i-1} rounds the robber is in the set A_i . Consider the next 2^{i-1} rounds (so in total $2 \cdot 2^{i-1} = 2^i$ rounds). After this many additional rounds, the robber is in $B(A_i, 2^{i-1})$. Since we know that D_{i+1} is occupied by the cops by time 2^i , the robber must be in $B(A_i, 2^{i-1}) - D_{i+1} = A_{i+1}$. Thus, arguing by induction, the only place the robber can be after 2^{i-1} rounds without having been caught is in the set A_i . Since A_s is empty, we are done. This completes the proof of the lemma. \square

Proof of Theorem 1.1. Let G be a connected graph on n vertices. We prove by induction that $c(G) \leq f(n) = 2n(\log n)^3 2^{-\sqrt{\log n}}$. Since $c(G) \leq n$, the result holds trivially when $2(\log n)^3 2^{-\sqrt{\log n}} \geq 1$. Thus we can assume that $\log n \geq 400$. If the diameter of G is at most $2^{\sqrt{\log n}}/\log^3 n$, then we are immediately done by Lemma 2.2. Otherwise G contains two vertices such that the shortest path P between them has length at least $D = 2^{\sqrt{\log n}}/\log^3 n$. Put one cop on this path. By Lemma 2.1, after a finite number of

steps this cop can prevent the robber from entering path P . Thus we can continue the game on the graph obtained from G by deleting P . If this graph is disconnected, we continue playing on the connected component which contains the robber. Thus $c(G) \leq 1 + c(G - P)$. By induction we know that

$$(1) \quad c(G - P) \leq f(n - D) \leq 2(n - D)(\log n)^3 2^{-\sqrt{\log(n-D)}} \leq 2n(\log n)^3 2^{-\sqrt{\log(n-D)}} - 2.$$

Since $\log n \geq 400$, we have $D \leq 2^{-20}n$. Then $\log(n - D) \geq \log n - 2D/n$, and therefore

$$\sqrt{\log(n - D)} \geq \sqrt{\log n - 2D/n} \geq \sqrt{\log n} - 2D / \left(n\sqrt{\log n} \right).$$

We also have that $2^{2D/(n\sqrt{\log n})} \leq (1 + 2D/(n\sqrt{\log n}))$. Substituting this into (1) we get that

$$\begin{aligned} c(G - P) &\leq 2n(\log n)^3 2^{-\sqrt{\log(n-D)}} - 2 \leq 2n(\log n)^3 2^{-\sqrt{\log n - 2D/(n\sqrt{\log n})}} - 2 \\ &\leq 2n(\log n)^3 2^{-\sqrt{\log n}} \left(1 + 2D / \left(n\sqrt{\log n} \right) \right) - 2 \\ &= f(n) + \frac{4}{\sqrt{\log n}} - 2 < f(n) - 1. \end{aligned}$$

Therefore $c(G) \leq 1 + c(G - P) < 1 + f(n) - 1 = f(n)$, which completes the induction step. \square

Finally, let us note that the strategy in Lemma 2.2 does not require the cops to see the robber's movements after the first round. The cops therefore have the following randomized strategy against an invisible robber (see [9] for a discussion of cops and robbers games where there is a cop with limited vision). The cops move to their assigned positions, then guess the position of the robber and follow the strategy of Lemma 2.2, repeating this process until the robber is caught. This algorithm catches the robber in polynomial expected time. On a general graph, we can add a further cop on each path P that we have deleted in the proof of Theorem 1.1: If these cops move at random on their paths, the robber will be caught in finite (although not necessarily polynomial) expected time.

Note added in proof. When this paper was written, we learned that a similar result independently and slightly before us was also obtained by Lu and Peng [10]. We would like to thank Krivelevich for bringing their paper to our attention.

REFERENCES

- [1] M. AIGNER AND M. FROMME, *A game of cops and robbers*, Discrete Appl. Math., 8 (1984), pp. 1–11.
- [2] N. ALON AND J. SPENCER, *The Probabilistic Method*, 3rd ed., Wiley, Hoboken, NJ, 2008.
- [3] B. ALSPACH, *Searching and sweeping graphs: A brief survey*, Matematiche (Catania), 59 (2004), pp. 5–37.
- [4] T. ANDREA, *On a pursuit game played on graphs for which a minor is excluded*, J. Combin. Theory Ser. B, 41 (1986), pp. 37–47.
- [5] A. BERARDUCCI AND B. INTRIGILA, *On the cop number of a graph*, Adv. in Appl. Math., 14 (1993), pp. 389–403.
- [6] B. BOLLOBÁS, G. KUN, AND I. LEADER, *Cops and Robbers in Random Graphs*, ArXiv 0805.2709, 2008.
- [7] E. CHINIFOROOSHAN, *A better bound for the cop number of general graphs*, J. Graph Theory, 58 (2008), pp. 45–48.

- [8] P. FRANKL, *Cops and robbers in graphs with large girth and Cayley graphs*, Discrete Appl. Math., 17 (1987), pp. 301–305.
- [9] V. ISLER AND N. KARNAD, *The role of information in the cop-robber game*, Theoret. Comput. Sci., 399 (2008), pp. 179–190.
- [10] L. LU AND X. PENG, *On Meyniel’s conjecture of the cop number*, J. Graph Theory, to appear.
- [11] T. ŁUCZAK AND P. PRALAT, *Chasing robbers on random graphs: Zigzag theorem*, Random Structures Algorithms, 37 (2010), pp. 516–524.
- [12] R. NOWAKOWSKI AND P. WINKLER, *Vertex-to-vertex pursuit in a graph*, Discrete Math., 43 (1983), pp. 235–239.
- [13] A. QUILLIOT, *Jeux et pointes fixes sur les graphes*, Ph.D. dissertation, Université de Paris VI, Paris, 1978.