

On Graphs with Subgraphs Having Large Independence Numbers

Noga Alon¹ and Benny Sudakov^{2,3}

¹SCHOOLS OF MATHEMATICS AND COMPUTER SCIENCE
RAYMOND AND BEVERLY SACKLER FACULTY OF EXACT SCIENCES
TEL AVIV UNIVERSITY
TEL AVIV 69978, ISRAEL
E-mail: nogaa@tau.ac.il

²DEPARTMENT OF MATHEMATICS
PRINCETON UNIVERSITY, PRINCETON
NEW JERSEY 08544
E-mail: bsudakov@math.princeton.edu

³INSTITUTE FOR ADVANCED STUDY PRINCETON
NEW JERSEY

Received August 8, 2006; Revised July 9, 2007

Published online 9 August 2007 in Wiley InterScience(www.interscience.wiley.com).
DOI 10.1002/jgt.20264

Abstract: Let G be a graph on n vertices in which every induced subgraph on $s = \log^3 n$ vertices has an independent set of size at least $t = \log n$. What is the largest $q = q(n)$ so that every such G must contain an independent set of size at least q ? This is one of the several related questions raised by Erdős and Hajnal. We show that $q(n) = \Theta(\log^2 n / \log \log n)$, investigate

Contract grant sponsor: Israel Science Foundation (to N.A.); Contract grant sponsor: USA-Israeli BSF grant (to N.A.); Contract grant sponsor: Hermann Minkowski Minerva Center for Geometry (to N.A.); Contract grant sponsor: NSF CAREER award (to B.S.); Contract grant number: DMS-0546523; Contract grant sponsor: NSF (to B.S.); Contract grant number: DMS-0355497; Contract grant sponsor: Alfred P. Sloan fellowship; Contract grant sponsor: State of New Jersey.

Journal of Graph Theory
© 2007 Wiley Periodicals, Inc.



the more general problem obtained by changing the parameters s and t , and discuss the connection to a related Ramsey-type problem. © 2007 Wiley Periodicals, Inc. *J Graph Theory* 56: 149–157, 2007

Keywords: *Ramsey-type problems; independence numbers*

1. INTRODUCTION

What is the largest $f = f(n)$ so that every graph G on n vertices in which every induced subgraph on $\log^2 n$ vertices has an independent set of size at least $\log n$, must contain an independent set of size at least f ? This is one of the several related questions considered by Erdős and Hajnal in the late 80s. The question appears in [3], where Erdős mentions that they thought that $f(n)$ must be at least $n^{1/2-\epsilon}$, but they could not even prove that it is at least $2 \log n$. As a special case of our main results, here, we determine the asymptotic behavior of $f(n)$ up to a factor of $\log \log n$, showing that in fact it is much smaller than one may suspect (and yet much bigger than $\log n$):

$$\Omega\left(\frac{\log^2 n}{\log \log n}\right) \leq f(n) \leq O(\log^2 n). \quad (1)$$

Another specific variant of the above question, discussed in [3], is the problem of estimating the largest $q = q(n)$ so that every graph G on n vertices in which every induced subgraph on $\log^3 n$ vertices has an independent set of size at least $\log n$, must contain an independent set of size at least q . Here, too, one may tend to believe that $q(n)$ is large, and specifically it is mentioned in [3] that probably $q(n) > \log^3 n$, but the correct asymptotic behavior of $q(n)$ is smaller. In this case, our results determine the asymptotic behavior of $q(n)$ up to a constant factor, implying that

$$q(n) = \Theta\left(\frac{\log^2 n}{\log \log n}\right). \quad (2)$$

Both problems above are special instances of the general problem of understanding the asymptotic behavior of the function $f(n, s, t)$ defined as follows. For $n > s > t$, let $f = f(n, s, t)$ denote the largest integer f so that every graph G on n vertices in which every induced subgraph on s vertices has an independent set of size at least t , must contain an independent set of size at least f . In this note, we investigate the asymptotic behavior of f , and obtain rather tight bounds for this behavior for most interesting values of the parameters. Our results provide much less satisfactory information about a closely related Ramsey-type problem of Erdős and Hajnal discussed in [3], which is the following. For which functions

$h(n)$ and $g(n)$, where $n > g(n) \geq h(n)^2 \gg 1$, is there a graph on n vertices in which every induced subgraph on $g(n)$ vertices contains a clique of size $h(n)$ as well as an independent set of size $h(n)$? In particular, Erdős and Hajnal conjectured that there is no such graph for $g(n) = \log^3 n$ and $h(n) = \log n$; our results here do not settle this conjecture, and only suffice to show that there is no such graph with $g(n) = c \log^3 n / \log \log n$ and $h(n) = \log n$, for some absolute positive constant c .

The rest of this note is organized as follows. In Section 2, we state our main results concerning the behavior of the function $f(n, s, t)$. The proofs are described in Section 3. The final Section 4 contains a few remarks, including the (simple) connection between the study of f and the Ramsey-type question discussed above.

Throughout the note, we omit all floor and ceiling signs, whenever these are not crucial. We always assume that the number n of vertices of the graphs considered here is large. All logarithms are in the natural base e .

2. THE MAIN RESULTS

The following two theorems provide lower bounds for the independence numbers of graphs in which every induced subgraph of size s contains an independent set of size t .

Theorem 2.1. *Let $t < s < n/2$, and let G be a graph of order n such that every induced subgraph of G on s vertices contains an independent set of size t . Denote $k = \lfloor \frac{s}{t-1} \rfloor$. Then G contains an independent set of size at least $\Omega(kn^{1/k})$ if $k \leq 2 \log n$ and of size at least $\Omega\left(\frac{\log n}{\log(k/\log n)}\right)$ if $k > 2 \log n$.*

Theorem 2.2. *Let $2t \leq s < n/2$, and let G be a graph of order n such that every induced subgraph of G on s vertices contains an independent set of size t . Then G contains an independent set of size at least $\Omega\left(t \frac{\log(n/s)}{\log(s/t)}\right)$.*

The next result shows that there are graphs with relatively small independence numbers in which every induced subgraph of size s contains an independent set of size t .

Theorem 2.3. *For every sufficiently large t and $2t \leq s \leq n/2$, there exists a graph G on n vertices with independence number*

$$\alpha(G) \leq O\left(t \left(\frac{n}{s}\right)^{2t/(s-t)} \log(n/t)\right)$$

such that every induced subgraph of G of order s contains an independent set of size t .

For certain values of t and s , one can improve the previous result as follows.

Theorem 2.4. *Let $t < s \leq n/2$, where $s \leq e^{2t}$, and assume, further, that either there exists a constant $\delta > 0$ such that $(s/t)^{1-\delta} \geq \log n$ or $s/t = \Omega(\log n)$ and there exists a constant $\gamma > 0$ such that $\log t \geq \log^\gamma n$. Then there exists a graph G on n*

vertices with

$$\alpha(G) \leq O\left(\frac{t}{\log(s/t)} \log(n/t)\right)$$

such that every induced subgraph of G of order s contains an independent set of size t .

3. THE PROOFS

Proof of Theorem 2.1. Suppose that G contains $t - 1$ disjoint cliques whose union U has size at least s . Then the size of the largest independent set in U is bounded by $t - 1$, since an independent set can intersect each of the $t - 1$ cliques in at most one vertex. This contradicts the property of G . Therefore, if G' is a graph obtained from G by deleting the vertices of $t - 1$ disjoint cliques with maximum union we have that $|V(G')| > n - s \geq n/2$. We also have that the largest clique in G' has size at most $k = \lfloor \frac{s}{t-1} \rfloor$. Otherwise, by the above discussion, G will contain $t - 1$ disjoint cliques each of size at least $k + 1$, whose union has at least $(k + 1)(t - 1) \geq s$ vertices. To finish the proof, we apply the classical bound of Erdős and Szekeres [4] (see also [5]) for the usual graph Ramsey numbers. This result asserts that the maximum possible number of vertices in a graph with neither a clique of size $k + 1$ nor an independent set of size $\ell + 1$ is at most $\binom{k+\ell}{k}$. By this estimate, G' contains an independent set of size ℓ , where $\binom{k+\ell}{k} \geq n/2$. Thus, $(\frac{e(k+\ell)}{k})^k > n/2$ and therefore, if $k \leq 2 \log n$, we get $\alpha(G) \geq \ell \geq \Omega(kn^{1/k})$. On the other hand, if $k > 2 \log n$, we use the estimate $(\frac{e(k+\ell)}{\ell})^\ell \geq \binom{k+\ell}{\ell} = \binom{k+\ell}{k} \geq n/2$. This gives that $\ell \geq \Omega(\frac{\log n}{\log(k/\log n)})$ and completes the proof of the theorem. ■

Proof of Theorem 2.2. Let r be the largest integer such that

$$n \left(\frac{t^2}{4e^2s^2} \right)^{r-1} \geq s.$$

It is easy to check that $r = \Omega(\frac{\log(n/s)}{\log(s/t)})$. To prove the theorem, we construct a sequence of pairwise disjoint independent sets X_1, \dots, X_r together with a sequence of nested subsets $V_0 = V(G) \supset V_1 \supset \dots \supset V_{r-1}$ such that the following holds. Each X_i is a subset of V_{i-1} of size $t/2$, $X_i \cap V_i = \emptyset$, there are no edges between X_i and V_i and $|V_i| \geq \frac{t^2}{4e^2s^2} |V_{i-1}|$ for all $1 \leq i \leq r - 1$. Then the union of all sets X_i forms an independent set in G of size $rt/2 = \Omega(t \frac{\log(n/s)}{\log(s/t)})$.

Assuming the sets X_j, V_j have already been constructed for all $j < i$, construct X_i and V_i as follows. Let m be the size of V_{i-1} . The inductive hypothesis implies that

$$m \geq \left(\frac{t^2}{4e^2s^2} \right)^{i-1} |V_0| \geq n \left(\frac{t^2}{4e^2s^2} \right)^{r-1} \geq s.$$

Since every subset of order s in V_{i-1} contains t independent vertices and every independent set of size t belongs to at most $\binom{m-s}{s-t}$ subsets of size s we have that V_{i-1} contains at least $\binom{m}{s} / \binom{m-t}{s-t}$ independent sets of size t . Therefore, using that $m \geq s \geq 2t$ and $\binom{a}{b} \leq \left(\frac{ea}{b}\right)^b$, we conclude that there exists a subset X_i of V_{i-1} of size $t/2$ which is contained in at least

$$\begin{aligned} \frac{\binom{m}{s}}{\binom{m-t}{s-t}\binom{m}{t/2}} &= \frac{m!}{(m-t)!} \frac{(s-t)!}{s!} \binom{m}{t/2}^{-1} \geq \left(\frac{m}{s}\right)^t \left(\frac{t}{2em}\right)^{t/2} \\ &= \left(\frac{mt}{2es^2}\right)^{t/2} = \left(\frac{e\left(\frac{t^2}{4e^2s^2}m\right)}{t/2}\right)^{t/2} \\ &\geq \binom{\frac{t^2}{4e^2s^2}m}{t/2} \end{aligned}$$

independent sets of size t . This implies that V_{i-1} contains at least $\binom{\frac{t^2}{4e^2s^2}m}{t/2}$ subsets of size $t/2$ whose union with X_i forms an independent set of size t . Let V_i be the union of all these subsets. By definition, for every vertex of V_i there is an independent set that contains it together with X_i , so there are no edges from X_i to V_i . Also, it is easy to see that the size of V_i must be at least $\frac{t^2}{4e^2s^2}m = \frac{t^2}{4e^2s^2}|V_{i-1}|$. This completes the construction step and the proof of the theorem. ■

Proof of Theorem 2.3. We prove the theorem by considering an appropriate random graph. As usual, let $G_{n,p}$ denote the probability space of all labeled graphs on n vertices, where every edge appears randomly and independently with probability $p = p(n)$. We say that the random graph possesses a graph property \mathcal{P} almost surely, or a.s. for brevity, if the probability that $G_{n,p}$ satisfies \mathcal{P} tends to 1 as n tends to infinity. Clearly, it is enough to show that there is a value of the edge probability p such that $G_{n,p}$ satisfies the assertion of the theorem with positive probability.

Let $p = \frac{1}{4e^3t} \left(\frac{n}{s}\right)^{-2t/(s-t)}$. We claim that almost surely every subset of $G = G_{n,p}$ of size s spans at most $s^2/(2t) - s/2$ edges. Indeed, the probability that there is a subset of size s that violates this assertion is at most

$$\begin{aligned} \mathbb{P} &\leq \binom{n}{s} \binom{s^2/2}{s^2/(2t) - s/2} p^{s^2/(2t) - s/2} \leq \left(\frac{en}{s}\right)^s \left(e \frac{s}{s-t} tp\right)^{s^2/(2t) - s/2} \\ &\leq \left(\frac{en}{s}\right)^s \left(\frac{1}{2e^2} \left(\frac{n}{s}\right)^{-2t/(s-t)}\right)^{s^2/(2t) - s/2} \leq 2^{-s/2} = o(1), \end{aligned}$$

(where the $o(1)$ -term tends to zero as s tends to infinity). This implies that with high probability every subgraph of G on s vertices has average degree $d \leq s/t -$

1. Therefore, by Turán’s theorem, it contains an independent set of size at least $\frac{s}{d+1} \geq t$. On the other hand, it is well known (see, e.g., [2]) that almost surely the independence number of $G_{n,p}$ is bounded by

$$\alpha(G_{n,p}) \leq O(p^{-1} \log np) \leq O\left(t \left(\frac{n}{s}\right)^{2t/(s-t)} \log(n/t)\right).$$

This implies that a.s. $G_{n,p}$ satisfies the assertion of the theorem and completes the proof. ■

For the proof of Theorem 2.4, we need the following lemma.

Lemma 3.1. *Let $G = G_{s,p}$ be a random graph, assume $sp \rightarrow \infty$ and fix $\epsilon > 0$. Then the probability that the independence number of G is at most $\frac{\epsilon}{p} \log(sp)$ is less than $e^{-s(sp)^{1-3\epsilon/2}}$.*

Proof. Let $k = \frac{\epsilon}{p} \log(sp)$. To prove the lemma, we use the standard greedy algorithm which constructs an independent set by examining the vertices of the graph in some fixed order and by adding a vertex to the current independent set whenever possible. The behavior of this algorithm for random graphs can be analyzed rather accurately (see, e.g., [2]). At iteration i of our procedure, we use the greedy algorithm to find a maximal (with respect to inclusion) independent set I_i in the remaining vertices of G . If $|I_i| \geq k$, we stop. Otherwise, we delete the vertices of I_i from G and continue. We stop when the number of remaining vertices drops below $s/2$. Note that during iteration number i , we only expose edges incident to I_i ; therefore, the remaining vertices still form a truly random graph. Given a set I the probability that a fixed vertex of G is adjacent to some vertex of I is $1 - (1 - p)^{|I|}$. Therefore, the probability that a fixed I is maximal, where $|I| \leq k$, is at most $(1 - (1 - p)^k)^{s/2}$. By definition, when the iteration fails, the random graph must contain a maximal independent set of size less than k (and hence also a set of size exactly k) so that every remaining vertex is adjacent to at least one of its members. Thus, the probability of this event is at most $\binom{s}{k} (1 - (1 - p)^k)^{s/2}$. Moreover, the outcomes of different iterations depend on disjoint sets of edges and therefore are independent. Finally, note that if G has no independent set of size k , then the number of iterations is at least $s/(2k)$. This implies that the probability of such an event is bounded by

$$\begin{aligned} \mathbb{P} &\leq \left(\binom{s}{k} (1 - (1 - p)^k)^{s/2}\right)^{s/(2k)} \leq \left(\frac{eS}{k}\right)^{s/2} e^{-\frac{s^2}{4k}(1-p)^k} \\ &\leq e^{-\Omega\left(s \frac{(sp)^{1-\epsilon}}{\log sp}\right) + s \log(sp)} \leq e^{-s(sp)^{1-3\epsilon/2}}. \end{aligned}$$

■

Proof of Theorem 2.4. Suppose that $(s/t)^{1-\delta} \geq \log n$ for some fixed $\delta > 0$ and consider the random graph $G = G_{n,p}$ with $p = \frac{\delta \log(s/t)}{2t}$. (Note that $p < 1$

as $s \leq e^{2t}$.) As already mentioned above, a.s. the independence number of this graph is bounded by $O(\frac{1}{p} \log(np)) = O(\frac{t}{\log(s/t)} \log(n/t))$. Here, we used that $\log(s/t) < s/t < n/t$ and hence $\log(\frac{n}{t} \log(s/t)) < 2 \log(n/t)$. Also, by Lemma 3.1 (with $\epsilon = \delta/2$), the probability that G contains an induced subgraph of order s with no independent set of size $\frac{\delta}{2p} \log(sp) \geq t$ is at most

$$\binom{n}{s} e^{-s(sp)^{1-3\delta/4}} \leq n^s e^{-s \log^{1+\delta/4} n} = o(1).$$

Here, we used that $\log(s/t) \rightarrow \infty$ and hence $(sp)^{1-3\delta/4} \geq (s/t)^{1-3\delta/4} \geq \log^{1+\delta/4} n$. Therefore, with high probability G satisfies the first assertion of the theorem.

To prove the second part of the theorem, suppose that $s/t = \Omega(\log n)$ and $\log t \geq \log^\gamma n$ for some constant $\gamma > 0$. By the previous paragraph we can also assume that $s/t \leq \log^2 n$. Let $G = G_{n,p}$ with $p = \frac{\gamma \log(s/t)}{640t}$. Again, we have that a.s. $\alpha(G) \leq O(\frac{t}{\log(s/t)} \log(n/t))$. Since $sp = \Omega(\log n \log \log n)$, the probability that there is a subset of size s in G which spans at least $2s^2 p$ edges is bounded by

$$\begin{aligned} \binom{n}{s} \binom{s^2/2}{2s^2 p} p^{2s^2 p} &\leq n^s \left(\frac{e}{4p}\right)^{2s^2 p} p^{2s^2 p} = n^s \left(\frac{e}{4}\right)^{2s^2 p} \\ &\leq e^{s \log n} e^{-\Omega(s \log n \log \log n)} = o(1). \end{aligned}$$

Also, since $s/t \leq sp \leq t^{o(1)}$, we have that the probability that G contains a subset of order $k = \Theta(sp)$ which spans more than $k^{2-\gamma/2}$ edges is at most

$$\begin{aligned} \binom{n}{k} \binom{k^2/2}{k^{2-\gamma/2}} p^{k^{2-\gamma/2}} &\leq n^k (kp)^{k^{2-\gamma/2}} \leq e^{k \log n} e^{-\Omega(k(s/t)^{1-\gamma/2} \log t)} \\ &\leq e^{k \log n} e^{-\Omega(k \log^{1+\gamma/2} n)} = o(1). \end{aligned}$$

Let H be an induced subgraph of G of order s . Since the number of edges in H is a.s. at most $2s^2 p$, it contains an induced subgraph H' of order $s/2$ with maximum degree at most $d = 8sp$. By the above discussion, we also have that the neighborhood of every vertex of H' spans at most $d^{2-\gamma/2}$ edges and therefore every vertex of H' is in at most $T \leq d^{2-\gamma/2}$ triangles. Now, we can use the known estimate (Lemma 12.16 in [2], see also [1] for a more general result) on the independence number of a graph containing a small number of triangles. It implies that

$$\begin{aligned} \alpha(H') &\geq 0.1 \frac{|V(H')|}{d} \left(\log d - \frac{1}{2} \log T \right) \geq 0.1 \frac{s/2}{d} \left(\log d - \frac{1}{2} \log d^{2-\gamma/2} \right) \\ &= \frac{\gamma s}{80d} \log d \geq \frac{\gamma s}{640sp} \log(8sp) \geq t. \end{aligned}$$

This shows that G a.s. satisfies the second assertion of our theorem and completes the proof. ■

4. REMARKS

Theorems 2.1 and 2.3 show that if $s/t = O(1)$ and $t > 1$ then $f(n, s, t) = n^{\Theta(1)}$, whereas if $s/t \gg 1$ and $t = n^{o(1)}$ then $f(n, s, t) = n^{o(1)}$, and if $s/t \geq \Omega(\log n)$ and $t \leq (\log n)^{O(1)}$ then $f(n, s, t) \leq (\log n)^{O(1)}$.

Theorems 2.2 and 2.4 determine the asymptotic behavior of the function $f(n, s, t)$ up to a constant factor for a wide range of the parameters. We do not specify here all this range, and only observe that in particular, for every fixed $\mu > 0$, $f(n, \log^{2+\mu} n, \log n) = \Theta\left(\frac{\log^2 n}{\log \log n}\right)$. For $\mu = 1$, this implies (2). The estimate (1) follows from Theorems 2.2 and 2.3.

The connection between the Ramsey-type question described in Section 1 and the function f is the following simple fact.

Claim. *If*

$$n/2 > s > t \quad \text{and} \quad (t-1)f(n/2, s, t) \geq s, \quad (3)$$

then there is no graph on n vertices in which every induced subgraph on s vertices contains a clique of size t and an independent set of size t .

Proof. Assuming there is such a graph G , observe that by the definition of f it contains an independent set I_1 of size $f = f(n/2, s, t)$. Omit this set, and observe that the induced graph on the remaining vertices, assuming there are at least $n/2$ of them, contains another independent set I_2 of size f . Repeating this process $t-1$ times (or until the union of the independent sets obtained is of size at least $n/2 > s$), we get an induced subgraph of G on $\min\{n/2, (t-1)f\} \geq s$ vertices, which is $(t-1)$ -colorable (as it is the union of $t-1$ independent sets), and hence cannot contain a clique of size t . This is a contradiction, proving the assertion of the claim. ■

In particular, for $t = \log n$ and $s = c \log^3 n / \log \log n$, where c is a sufficiently small positive absolute constant, it is not difficult to check that the assumption in (3) holds, by Theorem 2.2.

It will be interesting to close the gap between our upper and lower bounds for the function $f(n, s, t)$. It will also be interesting to know more about the Ramsey-type question of Erdős and Hajnal described in the introduction, and in particular, to decide if there exists a graph on n vertices in which every induced subgraph on, say, $\log^{100} n$ vertices, contains a clique of size at least $\log n$ and an independent set of size at least $\log n$.

ACKNOWLEDGMENTS

The research work of N.A. was supported in part by the Israel Science Foundation and the Hermann Minkowski Minerva Center for Geometry at Tel Aviv University. B.S. gratefully acknowledges partial support for this research from NSF CAREER award DMS-0546523, NSF grant DMS-0355497, Alfred P. Sloan fellowship, and the State of New Jersey. A USA-Israel BSF grant supported the research work of both the authors.

REFERENCES

- [1] N. Alon, M. Krivelevich, and B. Sudakov, Coloring graphs with sparse neighborhoods, *J Combin Theory B* 77 (1999), 73–82.
- [2] B. Bollobás, *Random graphs*, Second Edition, Cambridge studies in advanced mathematics, Vol. 73, Cambridge University Press, Cambridge, 2001.
- [3] P. Erdős, Problems and results in combinatorial analysis and combinatorial number theory, In: *Graph theory, combinatorics, and applications*, Vol. 1 (Kalamazoo, MI, 1988), Wiley, New York, 1991, pp. 397–406.
- [4] P. Erdős and G. Szekeres, A combinatorial problem in geometry, *Compos Math* 2 (1935), 463–470.
- [5] R. Graham, B. Rothschild, and J. Spencer, *Ramsey theory*, Second Edition, Wiley, New York, 1990.