

THE NUMBER OF
HAMILTONIAN DECOMPOSITIONS OF REGULAR
GRAPHS

BY

ROMAN GLEBOV*

*School of Computer Science and Engineering
The Hebrew University of Jerusalem, Jerusalem 9190401, Israel
and
Department of Mathematics, ETH, 8092 Zurich, Switzerland
e-mail: roman.l.glebov@gmail.com*

AND

ZUR LURIA**

*Institute of Theoretical Studies, ETH, 8092 Zurich, Switzerland
e-mail: zluria@gmail.com*

AND

BENNY SUDAKOV†

*Department of Mathematics, ETH, 8092 Zurich, Switzerland
e-mail: benjamin.sudakov@math.ethz.ch*

* Research supported by the ERC grant “High-dimensional combinatorics” at the Hebrew University of Jerusalem.

** Research supported by Dr. Max Rössler, the Walter Haefner Foundation and the ETH Foundation.

† Research supported in part by SNSF grant 200021-149111.

Received December 21, 2015 and in revised form June 20, 2016

ABSTRACT

A Hamilton cycle in a graph Γ is a cycle passing through every vertex of Γ . A Hamiltonian decomposition of Γ is a partition of its edge set into disjoint Hamilton cycles. One of the oldest results in graph theory is Walecki's theorem from the 19th century, showing that a complete graph K_n on an odd number of vertices n has a Hamiltonian decomposition. This result was recently greatly extended by Kühn and Osthus. They proved that every r -regular n -vertex graph Γ with even degree $r = cn$ for some fixed $c > 1/2$ has a Hamiltonian decomposition, provided $n = n(c)$ is sufficiently large. In this paper we address the natural question of estimating $H(\Gamma)$, the number of such decompositions of Γ . Our main result is that $H(\Gamma) = r^{(1+o(1))nr/2}$. In particular, the number of Hamiltonian decompositions of K_n is $n^{(1+o(1))n^2/2}$.

1. Introduction

A **Hamilton cycle** in a graph Γ is a cycle passing through each vertex of Γ , and a graph is **Hamiltonian** if it contains a Hamilton cycle. Hamiltonicity, named after Sir Rowan Hamilton who studied it in the 1850s, is one of the most important and extensively studied concepts in graph theory. It is well known that deciding Hamiltonicity is an NP-complete problem, and thus one does not expect a simple sufficient condition for Hamiltonicity for general graphs. Once Hamiltonicity is established, it is very natural to strengthen such a result by showing that the graph in question has many edge-disjoint Hamilton cycles, or even has a **Hamiltonian decomposition**, which is a partition of the edge set of the graph into disjoint Hamilton cycles. Clearly a Hamiltonian decomposition is only possible when Γ is r -regular for some even r . In 1890, Walecki gave a celebrated construction of a Hamiltonian decomposition of the complete graph K_n for every odd n . For a description of his construction; see, e.g., [2].

The work of Walecki was extended by various researchers, who proved that more general families of graphs admit a Hamiltonian decomposition; see, e.g., [21, 4, 14, 19, 6] and a survey [3], which gives an overview of many results on this topic. One of the classical results on Hamiltonicity of graphs is a theorem of Dirac which says that every n -vertex graph with minimum degree $n/2$ has a Hamilton cycle. Recently Kühn and Osthus [19] obtained a far reaching generalization of both Dirac's result for regular graphs and Walecki's decomposition. They proved that every r -regular n -vertex graph Γ with even degree $r = cn$

for some fixed $c > 1/2$ has a Hamiltonian decomposition, provided $n = n(c)$ is sufficiently large.

The question of estimating the number of Hamilton cycles in various classes of graphs has also been intensively studied; see, e.g., [13, 10, 7, 16, 12, 8] and the references therein. However, to the best of our knowledge, so far only few results have been established on the number of Hamiltonian decompositions; see, e.g., [9] and the references therein. In this paper we study the first natural question of this sort, providing counting versions of Walecki’s and Kühn-Osthus’ results. We are interested in $H(\Gamma)$, the number of Hamiltonian decompositions of a given r -regular n -vertex graph Γ with even degree $r = cn$ for some fixed constant $c > 1/2$.

To upper bound the number of Hamilton cycles in a regular graph, one can use the standard upper bound for the permanent of a 0-1 matrix proved by Brégman [5] (solving the famous Minc conjecture). This permanent-based approach to Hamiltonicity problems was used for the first time by Alon [1] to bound the number of Hamilton paths in tournaments (see also [11, 16, 12, 15, 8] for additional applications). Let S_n be the set of all permutations of the set $[n]$. The **permanent** of an $n \times n$ matrix A is defined as

$$\text{per}(A) = \sum_{\sigma \in S_n} \prod_{i=1}^n A_{i\sigma(i)}.$$

Note that when $A = A_\Gamma$ is the 0-1 adjacency matrix of a graph Γ , a summand in the permanent is 1 if it corresponds to a spanning subgraph of Γ whose connected components are either cycles or isolated edges, and the summand is 0 otherwise. Therefore, the number of Hamilton cycles in Γ is bounded from above by $\text{per}(A_\Gamma)$. Combining this observation with Brégman’s [5] upper bound, we see that an r -regular graph has at most $(r!)^{n/r}$ Hamilton cycles. By choosing any one of these Hamilton cycles and deleting it, we obtain an $(r - 2)$ -regular graph. The number of Hamilton cycles in the new graph can again be bounded from above using Brégman’s theorem. Continuing this process and multiplying all the estimates for regularities $r, r - 2, r - 4, \dots$ we can use Stirling’s formula to deduce the following upper bound.

PROPOSITION 1.1: *For every $r = r(n) \rightarrow \infty$, the number of Hamiltonian decompositions of an r -regular graph Γ of order n is at most*

$$\left((1 + o(1)) \frac{r}{e^2} \right)^{nr/2}.$$

Our main result is the corresponding lower bound, which together with the last proposition determines asymptotically the number of Hamiltonian decompositions of dense regular graphs.

THEOREM 1.2: *Let $c > 1/2$ be a constant and let Γ be an n -vertex r -regular graph with even degree $r \geq cn$. Then the number of Hamiltonian decompositions of Γ satisfies*

$$H(\Gamma) = r^{(1+o(1))rn/2}.$$

In particular the number of Hamiltonian decompositions of K_n is $n^{(1+o(1))n^2/2}$.

1.1. OUTLINE OF THE PROOF. Let Γ be an n -vertex r -regular graph with even degree $r \geq cn$ and $c > 1/2$. Let δ be an arbitrary constant $\delta < \frac{1}{2}(c - 1/2)$ and let A and B be two subsets of $V(\Gamma)$ such that $|A| \geq \delta^2 n$ and $|B| \geq (1/2 - \delta)n$. Every vertex in A has at least $cn - (n - |B|) > \delta n$ neighbors in B . Therefore the number of edges between A and B satisfies

$$e_\Gamma(A, B) \geq \delta n|A|/2 \geq \delta^3 n^2/2.$$

(If A and B have a nonempty intersection, then $e_\Gamma(A, B)$ counts the edges inside $A \cap B$ with multiplicity one.) Therefore, our main result follows from the following more general statement (by taking $\gamma = \delta^3/2$ and $\varepsilon \rightarrow 0$).

THEOREM 1.3: *For every $c > 0$ and $0 < \varepsilon < \frac{1}{10}$, there exists $\delta > 0$ such that for any constant $\gamma > 0$ and sufficiently large n the following holds. Let Γ be an n -vertex r -regular graph with even degree $r \geq cn$ such that there are at least γn^2 edges between any two subsets $A, B \subseteq V(G)$ satisfying $|A| \geq \delta^2 n$ and $|B| \geq (1/2 - \delta)n$. Then the number of Hamiltonian decompositions of Γ is at least $r^{(1-5\varepsilon)rn/2}$.*

The proof of this statement follows the same general strategy as the proof of Proposition 1.1. Namely, we remove Hamilton cycles one by one, giving a lower bound on the number of possibilities at each step. The two additional ingredients we need are a way to construct many Hamilton cycles in a regular graph, and sufficient conditions for the existence of a Hamiltonian decomposition in a graph.

The proof consists of three parts. In the first part, we split Γ into three edge-disjoint spanning graphs G, F , and R . The graph G contains most of the

edges of Γ , F satisfies a pseudo-randomness condition similar to the expander mixing lemma, and R is a robust expander, a concept that is defined below.

In the second part, we use some ideas of Ferber, Krivelevich and Sudakov [8] to repeatedly find Hamilton cycles in $G \cup F$. Their approach, based on the permanent of the adjacency matrix, gives us at every step many 2-regular subgraphs of the current graph which are unions of a small number of cycles. Using edges of F we can transform each of these subgraphs into a Hamilton cycle. This part of the proof gives the asymptotic count in the theorem. We stop at some point when all but a small constant fraction of the edges of $G \cup F$ have been covered.

In the third part of the proof, we need to show that the remaining edges of Γ can be partitioned into Hamilton cycles. Here, we use a recent result of Kühn and Osthus [19]. Given parameters $0 < \nu \leq \tau < 1$, a graph \mathcal{G} and a set of vertices $S \subseteq V(\mathcal{G})$, the robust ν -neighborhood of S is the set of vertices that have at least νn neighbors in S , and is denoted by $RN_{\nu, \mathcal{G}}(S)$. A graph \mathcal{G} is a robust (ν, τ) -expander if

$$RN_{\nu, \mathcal{G}}(S) \geq |S| + \nu n \text{ for all } S \text{ such that } \tau n \leq |S| \leq (1 - \tau)n.$$

Note in particular that this is a monotone property under addition of edges. The following powerful theorem of Kühn and Osthus [19], which they derived from the corresponding digraph result [18], provides sufficient conditions for a graph to admit a Hamiltonian decomposition.

THEOREM 1.4 (Theorem 1.2 in [19]): *For every $\alpha > 0$ there exists $\tau > 0$ such that for every $\nu > 0$ there exists $n_0(\alpha, \nu, \tau)$ for which the following holds. Suppose that*

- \mathcal{G} is an r -regular graph on $n \geq n_0$ vertices, where $r \geq \alpha n$ is even, and
- \mathcal{G} is a robust (ν, τ) -expander.

Then \mathcal{G} has a Hamiltonian decomposition.

Since R is a robust expander, the graph consisting of all remaining edges of Γ that are not covered by the Hamilton cycles found in the second part satisfies the conditions of Theorem 1.4, and therefore we can complete our packing of edge-disjoint Hamilton cycles to a Hamiltonian decomposition of Γ .

Throughout the proof, we sometimes identify a graph with its edge set. For the sake of clarity of presentation, we omit floor and ceiling signs.

2. Proof of Theorem 1.3

Given $c, \gamma > 0$ and $0 < \varepsilon < \frac{1}{10}$, choose $\alpha = 3\varepsilon c$ and let $\tau = \tau(\alpha)$ be as in Theorem 1.4. Next, choose $\delta \leq \min(\varepsilon c/5, \tau/2)$ and set $\nu = \min(\delta, \varepsilon\gamma/2)$. Finally, let $n_0 = n_0(\alpha, \nu, \tau)$ be as in Theorem 1.4. Throughout this section we will always assume that $n > n_0$ and is sufficiently large whenever it is necessary.

We first show how to find three edge-disjoint graphs $G, F,$ and R with the desired properties.

LEMMA 2.1: *The edges of Γ can be partitioned into three subgraphs $G, F,$ and R such that:*

- *The graph G is d -regular with even degree $d \geq (1 - 2\varepsilon)r$.*
- *For any two sets $A, B \subseteq [n]$ of sizes $|A| \geq \delta^2 n$ and $|B| \geq (1/2 - \delta)n$, there exist at least $n^{1.6}$ edges in F between A and B .*
- *The graph R is a robust (ν, τ) -expander.*

Proof. Consider the edge partition of Γ into three edge-disjoint random subgraphs $F, R^*,$ and G^* , where every edge gets into F with probability $1/\log n$, into R^* with probability ε , and into G^* with probability $1 - \varepsilon - 1/\log n$.

First, we need to find a dense regular spanning subgraph $G \subseteq G^*$. To do so we will use the following lemma, which we prove in the Appendix to this paper.

LEMMA 2.2: *Let $\varepsilon_0, \gamma_0 > 0$ be constants, $c_0 \geq \varepsilon_0$, and let n be sufficiently large. Let G^* be an n -vertex graph whose vertex degrees are all between $c_0 n - n^{2/3}$ and $c_0 n + n^{2/3}$, such that there are at least $\gamma_0 n^2$ edges between any two subsets $A, B \subseteq V(G^*)$ satisfying $|A| \geq c_0 n/3$ and $|B| \geq n/2$. Then G^* has a regular spanning subgraph of degree $2d$, where $d = \lceil \frac{(c_0 - \varepsilon_0)n}{2} \rceil$.*

We claim that with high probability G^* satisfies the conditions of this Lemma, with the parameters $c_0 = (1 - \varepsilon - 1/\log(n))\frac{r}{n}$ and $\gamma_0 = \frac{\gamma}{2}$. Indeed, this follows from a straightforward application of Chernoff’s inequality and the union bound after observing that $c_0/3 > \delta^2$. We apply Lemma 2.2 with $\varepsilon_0 = \frac{\varepsilon r}{2n} < c_0$, obtaining the desired d -regular graph G with $d \geq (1 - 2\varepsilon)r$.

Next, consider the graph F . Between any two sets $A, B \subseteq [n]$ of sizes $|A| \geq \delta^2 n$ and $|B| \geq (1/2 - \delta)n$, there are at least γn^2 edges of Γ . Therefore, the number of edges between A and B in F is distributed binomially with parameters $|E_\Gamma(A, B)| \geq \gamma n^2$ and $\log^{-1} n$. Using Chernoff-type estimates again, we can see that the probability of having less than $n^{1.6}$ edges in F between A

and B is $\ll 2^{-2n}$. Taking a union bound over all pairs of subsets shows that with high probability Γ has the desired property.

Note that the same argument as above using Chernoff-type estimates and the union bound implies that with high probability there are at least $\varepsilon\gamma n^2/2$ edges of R^* between any two sets $A, B \subseteq [n]$ of sizes $|A| \geq \delta^2 n$ and $|B| \geq (1/2 - \delta)n$. Using this we prove that R^* is a robust expander. First fix a subset S of the vertices of R^* such that $\delta^2 n < \tau n \leq |S| < n/2$, and consider the set B of vertices of R^* which have less than $\nu n \leq \varepsilon\gamma n/2$ neighbors in S . If there are at least $(1/2 - \delta)n$ such vertices, then between B and S there are less than $\varepsilon\gamma n^2/2$ edges, contradiction. Therefore the robust ν -neighborhood of S has size at least $(1/2 + \delta)n \geq |S| + \nu n$. Now consider a subset S of size $n/2 \leq |S| \leq (1 - \tau)n \leq n - 2\delta n$ and let A be the set of vertices of R^* which have less than $\nu n \leq \varepsilon\gamma n/2$ neighbors in S . If there are at least $\delta^2 n$ such vertices then there are less than $\varepsilon\gamma n^2/2$ edges between A and S , again obtaining a contradiction. Therefore the robust ν -neighborhood of S has size at least $n - \delta^2 n \geq |S| + \nu n$ and so R^* is a robust (ν, τ) -expander. Finally, define R to be the union of R^* together with all the edges from $G^* \setminus G$. By monotonicity, R is a robust expander as well. ■

As we already mentioned above, our proof proceeds by successively removing Hamilton cycles from $G \cup F$. The following lemma provides a lower bound on the number of choices in each such step.

LEMMA 2.3: *Let $\varepsilon > 0$ and let G and F be edge-disjoint graphs on the vertex set $[n]$ such that the graph G is d -regular with $d \geq \varepsilon r = \varepsilon cn$ and, for any two sets $A, B \subseteq [n]$ of sizes at least $|A| \geq \delta^2 n$ and $|B| \geq (1/2 - \delta)n$, there are at least $dn^{0.6}$ edges of F between A and B . Then there exist at least $(1 + o(1))^n (d/e)^n$ Hamilton cycles H in $G \cup F$ with the following properties. For each cycle H there are edge sets $E_G \subseteq G$ and $E_F \subseteq F$ such that:*

- $G' := (G \cup E_F) \setminus (H \cup E_G)$ is $(d - 2)$ -regular.
- $F' := F \setminus (H \cup E_F)$ has the property that for any $A, B \subseteq [n]$ of sizes at least $|A| \geq \delta^2 n$ and $|B| \geq (1/2 - \delta)n$, there are at least $(d - 2)n^{0.6}$ edges of F' between A and B .

Remark: In order to keep the graph obtained from G after removing the cycle H regular we need to add and delete some additional edges. This is achieved by using the sets of edges E_G and E_F . Note that the resulting graphs G', F' , and H are edge-disjoint.

Proof. We call a spanning subgraph of $G \cup F$ a **partial HC** (for Hamilton cycle) if one of its components is a path, and all other components are cycles or isolated edges. We also call a spanning collection of vertex-disjoint cycles and isolated edges a (≤ 2) -**factor**. The following claim allows us to reduce the number of connected components in a 2-factor by concatenating two of its cycles into a path.

CLAIM 2.4: *Let H be a 2-factor in $G \cup F$ with more than one cycle. There exist edges $e \in (G \cup F) \setminus H$ and $e'_1, e'_2 \in H$ such that*

$$H \cup \{e\} \setminus \{e'_1, e'_2\}$$

is a partial HC with one less component (and one less edge) than H .

Proof. Consider an arbitrary cycle C of H on at most $n/2$ vertices. If $|C| < d$, then there exists an edge $e \in E(G)$ connecting C with a different component of H . If $|C| \geq d \geq \varepsilon cn \geq \delta^2 n$, then again there exists an edge $e \in E(F)$ connecting C with a different component of H . In both cases, obviously $e \notin H$. Now any two edges $e'_1, e'_2 \in E(H)$ adjacent to different endpoints of e satisfy the requirements of the claim. ■

The following technical claim allows us to either extend a path in a partial HC, thereby reducing the number of connected components, or to close this path into a cycle.

CLAIM 2.5: *Let H be a partial HC in $G \cup F$. Then one of the following holds:*

- *There exist edges $e, e_1, e_2 \in (G \cup F) \setminus H$ and $e', e'_1, e'_2 \in H$ such that $H \cup \{e\} \setminus \{e'\}$, or $H \cup \{e\}$, or $H \cup \{e, e_1, e_2\} \setminus \{e', e'_1, e'_2\}$, or $H \cup \{e, e_1, e_2\} \setminus \{e'_1, e'_2\}$ is a partial HC with one less component than H (and at least as many edges as H).*
- *There exist edges $e_1, e_2, e_3 \in (G \cup F) \setminus H$, $e'_1, e'_2, e'_3 \in H$, and $f \in F \setminus H$ such that $H \cup \{e_1, e_2, e_3, f\} \setminus \{e'_1, e'_2, e'_3\}$ is a (≤ 2) -factor in $G \cup F$ with the same number of components as H (and one more edge than H).*

Proof. At least one component of H is a path; choose $P = v_1, \dots, v_t$ to be a longest such. If v_1 or v_t have one of their neighbors outside P , say $(v_1 z) \in E(G)$ with $z \notin P$, then $(v_1 z)$ connects P to a component D of H with $z \in D$. If D is an isolated edge, then $H \cup (v_1 z)$ satisfies the conditions of the first bullet. Otherwise, D is a cycle and we can choose an arbitrary edge $e' \in E(D)$ incident to z , and $H \cup (v_1 z) \setminus e'$ satisfies the conditions of the first bullet.

Therefore we can assume that all the neighbors of v_1, v_t in G lie in P . Then we will use a variant of the celebrated rotation-extension technique of Pósa [20], which was developed by the third author together with Vu (see the proof of Theorem 4.1 in [22]).

We split P into $s = 2\delta^{-1}$ segments I_1, \dots, I_s of size $|P|/s \leq \delta n/2$ each, and consider the graph G . By the pigeonhole principle, there exist $p, q \in [s]$ such that v_1 has at least $d/s \geq \varepsilon c\delta n/2$ neighbors in I_p , and v_t has at least $\varepsilon c\delta n/2$ neighbors in I_q . If $p = q$, split I_p further into two segments J_1 and J_2 such that the **interior** of J_1 , i.e., all its vertices except its endpoints, contain at least $\varepsilon c\delta n/5$ neighbors of v_1 , and v_t has at least $\varepsilon c\delta n/5$ neighbors in the interior of J_2 . If $p \neq q$, simply set $J_1 = I_p$ and $J_2 = I_q$. For every neighbor v_i of v_t from the interior of J_2 , we can **rotate** the path P keeping its endpoint v_1 fixed and using v_i as the **pivot**. This means that we add the edge (v_i, v_t) , delete the edge (v_i, v_{i+1}) , and consider the new path $v_1, \dots, v_i, v_t, \dots, v_{i+1}$ with endpoints v_1 and v_{i+1} . Setting A to be the set of successors of the neighbors of v_t in the interior of J_2 , then for every $a \in A$, we obtain a (v_1, a) -path on the vertex set $V(P)$ differing from P in exactly one edge from J_2 . Similarly, for every neighbor v_j of v_1 in the interior of J_1 , every (v_1, a) -path can be further rotated using the pivot v_j . Note that since J_1 and J_2 are disjoint, all these paths we constructed using pivots in J_1 traverse J_2 in the same direction. Therefore the new endpoints we will obtain do not depend on a . Thus we have constructed a set B such that for every $a \in A, b \in B$, there exists a (b, a) -path in G on the vertex set $V(P)$ differing from P in exactly two edges from J_1 and J_2 . Furthermore, $|A|, |B| \geq \varepsilon c\delta n/5 \geq \delta^2 n$, and all these paths traverse all other segments of P outside of J_1 and J_2 in the same direction.

If a vertex $x \in A \cup B$ has a neighbor outside $V(P)$ in $F \cup G$, we extend P similarly to the case discussed above, obtaining a partial HC with one less component than H (and at least as many edges as H), as desired in the first bullet of the claim. Therefore we can assume that all the neighbors of vertices in $A \cup B$ in G lie in P . Then for every $a \in A, b \in B$ and every neighbor v_ℓ of a in G outside of J_1 and J_2 , we can rotate every (b, a) -path further using the pivot v_ℓ to obtain a new endpoint c and a (b, c) -path. Note that, since we did not touch any edge outside J_1 and J_2 before and the parts of P outside these intervals are traversed by all paths in the same direction, the vertex c is determined only by the neighbor v_ℓ of a and does not depend on the other endpoint b . Therefore, we obtain a set S such that for every $b \in B$ and $c \in S$, there exists a (b, c) -path




Figure 1. An illustration of the second bullet of Claim 2.5: Closing the partial HC into a (≤ 2) -factor. The edges e_1, e_2, e_3, f are added and the edges e'_1, e'_2, e'_3 are deleted. The edge set of the new cycle is $P \cup \{e_1, e_2, e_3, f\} \setminus \{e'_1, e'_2, e'_3\}$.

in G on the same vertex set as P , which differs from P in exactly three edges. Furthermore, we know that F has an edge between any two subsets of size $\delta^2 n$ and $(1/2 - \delta)n$. This implies that $|N_F(A)| \geq n - (1/2 - \delta)n = (1/2 + \delta)n$. Therefore

$$|S| \geq |N_F(A)| - |J_1 \cup J_2| \geq (1/2 + \delta)n - \delta n \geq n/2.$$

Similarly we also have that $|N_F(B)| \geq (1/2 + \delta)n$. Thus there exists an edge $f = (b, c) \in E(F)$ between vertices $b \in B$ and $c \in S$. This edge closes the (b, c) -path into a cycle; see Figure 1. Hence we can use the edge f , together with the three edges from the rotations of P which produced the (b, c) -path, to transform a partial HC into a (≤ 2) -factor, satisfying the second bullet of the claim. ■

Observation 2.6: For any vertices $x, y \in [n]$ and any $S \subseteq [n]$ of size at most $|S| \leq n^{0.6}$, there exist vertices $x_1, x_2, y_1, y_2 \in [n] \setminus S$ such that $xx_1, yy_1, x_2y_2 \in E(G)$ and $x_1x_2, y_1y_2 \in E(F)$.

Proof. Since G is d -regular with $d \geq \varepsilon cn \geq 2\delta^2 n + n^{0.6}$ we can choose disjoint sets $A_x \subset N_G(x)$ of size $\delta^2 n$ and $A_y \subset N_G(y)$ of size $\delta^2 n$ which are also disjoint from S . Further, we denote

$$B_x = N_F(A_x) \setminus (S \cup A_x \cup A_y) \quad \text{and} \quad B_y = N_F(A_y) \setminus (S \cup A_x \cup A_y).$$

Since, as we already explained above, the neighborhood in F of any set of size at least $\delta^2 n$ has size at least $(1/2 + \delta)n$, we conclude that $|B_x|, |B_y| > n/2$.

Since G is regular, every subset X of G satisfies $|N_G(X)| \geq |X|$. Indeed

$$d|X| = \sum_{v \in X} d(v) \leq \sum_{u \in N_G(X)} d(u) = d|N_G(X)|.$$

Therefore there exists an edge x_2y_2 in G with $x_2 \in B_x$ and $y_2 \in B_y$. Choosing a neighbor $x_1 \in A_x$ of x_2 and a neighbor $y_1 \in A_y$ of y_2 finishes the proof. ■

We are now ready to complete the proof of Lemma 2.3. Let $s^* = \sqrt{n \log n}$. To lower bound the number of (≤ 2) -factors with at most s^* components in G , we use Corollary 2.8 in [8].

PROPOSITION 2.7 (Corollary 2.8 in [8]): *Let $\tilde{\alpha} > 0$ be a constant and let n be a positive integer. Suppose that:*

- \tilde{G} is a graph on n vertices, and
- \tilde{G} is $\tilde{\alpha}n$ -regular.

Then the number of (≤ 2) -factors of \tilde{G} with at most s^ cycles is $(1+o(1))^n (\frac{\tilde{\alpha}n}{e})^n$.*

Therefore, the number of (≤ 2) -factors with at most s^* components in G is at least $(1+o(1))^n (d/e)^n$. Fix one such factor and call it H . We will transform H into a Hamilton cycle, using edges from G and F . During this process, for the sake of clarity and convenience we will still use H to denote all the new factors and partial HC which we obtain from it.

As long as H is not a Hamilton cycle, we will modify it using Claim 2.5 if it is a partial HC, and Claim 2.4 if H is a 2-factor. Note that, after at most every two such steps, the number of components in H goes down, and therefore after at most $2s^* + 1$ iterations we end up with a Hamilton cycle H . Since the Hamming distance between the original (≤ 2) -factor and the final Hamilton cycle is $O(s^*)$, the number of possible Hamilton cycles H which we obtain in this process is at least

$$\frac{(1+o(1))^n (d/e)^n}{n^{O(s^*)}} = (1+o(1))^n (d/e)^n.$$

It remains to construct the edge sets E_F and E_G , whose role is to keep the resulting graph regular. We denote

$$\mathcal{E}_F = H \cap F,$$

and note that $|\mathcal{E}_F| = O(s^*)$. One by one, for every pair of vertices $x, y \in [n]$ with $xy \in \mathcal{E}_F$, we apply Observation 2.6 to find vertices x_1, x_2, y_1, y_2 such that $xx_1, yy_1, x_2y_2 \in E(G)$ and $x_1x_2, y_1y_2 \in E(F)$. We choose it so that none of the

vertices x_1, x_2, y_1, y_2 is incident to an edge from \mathcal{E}_F or a previously chosen edge from E_F . We then add the edges x_1x_2, y_1y_2 to E_F and the edges xx_1, yy_1, x_2y_2 to E_G . Since at most $O(s^*)$ edges from \mathcal{E}_F and E_F were chosen before, we can always put all endpoints of such edges into the set S for the further applications of Observation 2.6.

We claim that the sets E_F and E_G together with the Hamilton cycle H satisfy the requirements of the lemma. Indeed, in constructing $F' = F \setminus (H \cup E_F)$ we delete at most $O(s^*) = o(n^{0.6})$ edges from F , and so the number of edges in F between two sets A and B of sizes $|A| \geq \delta^2 n$ and $|B| \geq (1/2 - \delta)n$ remains at least $dn^{0.6} - O(s^*) > (d-2)n^{0.6}$. In addition, note that $G' = (G \cup E_F) \setminus (H \cup E_G)$ is $(d-2)$ -regular. In the transition from G to G' , every vertex that isn't incident to an edge of \mathcal{E}_F loses the two edges of H that pass through it. On the other hand, for every vertex incident to an edge of \mathcal{E}_F we remove one edge from $G \cap H$ incident to it, and then delete an edge from E_G incident to it. The edges from E_F ensure that this doesn't change the degree of other vertices. ■

Proof of Theorem 1.3. Fix an edge partition of the graph Γ into graphs $G, F,$ and R which satisfy the assertions of Lemma 2.1. Starting with the d_0 -regular graph $G_0 = G$ with $d_0 \geq (1 - 2\varepsilon)r$ and the graph $F_0 = F$, we apply Lemma 2.3 repeatedly to remove Hamilton cycles from $G_i \cup F_i$. After the i 'th iteration, we will have a d_i -regular graph G_i with $d_i = d_0 - 2i$ and a graph F_i satisfying the second bullet of Lemma 2.3. Therefore the process can be continued. We stop after $t = \frac{d_0 - \varepsilon r}{2}$ steps, at which point G_t is an εr -regular graph. The bound from Lemma 2.3 guarantees that there are at least $((1 + o(1))\frac{d_i}{e})^n$ choices for a Hamilton cycle in the i 'th step. Therefore, this procedure results in at least

$$\begin{aligned} \prod_{i=0}^{t-1} \left((1 + o(1))\frac{d_i}{e} \right)^n &= \left((1 + o(1))^t e^{-t} \frac{2^{d_0/2} (d_0/2)!}{2^{\varepsilon r/2} (\varepsilon r/2)!} \right)^n \\ &\geq \left(e^{-r/2} \frac{\left(\frac{(1-2\varepsilon)r}{2}\right)!}{\left(\frac{\varepsilon r}{2}\right)!} \right)^n \geq r^{(1-4\varepsilon)\frac{r}{2}} \end{aligned}$$

possible ordered tuples of t edge-disjoint Hamilton cycles.

Let us fix one such collection of t -edge disjoint Hamilton cycles and let R' denote the union of R and the edges of $G \cup F$ that do not belong to any of these cycles. The graph R' is αn -regular ($\alpha = 3\varepsilon c$), since it is obtained from Γ by deleting t edge-disjoint Hamilton cycles. The graph R' is also a robust (ν, τ) -expander, since, as we mentioned before, this property is monotone and R

satisfies it. Since $n \geq n_0$, it follows from Theorem 1.4 that R' admits (at least one) Hamiltonian decomposition. Together with the above Hamilton cycles it gives us the desired Hamiltonian decomposition of Γ .

Clearly the same decompositions of Γ can appear in our process in at most $(\frac{n-1}{2})! < n^n$ different orderings. Therefore, we have that

$$H(\Gamma) \geq r^{(1-4\varepsilon)\frac{rn}{2}} / n^n \geq r^{(1-5\varepsilon)\frac{rn}{2}}. \quad \blacksquare$$

3. Concluding remarks and open questions

- A substantial gap remains between the upper bound of Proposition 1.1 and the main result that we prove. We tend to believe that the upper bound of the proposition is close to the truth, i.e,

$$H(\Gamma) = \left((1 + o(1)) \frac{r}{e^2} \right)^{rn/2}$$

for an n -vertex r -regular graph Γ with $r = cn, c > 1/2$. To prove such an estimate using our methods, one needs a Hamiltonian decomposition result for a sparse analogue of robust expanders.

- When the degree r of the graph Γ is odd, the result of Kühn and Osthus also implies that the edges of Γ can be decomposed into $(r - 1)/2$ Hamilton cycles and a perfect matching. Our proof can be easily adapted to show that the number of such decompositions is $r^{(1+o(1))rn/2}$.
- It would be interesting to estimate the number of Hamiltonian decompositions of other families of graphs. In particular, Theorem 1.1.3. in [6] states that for even $r \geq \lfloor \frac{n}{2} \rfloor$, every r -regular graph on n vertices has a Hamiltonian decomposition. Can one strengthen our theorem by giving an accurate estimate of the number of such decompositions for all r , starting with $\lfloor \frac{n}{2} \rfloor$?
- Finally, our main result can be further generalized to subgraphs of quasirandom graphs. A graph G on n vertices is called (α, β) -regular if its minimum degree is at least $\alpha n - 1$ and

$$\left| \frac{e_G(S, T)}{|S||T|} - \alpha \right| \leq \beta$$

for every pair of disjoint sets S, T of size at least βn . It is easy to see that a complete graph on n vertices is $(1, \beta)$ -regular for any $\beta > 0$. Another example of such a graph is an αn -regular graph on n vertices

whose nontrivial eigenvalues in absolute value are all $o(n)$; see, e.g., the survey [17] for more details. We can obtain the following result on the number of Hamiltonian decompositions of regular subgraphs of (α, β) -regular graphs.

THEOREM 3.1: *For all constants c, α such that $c > \alpha/2$ there exists $\beta > 0$ such that the following holds. Let Γ be an r -regular subgraph of an (α, β) -regular graph G such that $r = cn$ is even. Then the number of Hamiltonian decompositions of Γ satisfies $H(\Gamma) = r^{(1+o(1))rn/2}$.*

When G is a complete graph K_n this implies Theorem 1.2.

To prove this theorem, using our Theorem 1.3, one needs only to show that there are at least γn^2 edges (for some very small but fixed γ) between any two subsets $A, |A| = \delta^2 n$ and $B, |B| = (1/2 - \delta)n$ of $V(\Gamma)$. Choose $\delta \ll (c - \alpha/2)$ and $\beta \ll \delta^2$. If Γ contains two subsets A, B violating the above condition, then by the regularity of Γ the number of its edges between A and $V(\Gamma) - B$ is very close to $cn|A|$. On the other hand the graph G , containing Γ , has roughly

$$\alpha(n - |B|)|A| = \alpha(1/2 + \delta)|A|$$

edges between these sets (which is less), a contradiction. We omit further calculation.

ACKNOWLEDGMENT. We would like to thank an anonymous referee for the thorough review of the paper and for pointing out an error in the proof of Lemma 2.1. Part of this work was done when the third author visited the Freie University Berlin. He would like to thank the Humboldt Foundation for its generous support during this visit and the Freie University for its hospitality and stimulating research environment.

Appendix

Proof of Lemma 2.2. The proof uses the same ideas as in [18], Lemma 5.2 and also [11], Section 3.1.

Let D^* be a random orientation of G^* , where each edge chooses an orientation uniformly at random and independently of the other edges. We claim that with high probability, D^* satisfies the following.

- For every vertex v we have the following concentration on the indegrees and outdegrees:

$$c_0n/2 - n^{2/3} \leq \text{deg}_{in}(v) \leq c_0n/2 + n^{2/3},$$

$$c_0n/2 - n^{2/3} \leq \text{deg}_{out}(v) \leq c_0n/2 + n^{2/3}.$$

- For every two subsets $A, B \subset V$ such that $|A| \geq c_0n/3$ and $|B| \geq n/2$ there are at least $\gamma_0n^2/3$ edges from A to B , and at least $\gamma_0n^2/3$ edges from B to A .

This follows from a straightforward application of Chernoff’s inequality and the union bound.

Recall that

$$d = \left\lceil \left(\frac{c_0 - \varepsilon_0}{2} \right) n \right\rceil.$$

We construct a subdigraph of D^* whose indegrees and outdegrees are all equal to d . By forgetting the orientation of this subgraph we obtain the desired subgraph of G^* .

Consider the following flow network H . The vertex set is $\{s, t\} \cup X \cup Y$, where X and Y are both copies of $V(G^*)$. We add an edge (x, y) of capacity 1 for every edge (x, y) in D^* . Furthermore, we connect s to all vertices in X and also connect all vertices in Y to t using edges of capacity d . By the above discussion and the definition of d , note that the outdegrees of the vertices in X and the indegrees of the vertices in Y are all at least $c_0n/2n^{2/3} > d$ and at most $c_0n/2 + n^{2/3}$. Our aim is to show that there is a flow of value dn in H . Clearly, the edges used in such a flow will correspond to a subdigraph of D^* whose indegrees and outdegrees are all d , as desired.

By the Max-Flow-Min-Cut Theorem, it is sufficient to show that the capacity of every cut is at least dn . Let $C \subset V(H)$ such that $s \in C$ and $t \notin C$, and let $S = C \cap X$ and $T = C \cap Y$. The capacity of the cut defined by C is

$$d(n - |S|) + e(S, Y \setminus T) + d|T|,$$

which is at least dn if and only if

$$e(S, Y \setminus T) \geq d|S| - d|T|.$$

To show that this holds for any S and T , we must consider several cases. Note that we can assume that $|S| > |T|$, since otherwise $d|S| - d|T| \leq 0$ and we are done.

(1) $|S| < d$: In this case, we have

$$\begin{aligned} e(S, Y \setminus T) &= e(S, Y) - e(S, T) \\ &\geq (c_0n/2 - n^{2/3})|S| - |S||T| \\ &\geq d|S| - d|T|. \end{aligned}$$

(2) $|S| - |T| \geq \frac{4n^{2/3}}{\varepsilon_0}$: Here, we use the fact that d is substantially smaller than the indegrees and outdegrees of the vertices,

$$\begin{aligned} e(S, Y \setminus T) &\geq e(S, Y) - e(X, T) \\ &\geq (c_0n/2 - n^{2/3})|S| - (c_0n/2 + n^{2/3})|T| \\ &= d|S| - d|T| + \left(\frac{\varepsilon_0n}{2}\right)(|S| - |T|) - 2n^{2/3}|T| \\ &\geq d|S| - d|T|. \end{aligned}$$

(3) $|S| \leq (1 - c_0/3)n$: If $|T| \geq n/2$, since $|X \setminus S| \geq c_0n/3$, by the second property of D^* we have $e(X \setminus S, T) \geq \frac{\gamma_0n^2}{3}$. Otherwise, $|T| < n/2$. Since, after the above cases, we can assume that $|T| \geq |S| - \frac{4n^{2/3}}{\varepsilon_0}$ and $|S| \geq d$, we have that $|X \setminus S| \geq (1 - o(1))n/2$ and $T \geq c_0n/3$. Thus, again by the second property of D^* , we have $e(X \setminus S, T) \geq (1 - o(1))\frac{\gamma_0n^2}{3}$. Indeed, by adding $o(n)$ vertices to $X \setminus S$ we obtain a set of size at least $n/2$, and this can change the number of edges to T by at most $o(n^2)$. Hence, in both cases we have $e(X \setminus S, T) \geq (1 - o(1))\frac{\gamma_0n^2}{3}$, and therefore

$$\begin{aligned} e(S, Y \setminus T) &= e(S, Y) - e(X, T) + e(X \setminus S, T) \\ &\geq (c_0n/2 - n^{2/3})|S| - (c_0n/2 + n^{2/3})|T| + (1 - o(1))\frac{\gamma_0n^2}{3} \\ &\geq d|S| - d|T|. \end{aligned}$$

(4) $S > (1 - c_0/3)n$: Here, by the above we assume that $|T| \geq |S| - \frac{4n^{2/3}}{\varepsilon_0}$,

$$\begin{aligned} e(S, Y \setminus T) &= e(X, Y \setminus T) - e(X \setminus S, Y \setminus T) \\ &\geq d|Y \setminus T| - |X \setminus S||Y \setminus T| \\ &\geq d|Y \setminus T| - |X \setminus S|\left(\frac{c_0n}{3} + \frac{4n^{2/3}}{\varepsilon_0}\right) \\ &\geq d|Y \setminus T| - d|X \setminus S| \\ &= d|S| - d|T|. \quad \blacksquare \end{aligned}$$

References

- [1] N. Alon, *The maximum number of Hamiltonian paths in tournaments*, *Combinatorica* **10** (1990), 319–324.
- [2] B. Alspach, *The wonderful Walecki construction*, *Bulletin of the Institute of Combinatorics and its Applications* **52** (2008), 7–20.
- [3] B. Alspach, J. C. Bermond and D. Sotteau, *Decomposition into cycles I: Hamilton decompositions*, in: *Cycles and Rays (Montreal, PQ, 1987)*, NATO Advanced Science Institutes Series C: Mathematical and Physical Sciences, Vol. 301, Kluwer Academic Publishers, Dordrecht, 1990, pp. 9–18.
- [4] J. C. Bermond, O. Favaron and M. Maheo, *Hamiltonian decomposition of Cayley graphs of degree 4*, *Journal of Combinatorial Theory. Series B* **46** (1989), 142–153.
- [5] L. M. Brégman, *Certain properties of nonnegative matrices and their permanents*, *Doklady Akademii Nauk SSSR* **211** (1973), 27–30.
- [6] B. Csaba, A. Lo, D. Kühn, D. Osthus and A. Treglown, *Proof of the 1-factorization and Hamilton decomposition conjectures*, *Memoirs of the American Mathematical Society* **244** (2016).
- [7] B. Cuckler and J. Kahn, *Hamilton cycles in Dirac graphs*, *Combinatorica* **29** (2009), 299–326.
- [8] A. Ferber, M. Krivelevich and B. Sudakov, *Counting and packing Hamilton cycles in dense graphs and oriented graphs*, *Journal of Combinatorial Theory. Series B* **122** (2017), 196–220.
- [9] A. Fortuna, Z. Skupień and A. Žak, *Maximizing hamiltonian pairs and k -sets via numerous leaves in a tree*, *Discrete Mathematics* **309** (2009), 1788–1792.
- [10] A. Frieze, *On the number of perfect matchings and Hamilton cycles in ϵ -regular non-bipartite graphs*, *Electronic Journal of Combinatorics* **7** (2000), Research Paper 57.
- [11] A. Frieze and M. Krivelevich, *On packing Hamilton cycles in ϵ -regular graphs*, *Journal of Combinatorial Theory. Series B* **94** (2005), 159–172.
- [12] R. Glebov and M. Krivelevich, *On the number of Hamilton cycles in sparse random graphs*, *SIAM Journal on Discrete Mathematics* **27** (2013), 27–42.
- [13] S. Janson, *The numbers of spanning trees, Hamilton cycles and perfect matchings in a random graph*, *Combinatorics, Probability and Computing* **3** (1994), 97–126.
- [14] J. H. Kim and N. Wormald, *Random matchings which induce Hamilton cycles and Hamiltonian decompositions of random regular graphs*, *Journal of Combinatorial Theory. Series B* **81** (2001), 20–44.
- [15] F. Knox, D. Kühn and D. Osthus, *Edge-disjoint Hamilton cycles in random graphs*, *Random Structures & Algorithms* **46** (2015), 397–445.
- [16] M. Krivelevich, *On the number of Hamilton cycles in pseudo-random graphs*, *Electronic Journal of Combinatorics* **19** (2012), publication P25.
- [17] M. Krivelevich and B. Sudakov, *Pseudo-random graphs*, in *More Sets, Graphs and Numbers*, Bolyai Society Mathematical Studies, Vol. 15, Springer, Berlin, 2006, pp. 199–262.
- [18] D. Kühn and D. Osthus, *Hamilton decompositions of regular expanders: a proof of Kelly’s conjecture for large tournaments*, *Advances in Mathematics* **237** (2013), 62–146.
- [19] D. Kühn and D. Osthus, *Hamilton decompositions of regular expanders: applications*, *Journal of Combinatorial Theory. Series B* **104** (2014), 1–27.

- [20] L. Pósa, *Hamiltonian circuits in random graphs*, Discrete Mathematics **14** (1976), 359–364.
- [21] G. Ringel, *Über drei kombinatorische Probleme am n -dimensionalen Würfel und Würfelgitter*, Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg **20** (1955), 10–19.
- [22] B. Sudakov and V. Vu, *Local resilience of graphs*, Random Structures & Algorithms **33** (2008), 409–433.