

Directed Graphs Without Short Cycles

JACOB FOX^{1†}, PETER KEEVASH^{2‡} and BENNY SUDAKOV^{3§}

¹Department of Mathematics, Princeton, NJ 08544, USA
(e-mail: jacobfox@math.princeton.edu)

²School of Mathematical Sciences, Queen Mary, University of London,
Mile End Road, London E1 4NS, UK
(e-mail: p.keevash@qmul.ac.uk)

³Department of Mathematics, UCLA, Los Angeles, CA 90095, USA
(e-mail: bsudakov@math.ucla.edu)

Received 29 September 2008; first published online 1 October 2009

For a directed graph G without loops or parallel edges, let $\beta(G)$ denote the size of the smallest feedback arc set, *i.e.*, the smallest subset $X \subset E(G)$ such that $G \setminus X$ has no directed cycles. Let $\gamma(G)$ be the number of unordered pairs of vertices of G which are not adjacent. We prove that every directed graph whose shortest directed cycle has length at least $r \geq 4$ satisfies $\beta(G) \leq c\gamma(G)/r^2$, where c is an absolute constant. This is tight up to the constant factor and extends a result of Chudnovsky, Seymour and Sullivan.

This result can also be used to answer a question of Yuster concerning almost given length cycles in digraphs. We show that for any fixed $0 < \theta < 1/2$ and sufficiently large n , if G is a digraph with n vertices and $\beta(G) \geq \theta n^2$, then for any $0 \leq m \leq \theta n - o(n)$ it contains a directed cycle whose length is between m and $m + 6\theta^{-1/2}$. Moreover, there is a constant C such that either G contains directed cycles of every length between C and $\theta n - o(n)$ or it is close to a digraph G' with a simple structure: every strong component of G' is periodic. These results are also tight up to the constant factors.

1. Introduction

A digraph (directed graph) G is a pair (V_G, E_G) where V_G is a finite set of vertices and E_G is a set of ordered pairs (u, v) of vertices called edges. All digraphs we consider in this paper are simple, *i.e.*, they do not have loops or parallel edges. A path of length r in G is a collection of distinct vertices v_1, \dots, v_r together with edges (v_i, v_{i+1}) for $1 \leq i \leq r - 1$. Moreover, if (v_r, v_1) is also an edge, then it is an r -cycle.

[†] Research supported by an NSF Graduate Research Fellowship and a Princeton Centennial Fellowship.

[‡] Research supported in part by NSF grant DMS-0555755 and EPSRC grant EP/G056730/1.

[§] Research supported in part by NSF CAREER award DMS-0812005, and a USA–Israeli BSF grant.

The concept of cycles plays a fundamental role in graph theory, and there are numerous papers which study cycles in graphs. In contrast, the literature on cycles in directed graphs is not so extensive. It seems the main reason for this is that questions concerning cycles in directed graphs are often much more challenging than the corresponding questions in graphs. An excellent example of this difficulty is the well-known Caccetta–Häggkvist conjecture [4]. For $r \geq 2$, we say that a digraph is r -free if it does not contain a directed cycle of length at most r . The Caccetta–Häggkvist conjecture states that every r -free digraph on n vertices has a vertex of outdegree less than n/r . This notorious conjecture is still open even for $r = 3$, and we refer the interested reader to the recent surveys [11, 14], which discuss known results on this problem and other related open questions.

In approaching the Caccetta–Häggkvist conjecture it is natural to see what properties of an r -free digraph one can prove. A *feedback arc set* in a digraph is a collection of edges whose removal makes the digraph acyclic. For a digraph G , let $\beta(G)$ denote the size of the smallest feedback arc set. This parameter appears naturally in testing of electronic circuits and in efficient deadlock resolution (see, e.g., [10, 12]). It is also known that it is NP-hard to compute the minimum size of a feedback arc set even for tournaments [1, 5] (a tournament is an oriented complete graph). Let $\gamma(G)$ be the number of unordered pairs of vertices of G which are not adjacent. Chudnovsky, Seymour and Sullivan [7] conjectured that if G is a 3-free digraph then $\beta(G)$ is bounded from above by $\gamma(G)/2$. They proved this conjecture in two special cases, when the digraph is the union of two cliques or is a circular interval digraph. Moreover, for general 3-free digraphs G , they showed that $\beta(G) \leq \gamma(G)$.

Generalizing this conjecture, Sullivan [13] suggested that every r -free digraph G satisfies $\beta(G) \leq 2\gamma(G)/(r+1)(r-2)$, and gave an example showing that this would be best possible. She posed an open problem to prove that $\beta(G) \leq f(r)\gamma(G)$ for every r -free digraph G , for some function $f(r)$ tending to 0 as $r \rightarrow \infty$. Here we establish a stronger bound which shows that Sullivan’s conjecture is true up to a constant factor. This extends the result of Chudnovsky, Seymour and Sullivan to general r .

Theorem 1.1. *For $r \geq 3$, every r -free digraph G satisfies $\beta(G) \leq 800\gamma(G)/r^2$.*

The above result is tight up to a constant factor. Indeed, consider a blow-up of an $(r+1)$ -cycle, obtained by taking disjoint sets V_1, \dots, V_{r+1} of size $n/(r+1)$ and all edges from V_i to V_{i+1} , $1 \leq i \leq r+1$ (where $V_{r+2} = V_1$). This digraph on n vertices is clearly r -free, has

$$\gamma(G) = \binom{n}{2} - \frac{n^2}{r+1} \geq \frac{n(n-2)}{4}, \quad \text{and} \quad \beta(G) \geq \frac{n^2}{(r+1)^2}.$$

Indeed, G contains $\frac{n^2}{(r+1)^2}$ edge-disjoint cycles of length $r+1$, and one needs to delete at least one edge from each cycle to make G acyclic.

In order to prove Theorem 1.1, we obtain a bound on the edge expansion of r -free digraphs which may be of independent interest. For vertex subsets $S, T \subset V_G$, let $e_G(S, T)$ be the number of edges in G that go from S to T . The *edge expansion* $\mu(S)$ of a vertex

subset $S \subset V_G$ with cardinality $|S| \leq |V_G|/2$ is defined to be

$$\frac{1}{|S|} \min \{e_G(S, V_G \setminus S), e_G(V_G \setminus S, S)\}.$$

The edge expansion $\mu = \mu(G)$ of G is the minimum of $\mu(S)$ over all vertex subsets S of G with $|S| \leq |V_G|/2$. We show that r -free digraphs cannot have large edge expansion.

Theorem 1.2. *Suppose G is a digraph on n vertices, $r \geq 9$ and $\mu = \mu(G) \geq 25n/r^2$. Then every vertex of G is contained in a directed cycle of length at most r .*

Using this result, it is easy to deduce the following corollary, which implies Theorem 1.1 in the case when G is not too dense.

Corollary 1.3. *Every r -free digraph G on n vertices satisfies $\beta(G) \leq 25n^2/r^2$.*

Corollary 1.3 will also enable us to answer the following question posed by Yuster [15]. Suppose that a digraph G on n vertices is far from being acyclic, in that $\beta(G) \geq \theta n^2$. What lengths of directed cycles can we find in G ? Yuster [15] showed that for any $\theta > 0$ there exist constants K and η such that for any $m \in (0, \eta n)$ there is a directed cycle whose length is between m and $m + K$. He gave examples showing that one must have $K \geq \theta^{-1/2}$ and $\eta \leq 4\theta$, and posed the problem of determining the correct order of magnitude of these parameters as a function of θ . The following theorem, which is tight up to constant factors for both K and η , answers Yuster’s question.

Theorem 1.4. *For any $0 < \delta, \theta < 1$, the following holds for n sufficiently large. Suppose G is a digraph on n vertices with $\beta(G) \geq \theta n^2$. Then, for any $0 \leq m \leq (1 - \delta)\theta n$ there exists $m \leq \ell \leq m + (5 + \delta)\theta^{-1/2}$ such that G contains a directed cycle of length ℓ .*

Moreover, we can show that G either contains directed cycles of all lengths between some constant C and $\theta n - o(n)$ or is highly structured in the following sense. Say that G is *periodic* if the length of every directed cycle in G is divisible by some number $p \geq 2$, and *pseudoperiodic* if every strong component C is periodic (possibly with differing periods). A digraph is *strong* if, for every pair u, v of vertices, there is a path from u to v and a path from v to u . A *strong component* of a digraph G is a maximal strong subgraph of G . A pseudoperiodic digraph G is highly structured, as Theorem 10.5.1 of [3] shows that a strongly connected digraph with period p is contained in the blow-up of a p -cycle. Let $\lambda(G)$ denote the minimum number of edges of G that need to be deleted from G to obtain a pseudoperiodic digraph. Note that $\beta(G) \geq \lambda(G)$, as every acyclic digraph is pseudoperiodic.

Theorem 1.5. *For any $0 < \delta, \theta < 1$, there exist numbers C and n_0 such that the following holds for $n \geq n_0$. If G is a digraph on n vertices with $\lambda(G) \geq \theta n^2$, then G contains a directed cycle of length ℓ for any $C \leq \ell \leq (1 - \delta)\theta n$.*

The rest of this paper is organized as follows. In the next section we collect two simple lemmas concerning nearly complete digraphs. We need these lemmas in Section 3 to prove Theorems 1.1, 1.2 and Corollary 1.3. In Section 4, we discuss Szemerédi's Regularity Lemma for digraphs and some of its consequences. We use these results together with Corollary 1.3 in Section 5 to prove Theorems 1.4 and 1.5. The final section contains some concluding remarks.

Notation. An oriented graph is a digraph which can be obtained from a simple undirected graph by orienting its edges. Note that for $r \geq 2$, every r -free digraph is an oriented graph, as two opposite edges on the same pair of vertices form a 2-cycle. We write xy for an edge oriented from x to y , and say that the edge xy is incident to the vertices x and y . Suppose G is an oriented graph and S and T are subsets of its vertex set V_G . Let $E_G(S, T)$ be the set of edges in G that go from S to T , so $e_G(S, T) = |E_G(S, T)|$. We drop the subscript G if there is no danger of confusion. Let $G[S]$ denote the restriction of G to S , in which the vertex set is S and the edges are all those edges of G with both endpoints in S , and let $G \setminus S = G[V_G \setminus S]$ be the restriction of G to the complement of S . We use the notation $0 < \alpha \ll \beta$ to mean that there is a increasing function $f(x)$ so that the following argument is valid for $0 < \alpha < f(\beta)$. Throughout the paper, we systematically omit floor and ceiling signs whenever they are not crucial, for the sake of clarity of presentation. We also do not make any serious attempt to optimize absolute constants in our statements and proofs.

2. Basic facts

We start with two simple lemmas concerning oriented graphs that are nearly complete. First we prove a lemma which shows that such an oriented graph contains a vertex that has large indegree and large outdegree. Consider an oriented graph G whose vertex set is partitioned $V_G = V_1 \cup V_2$ with $|V_1| = |V_2| = n/2$, such that all edges go from V_1 to V_2 , and the restriction of G to each V_i is regular with indegree and outdegree of every vertex equal to $(1 - 2\epsilon)n/4$. The number of edges in G is $(1 - \epsilon)n^2/2$ and no vertex has indegree and outdegree both more than $(1 - 2\epsilon)n/4$. This example demonstrates tightness of the following lemma.

Lemma 2.1. *Let G be an oriented graph with n vertices and $(1 - \epsilon)n^2/2$ edges. Then G contains a vertex with indegree and outdegree at least $(1 - 2\epsilon)n/4$.*

Proof. Suppose for a contradiction that no vertex of G has indegree and outdegree at least $(1 - 2\epsilon)n/4$. Delete vertices one by one whose indegree and outdegree in the current oriented graph are both less than $(1 - 2\epsilon)n/4$. Let G' be the oriented graph that remains and let αn be the number of deleted vertices. Then G' has $(1 - \alpha)n$ vertices, at least $(1 - \epsilon)n^2/2 - \alpha n \cdot 2(1 - 2\epsilon)n/4$ edges, and every vertex has either indegree or outdegree at least $(1 - 2\epsilon)n/4$, but not both. Partition $V_{G'} = V_1 \cup V_2$, where V_1 consists of those vertices of G' that have indegree at least $(1 - 2\epsilon)n/4$. Since $|V_1| + |V_2| = (1 - \alpha)n$, we have $|V_1||V_2| \leq (1 - \alpha)^2 n^2/4$, and so

$$e(V_1) + e(V_2) \geq (1 - \epsilon)n^2/2 - (1 - 2\epsilon)\alpha n^2/2 - |V_1||V_2| \geq (1 - 2\epsilon + 4\alpha\epsilon - \alpha^2)n^2/4.$$

We may assume without loss of generality that $e(V_1)/e(V_2) \geq |V_1|/|V_2|$ (the other case can be treated similarly). In the first case,

$$\begin{aligned} e(V_1) &\geq \frac{|V_1|}{|V_1| + |V_2|} (e(V_1) + e(V_2)) \\ &\geq \frac{|V_1|}{|V_1| + |V_2|} (1 - 2\epsilon + 4\alpha\epsilon - \alpha^2) \frac{n^2}{4} \\ &= |V_1| (1 - 2\epsilon + 4\alpha\epsilon - \alpha^2) \frac{n}{4(1 - \alpha)}. \end{aligned}$$

Then the average outdegree of a vertex in V_1 is at least $(1 - 2\epsilon + 4\alpha\epsilon - \alpha^2) \frac{n}{4(1 - \alpha)}$. It is easy to check that as a function of α this is increasing for $\alpha \in [0, 1)$ and is therefore minimized when $\alpha = 0$. Therefore the average outdegree of a vertex in V_1 is at least $(1 - 2\epsilon)n/4$. Now we can choose a vertex in V_1 with outdegree at least the average, and then by definition of V_1 it has both indegree and outdegree at least $(1 - 2\epsilon)n/4$, a contradiction. \square

We can use this lemma to find in a nearly complete oriented graph a vertex of very large total degree and reasonably large indegree and outdegree.

Lemma 2.2. *Let G be an oriented graph with $n \geq 20$ vertices and $\gamma = \alpha n^2$ non-adjacent pairs, with $\alpha \leq 1/16$. Then G has a vertex v of total degree at least $(1 - 4\alpha)n$ and indegree and outdegree at least $n/10$.*

Proof. Let V' be those vertices of G with total degree at least $(1 - 4\alpha)n$. Then there are at least $|V \setminus V'|4\alpha n/2$ non-adjacent pairs with one or both vertices in $V \setminus V'$, so $(n - |V'|)2\alpha n \leq \gamma = \alpha n^2$, i.e., $|V'| \geq n/2$. Write $|V'| = \omega n$. The number of edges in the restriction $G[V']$ of G to V' is at least

$$\binom{|V'|}{2} - (\gamma - |V \setminus V'|4\alpha n/2) = (1 - (4\omega - 2)\alpha/\omega^2 - 1/|V'|)|V'|^2/2.$$

Applying Lemma 2.1 to $G[V']$, with $\epsilon = (4\omega - 2)\alpha/\omega^2 + 1/|V'|$, we find a vertex with indegree and outdegree at least

$$(1 - 2\epsilon)|V'|/4 = (1/4 - (2\omega - 1)\alpha/\omega^2)\omega n - 1/2 \geq n/8 - 1/2 \geq n/10,$$

where we use the fact that, for fixed $\alpha \leq 1/16$, the minimum of $f(\omega) = \omega/4 + (1 - 2\omega)\alpha/\omega$ for $\omega \in [1/2, 1]$ occurs at $\omega = 1/2$. Indeed, for $\omega \geq 1/2$, $f'(\omega) = 1/4 - \alpha/\omega^2 \geq 0$ and $f(\omega)$ is an increasing function. \square

3. Finding short cycles

We will prove Theorem 1.1 by proving that an r -free digraph cannot have large edge expansion. Recall that the edge expansion $\mu(S)$ of a set S of vertices of a digraph G with cardinality $|S| \leq |V_G|/2$ is defined to be

$$\frac{1}{|S|} \min \{e(S, V_G \setminus S), e(V_G \setminus S, S)\},$$

and the edge expansion $\mu = \mu(G)$ of G is the minimum of $\mu(S)$ over all subsets $S \subset V_G$ with $|S| \leq |V_G|/2$.

Consider a digraph G on n vertices and any vertex v of G . We say that a vertex w has *outdistance* i from v if the length of the shortest directed path from v to w is i . (*Indistance* is similarly defined.) Let N_i be the vertices at outdistance exactly i from v and let $M_i = \cup_{j \leq i} N_j$ be the vertices at outdistance at most i from v . It follows from these definitions that any edge from M_i to $V_G \setminus M_i$ is in fact an edge from N_i to N_{i+1} . We deduce that

$$\mu(M_i)|M_i| \leq e(M_i, V_G \setminus M_i) = e(N_i, N_{i+1}) \leq |N_i||N_{i+1}|.$$

Then the arithmetic–geometric mean inequality gives

$$|N_i| + |N_{i+1}| \geq 2\sqrt{\mu(M_i)|M_i|}. \tag{3.1}$$

The first step of the proof of Theorem 1.1 is Theorem 1.2, which shows that large edge expansion implies short cycles, and moreover we can find a short cycle through any specified vertex.

Proof of Theorem 1.2. Let v be any vertex of G . As before, let N_i be the vertices of outdistance exactly i from v and let M_i be the vertices of outdistance at most i from v . Also, let $a_i = (|N_i| + |N_{i+1}|)/\mu$ and $b_i = \sum_{1 \leq j \leq i} a_j$. Then $b_{i-1}\mu = 2|M_i| - |N_1| - |N_i| \leq 2|M_i|$, so dividing both sides of inequality (3.1) by μ and using $\mu(M_i) \geq \mu$ gives

$$a_i = (|N_i| + |N_{i+1}|)/\mu \geq 2\sqrt{\frac{\mu(M_i)|M_i|}{\mu}} \geq 2\sqrt{|M_i|/\mu} \geq \sqrt{2b_{i-1}}.$$

Adding b_{i-1} to both sides, we have $b_i \geq b_{i-1} + \sqrt{2b_{i-1}}$. Note that $b_1 = a_1 \geq |N_1|/\mu \geq 1$, as otherwise $|N_1| < \mu$, and taking $S = \{v\}$ we have $\mu(S) \leq |N_1| < \mu$, contradicting the definition of μ . Now we prove by induction that $b_i \geq \frac{2}{5}i^2$. This is easy to check for $i < 6$ using a calculator and $b_1 \geq 1$. For $i \geq 6$, the induction step is

$$b_i \geq b_{i-1} + \sqrt{2b_{i-1}} \geq \frac{2}{5}(i-1)^2 + \sqrt{4/5}(i-1) \geq \frac{2}{5}i^2.$$

Applying this with $i = \lfloor r/2 \rfloor$, we have $|M_i| \geq \mu b_{i-1}/2 \geq \mu(i-1)^2/5 > n/2$, since $\mu \geq 25n/r^2$ and $r \geq 9$. The same argument shows that there are more than $n/2$ vertices at indistance at most i from v . Therefore there is a vertex at indistance and outdistance at most i from v , which gives a directed cycle through v of length at most r . \square

Next we deduce Corollary 1.3, which implies our main theorem in the case when G is not almost complete.

Proof of Corollary 1.3. We suppose that G is r -free and prove that $\beta(G) \leq 25n^2/r^2$.

First we deal with the case $r \leq 10$. In any linear ordering of the vertices of G , deleting the forward edges or the backwards edges makes the digraph acyclic. Since the number of edges in G is $\binom{n}{2} - \gamma(G)$, we have $\beta(G) \leq \frac{1}{2}(\binom{n}{2} - \gamma(G)) < n^2/4$. Hence, $\beta(G) < 25n^2/r^2$ if $r \leq 10$.

Next, for $r \geq 11$ we use induction on n . Note that if $n \leq r$ then G is acyclic and $\beta(G) = 0$, so we can assume that $n > r$. By Theorem 1.2 and definition of μ we can find a set S with $|S| = s \leq n/2$ and $\mu(S) = \mu < 25n/r^2$. Note that a digraph formed by taking the disjoint union of two acyclic digraphs and adding some edges from the first acyclic digraph to the second acyclic digraph is acyclic. Therefore, using the inequality $n \leq 2(n - s)$, we obtain

$$\beta(G) \leq \beta(G[S]) + \beta(G \setminus S) + \mu s \leq 25s^2/r^2 + 25(n - s)^2/r^2 + 25n/r^2 \cdot s \leq 25n^2/r^2.$$

□

We need one more lemma before the proof of the main theorem, showing that an r -free oriented graph has a linear-sized subset S with small edge expansion.

Lemma 3.1. *Suppose $r \geq 15$, $0 \leq \alpha \leq 1/16$ and G is an r -free oriented graph on $n \geq 20$ vertices with $\gamma = \alpha n^2$ non-adjacent pairs. Then there exists $S \subset V(G)$ with $n/10 \leq |S| \leq n/2$ and $\mu(S) < 1500\alpha^2 n/r^2$.*

Proof. By Lemma 2.2 there is a vertex v of total degree at least $(1 - 4\alpha)n$ and indegree and outdegree at least $n/10$. As before, let N_i be the vertices of outdistance exactly i from v and let M_i be the vertices of outdistance at most i from v . Since G is r -free there is no vertex at indistance and outdistance at most $\lfloor r/2 \rfloor$ from v , so we can assume without loss of generality that $|M_i| \leq n/2$ for all $i \leq \lfloor r/2 \rfloor$. Also, by choice of v we have $|M_i| \geq |N_1| \geq n/10$, so we are done if we have $\mu(M_i) < 1500\alpha^2 n/r^2$ for some $i \leq \lfloor r/2 \rfloor$. Suppose for a contradiction that this is not the case. Then equation (3.1) gives

$$|N_i| + |N_{i+1}| \geq 2\sqrt{1500\alpha^2 n/r^2 \cdot n/10} > 24\alpha n/r.$$

Let $s = \lceil \frac{r-5}{4} \rceil \geq r/6$, so $2s + 1 \leq r/2$. The above inequality gives

$$|M_{2s+1}| - |N_1| = (|N_2| + |N_3|) + \dots + (|N_{2s}| + |N_{2s+1}|) > s \cdot 24\alpha n/r \geq 4\alpha n.$$

Let I_1 denote the inneighbourhood of v . By choice of v we have $|I_1| + |N_1| \geq (1 - 4\alpha)n$, and so $|I_1| + |M_{2s+1}| > n$, and hence there is a vertex in both I_1 and M_{1+2s} . This gives a cycle of length at most $2 + 2s \leq r$, contradiction. □

Proof of Theorem 1.1. We use induction on n to prove that every r -free digraph G on n vertices satisfies

$$\beta(G) \leq 800r^{-2}(\gamma(G) - \gamma(G)^2/n^2). \tag{3.2}$$

Note that the right-hand side of (3.2) is at least $400\gamma(G)/r^2$ and at most $800\gamma(G)/r^2$ as $0 \leq \gamma(G) \leq \binom{n}{2} \leq n^2/2$. We can assume that $\gamma(G) < n^2/16$, since otherwise we can apply Corollary 1.3 to get $\beta(G) \leq 25n^2/r^2 \leq 400\gamma(G)/r^2$. We can also assume that $r \geq 21$, as otherwise $r \leq 20$, and we can use the result of Chudnovsky, Seymour and Sullivan [7] that 3-free digraphs G satisfy $\beta(G) \leq \gamma(G) \leq 400\gamma(G)/r^2$. Then we can assume that $n \geq 22$, as otherwise $n \leq r$, G is acyclic, and $\beta(G) = 0$.

Let S be the set given by Lemma 3.1, $G_1 = G[S]$, $G_2 = G \setminus S$ and $n_i = |V(G_i)|$, $\gamma = \gamma(G)$, $\gamma_i = \gamma(G_i)$ for $i = 1, 2$, so that $n_1 + n_2 = n$ and $\gamma_+ := \gamma_1 + \gamma_2 \leq \gamma$. By choice of S we have

$\mu(S)|S| < 1600\gamma^2n_1/n^3r^2$. By deleting all edges from S to $V_G \setminus S$ or all edges from $V_G \setminus S$ to S , we get by the induction hypothesis that

$$\beta(G) \leq \beta(G_1) + \beta(G_2) + \mu(S)|S| \leq 800r^{-2}(\gamma_1 - \gamma_1^2/n_1^2 + \gamma_2 - \gamma_2^2/n_2^2 + 2\gamma^2n_1/n^3).$$

Now the Cauchy–Schwarz inequality gives

$$\gamma_+^2 = (n_1 \cdot \gamma_1/n_1 + n_2 \cdot \gamma_2/n_2)^2 \leq (n_1^2 + n_2^2)(\gamma_1^2/n_1^2 + \gamma_2^2/n_2^2),$$

so we have

$$\beta(G) \leq 800r^{-2}(\gamma_+ - \gamma_+^2/(n_1^2 + n_2^2) + 2\gamma^2n_1/n^3) \leq 800r^{-2}(\gamma - \gamma^2/(n_1^2 + n_2^2) + 2\gamma^2n_1/n^3).$$

Here we used $\gamma_+ \leq \gamma < n^2/16$ and $n_1^2 + n_2^2 \geq \frac{1}{2}(n_1 + n_2)^2 = n^2/2$, which give the inequality

$$\gamma - \frac{\gamma^2}{n_1^2 + n_2^2} - \gamma_+ + \frac{\gamma_+^2}{n_1^2 + n_2^2} = (\gamma - \gamma_+) \left(1 - \frac{\gamma_+ + \gamma}{n_1^2 + n_2^2} \right) \geq 0.$$

Now the desired bound on $\beta(G)$ follows from the inequality $\gamma^2/(n_1^2 + n_2^2) - 2\gamma^2n_1/n^3 \geq \gamma^2/n^2$. Set $n_1 = tn$, where $1/10 \leq t \leq 1/2$ by choice of S . It is required to show that $f(t) = \frac{1}{1+2t} - t^2 - (1-t)^2 \geq 0$. By computing $f'(t) = 2 - 4t - \frac{2}{(1+2t)^3}$ and $f''(t) = \frac{8}{(1+2t)^3} - 4$, we see that for $t \geq 0$, f'' is a decreasing function and $f''(0) > 0 > f''(1/2)$. Hence f' increases from $f'(0) = 0$ to a maximum and then decreases to $f'(1/2) < 0$, being first non-negative until some $t_0 < 1/2$ and then negative afterwards. Therefore, f increases from $f(0) = 0$ to a maximum $f(t_0)$ and then decreases to $f(1/2) = 0$, staying non-negative in the whole interval. This completes the proof. □

4. Regularity

For our second topic in the paper we will use the machinery of Szemerédi’s Regularity Lemma, which we will now describe. We will be quite brief, so for more details and motivation we refer the reader to the survey [9]. First we give some definitions. The density of a bipartite graph $G = (A, B)$ with vertex classes A and B is defined to be $d_G(A, B) := \frac{e_G(A, B)}{|A||B|}$. We write $d(A, B)$ if this is unambiguous. We also write $d_G(x)$, or simply $d(x)$, for the degree in G of a vertex x . Given $\epsilon > 0$, we say that G is ϵ -regular if, for all subsets $X \subseteq A$ and $Y \subseteq B$ with $|X| > \epsilon|A|$ and $|Y| > \epsilon|B|$, we have that $|d(X, Y) - d(A, B)| < \epsilon$. Given $d \in [0, 1]$, we say that G is (ϵ, d) -super-regular if it is ϵ -regular, and furthermore $d_G(a) \geq (d - \epsilon)|B|$ for all $a \in A$ and $d_G(b) \geq (d - \epsilon)|A|$ for all $b \in B$. If A and B are disjoint vertex subsets of a digraph G , we say that the pair $(A, B)_G$ is ϵ -regular if the bipartite graph with vertex sets A and B and edge set $E_G(A, B)$ is ϵ -regular. Similarly, we say that $(A, B)_G$ is (ϵ, d) -super-regular if the bipartite graph with vertex sets A and B and edge set $E_G(A, B)$ is (ϵ, d) -super-regular.

The Diregularity Lemma is a version of the Regularity Lemma for digraphs due to Alon and Shapira [2] (with a similar proof to the undirected version of Szemerédi).

Lemma 4.1 (Diregularity Lemma). *For every $\epsilon \in (0, 1)$ and $M' > 0$ there exist numbers M and n_0 such that, if G is a digraph on $n \geq n_0$ vertices, then there is a partition of the vertices of G into V_0, V_1, \dots, V_k for some $M' \leq k \leq M$ such that $|V_0| \leq \epsilon n$, $|V_1| = \dots = |V_k|$,*

and for all at but at most ek^2 ordered pairs $1 \leq i < j \leq k$ the underlying graph of $E_G(V_i, V_j)$ is ϵ -regular.

Given $0 \leq d \leq 1$ we define the reduced digraph R with parameters (ϵ, d) to have vertex set $[k] = \{1, \dots, k\}$ and an edge ij if and only if the underlying graph of $E_G(V_i, V_j)$ is ϵ -regular with density at least d . Note that if ϵ and d are small, M' is large, and G is a dense digraph, then most edges of G belong to pairs $E_G(V_i, V_j)$ for some edge $ij \in R$. Indeed, the exceptions are at most ϵn^2 edges incident to V_0 , at most n^2/M' edges lying within some V_i , at most ϵn^2 edges belonging to pairs $E_G(V_i, V_j)$ that are not ϵ -regular, and at most dn^2 edges belonging to $E_G(V_i, V_j)$ of density less than d : this gives a total less than $2dn^2$ if, say, $1/M' < \epsilon \ll d$. We also need the following path lemma.

Lemma 4.2. *For every $0 < d < 1$ there exists $\epsilon_0 > 0$ such that the following holds for $0 < \epsilon < \epsilon_0$. Let p, n be positive integers with $p \geq 4$, let U_1, \dots, U_p be pairwise disjoint sets of size n , and suppose G is a digraph on $U_1 \cup \dots \cup U_p$ such that each $(U_i, U_{i+1})_G$ is (ϵ, d) -super-regular. (Here, $U_{p+1} := U_1$.) Take any $x \in U_1$ and any $y \in U_p$. Then, for any $1 \leq \ell \leq n$ there is a path P in G of length $p\ell$, starting with x and ending with y , in which, for every vertex $v \in U_i$, the successor of v on P lies in U_{i+1} .*

This lemma can be easily deduced from the Blow-up Lemma of Komlós, Sarközy and Szemerédi (despite p being arbitrary), as shown in [6]. For the sake of completeness and the convenience of the reader we include the proof here. In fact, for our purposes it is sufficient to apply the result with $1 \leq \ell \leq (1 - \epsilon)n$; in that case it is not too hard to prove it directly with a random embedding procedure, but we omit the details. Note also that by applying the lemma when yx is an edge we can obtain a directed cycle of length $p\ell$ for any $1 \leq \ell \leq n$.

The requirement that $p \geq 4$ in Lemma 4.2 is necessary. Indeed, if $p = 2$ or $p = 3$, there may not be a path of length p from x to y . It is not difficult to show using Lemma 4.2 that, even in this case, we can find a path from x to y of length $p\ell$ for all $2 \leq \ell \leq n$. It is even easier to show that we can greedily find such paths for all $2 \leq \ell \leq dn/2$, and since this will be sufficient for our purposes, we do so now. In the following argument, if i does not satisfy $1 \leq i \leq p$, then we define $U_i := U_j$ with $1 \leq j \leq p$ and $i \equiv j \pmod{p}$. Since each pair $(U_i, U_{i+1})_G$ is (ϵ, d) -super-regular, each vertex in U_i has at least $(d - \epsilon)n$ outneighbours in U_{i+1} , and we can greedily find a path $P' = v_1 \dots v_{p\ell-3}$ with starting point $v_1 = x$ and with each v_i in U_i , as each such path only contains at most $\ell \leq dn/2$ vertices in each U_i . Let X be the outneighbours of $v_{p\ell-3}$ in $U_{p-2} \setminus P'$ and let Y be the inneighbours of y in $U_{p-1} \setminus P'$, so $|X| \geq (d - \epsilon)n - \ell \geq (\frac{d}{2} - \epsilon)n \geq \epsilon n$ and similarly $|Y| \geq \epsilon n$. Since the pair $(U_{p-2}, U_{p-1})_G$ is (ϵ, d) -super-regular, then there is at least one edge $(v_{p\ell-2}, v_{p\ell-1})$ from X to Y and $v_1 \dots v_{p\ell}$ for which $v_{p\ell} = y$ is the desired path P from x to y of length $p\ell$.

We start the proof of Lemma 4.2 by recalling the Blow-up Lemma of Komlós, Sárközy and Szemerédi [8].

Lemma 4.3. *Given a graph R of order k and parameters $d, \Delta > 0$, there exists an $\eta_0 = \eta_0(d, \Delta, k) > 0$ such that whenever $0 < \eta \leq \eta_0$, the following holds. Let V_1, \dots, V_k be disjoint*

sets and let R^* be the graph on $V_1 \cup \dots \cup V_k$ obtained by replacing each edge ij of R by the complete bipartite graph between V_i and V_j . Let G be a spanning subgraph of R^* such that, for each edge ij of R , the bipartite subgraph of G consisting of all edges between V_i and V_j is (η, d) -super-regular. Then G contains a copy of every subgraph H of R^* with maximum degree $\Delta(H) \leq \Delta$. Moreover, this copy of H in G maps the vertices of H to the same sets V_i as the copy of H in R^* , i.e., if $h \in V(H)$ is mapped to V_i by the copy of H in R^* , then it is also mapped to V_i by the copy of H in G .

From the Blow-up Lemma, we can quickly deduce the following lemma.

Lemma 4.4. *For every $0 < d < 1$ there exists $\epsilon_0 > 0$ such that the following holds for $0 < \epsilon < \epsilon_0$. Suppose $p \geq 4$, let U_1, \dots, U_p be pairwise disjoint sets of size n , for some n , and suppose G is a graph on $U_1 \cup \dots \cup U_p$ such that each pair (U_i, U_{i+1}) , $1 \leq i \leq p - 1$ is (ϵ, d) -super-regular. Let $f : U_1 \rightarrow U_p$ be any bijective map. Then there are n vertex-disjoint paths from U_1 to U_p such that, for every $x \in U_1$, the path starting from x ends at $f(x) \in U_p$.*

Proof. Choose a sequence $1 = i_1 < i_2 < \dots < i_t = p$ such that $3 \leq i_j - i_{j-1} \leq 5$ for $2 \leq j \leq t$. Let $f_j : U_{i_{j-1}} \rightarrow U_{i_j}$ be any bijective maps with $f = f_t \circ \dots \circ f_2$. Let G_j be the graph obtained from the restriction of G to $U_{i_{j-1}} \cup U_{i_{j-1}+1} \cup \dots \cup U_{i_j}$ by identifying each vertex $x \in U_{i_{j-1}}$ with $f_j(x) \in U_{i_j}$. By Lemma 4.3 we can find n vertex-disjoint cycles in G_j of length $i_j - i_{j-1}$, provided that $\epsilon_0 < \eta(d, 2, i_j - i_{j-1})$, which only depends on d as $i_j - i_{j-1} \leq 5$. These n cycles correspond to n vertex-disjoint paths in G from $U_{i_{j-1}}$ to U_{i_j} , such that, for every $x \in U_{i_{j-1}}$, the path starting from x ends at $f_j(x) \in U_{i_j}$. By concatenating these paths, we get the desired n vertex-disjoint paths from U_1 to U_p so that, for every $x \in U_1$, the path starting from x ends at $f(x) \in U_p$. \square

Now we give the proof of Lemma 4.2.

Proof of Lemma 4.2. Suppose G is a digraph on $U_1 \cup \dots \cup U_p$, where $|U_i| = n$, $1 \leq i \leq p$, such that each $(U_i, U_{i+1})_G$ is (ϵ, d) -super-regular, with $\epsilon < \epsilon_0$ given by Lemma 4.4. Suppose also $x \in U_1$, $y \in U_p$ and $1 \leq \ell \leq n$. We need to find a path P of length $p\ell$ from x to y . First we apply the Blow-up Lemma to find a perfect matching from $U_p \setminus y$ to $U_1 \setminus x$. We label U_1 as $\{x_1, \dots, x_n\}$ and U_p as $\{y_1, \dots, y_n\}$ with $x_1 = x$ and $y_1 = y$, so that the matching edges go from y_i to x_i for $2 \leq i \leq n$. Then we apply Lemma 4.4 to find n vertex-disjoint paths from U_1 to U_p so that the path P_i starting at x_i ends at y_{i+1} for $1 \leq i \leq \ell - 1$ and the path P_ℓ starting at x_ℓ ends at $y_1 = y$ (the other paths can be arbitrary). Now our required path P is $x_1 P_1 y_2 x_2 P_2 y_3 \dots x_\ell P_\ell y_1$. \square

We finish the section with two simple lemmas concerning super-regularity. The first lemma tells us that large induced subgraphs of super-regular bipartite graphs are also super-regular.

Lemma 4.5. *Let G be a bipartite graph with parts A and B that is (ϵ, d) -super-regular, $\epsilon < 1/2 \leq \alpha < 1$, $A' \subset A$ and $B' \subset B$ with $|A'|/|A|, |B'|/|B| \geq \alpha$, and let G' be the induced subgraph of G with parts A' and B' . Then G' is $(2\epsilon, d - 1 + \alpha)$ -super-regular.*

Proof. Super-regularity of G implies that each vertex $a \in A' \subset A$ satisfies $d_G(a) \geq (d - \epsilon)|B|$. Hence,

$$\begin{aligned} d_{G'}(a) &\geq d_G(a) - (|B| - |B'|) \geq (d - \epsilon)|B| - (|B| - |B'|) \\ &\geq (d - (1 - \alpha) - \epsilon)|B| \geq (d - (1 - \alpha) - \epsilon)|B'|. \end{aligned}$$

Likewise, each vertex $b \in B'$ satisfies $d_{G'}(b) \geq (d - (1 - \alpha) - \epsilon)|B'|$.

Let $X \subset A'$ and $Y \subset B'$ with $|X| > 2\epsilon|A'|$ and $|Y| > 2\epsilon|B'|$. Since $1/2 \leq \alpha \leq |A'|/|A|, |B'|/|B|$ we have $|X| > \epsilon|A|$ and $|Y| > \epsilon|B|$. Now the pair $(A, B)_G$ is ϵ -regular, so $|d(X, Y) - d(A, B)| < \epsilon$, and the triangle inequality gives

$$|d(X, Y) - d(A', B')| \leq |d(X, Y) - d(A, B)| + |d(A, B) - d(A', B')| < 2\epsilon.$$

Hence, G' is $(2\epsilon, d - 1 + \alpha)$ -super-regular. □

For any bounded degree subgraph H of a reduced graph R , the next lemma allows us to make the pairs $(V_i, V_j)_G$ corresponding to edges ij of H super-regular by deleting a few vertices from each V_i .

Lemma 4.6. *Suppose R is the reduced digraph with parameters (ϵ, d) of a Szemerédi partition $V_G = V_0 \cup V_1 \cup \dots \cup V_k$ of a digraph G and H is a subdigraph of R with maximum total degree at most Δ , where $\Delta \leq \frac{1}{2\epsilon}$. Then, for each $i, 1 \leq i \leq k$, there exists $U_i \subset V_i$ with $|U_i| = (1 - \Delta\epsilon)|V_i|$ such that, for each edge ij of H , the pair $(U_i, U_j)_G$ is $(2\epsilon, d - \Delta\epsilon)$ -super-regular.*

Proof. For each edge ij of H , delete all vertices in V_i with fewer than $(d - \epsilon)|V_j|$ outneighbours in V_j and all vertices in V_j with fewer than $(d - \epsilon)|V_i|$ inneighbours in V_i . For each edge ij of H , fewer than $\epsilon|V_i|$ elements are deleted from V_i and fewer than $\epsilon|V_j|$ elements are deleted from V_j . Indeed, if the subset $S \subset V_i$ of vertices with fewer than $(d - \epsilon)|V_j|$ outneighbours in V_j has cardinality $|S| \geq \epsilon|V_i|$, then $d_G(S, V_j) < d - \epsilon$, in contradiction to ij being an edge of the reduced graph R . Likewise, at most $\epsilon|V_j|$ elements are deleted from V_j for each edge ij . Hence, in total, at most $\Delta\epsilon|V_i|$ vertices are deleted from each V_i . Delete further vertices from each V_i until the resulting subset U_i has cardinality $(1 - \Delta\epsilon)|V_i|$. For each edge ij of H , each vertex in U_i has at least $(d - \epsilon)|V_j|$ outneighbours in V_j and hence at least

$$(d - \epsilon)|V_j| - (|V_j| - |U_j|) = (d - (\Delta + 1)\epsilon)|V_j| \geq (d - (\Delta + 1)\epsilon)|U_j|$$

outneighbours in U_j . Similarly, for each edge ij of H , each vertex in V_j has at least $(d - (\Delta + 1)\epsilon)|U_i|$ inneighbours in U_i . Letting $\alpha = |U_i|/|V_i| = 1 - \Delta\epsilon$, we have $\alpha \geq 1/2$. For each edge ij of H , since $(V_i, V_j)_G$ is ϵ -regular, Lemma 4.5 implies that $(U_i, U_j)_G$ is 2ϵ -regular and hence is $(2\epsilon, d - \Delta\epsilon)$ -super-regular. □

5. Cycles of almost given length

Now we will apply the Regularity Lemma and Corollary 1.3 to answer the question of Yuster mentioned in the Introduction.

Proof of Theorem 1.4. Choose parameters $0 < 1/n_0 \ll 1/M \ll \epsilon \ll d \ll \delta, \theta$ and $M' = \epsilon^{-1}$. Suppose G is a digraph on $n \geq n_0$ vertices with $\beta(G) \geq \theta n^2$. Note that $\theta < 1/2$ as in any linear ordering of the vertices of G , deleting all the forward edges or all the backward edges yields an acyclic digraph. Apply Lemma 4.1 to obtain a partition of the vertices of G into V_0, V_1, \dots, V_k for some $M' \leq k \leq M$ and let R be the reduced digraph on $[k]$ with parameters (ϵ, d) . As noted in the previous section, there are at most $2dn^2$ edges of G that do not belong to $E_G(V_i, V_j)$ for some edge $ij \in R$. We can make G acyclic by deleting these edges and at most $\beta(R)(n/k)^2$ edges corresponding to edges of R , so we must have $\beta(R) \geq (\theta - 2d)k^2$. Let S_1, \dots, S_g be the strong components of R and suppose $\beta(S_i) = \theta_i |S_i|^2$. Then

$$\sum_{i=1}^g |S_i| = k \quad \text{and} \quad \sum_{i=1}^g \theta_i |S_i|^2 = \sum_{i=1}^g \beta(S_i) = \beta(R) \geq (\theta - 2d)k^2.$$

It follows that we can choose some S_j with $\theta_j |S_j| \geq (\theta - 2d)k$ (otherwise we would have $\sum_{i=1}^g \theta_i |S_i|^2 < (\theta - 2d)k \sum |S_i| = (\theta - 2d)k^2$).

Next we restrict our attention to S_j and repeatedly delete any vertex with outdegree less than $\theta_j |S_j|$ in S_j . We must arrive at some digraph R_0 on $k_0 \leq |S_j|$ vertices with minimum outdegree at least $\theta_j |S_j| \geq (\theta - 2d)k$ and $\beta(R_0) \geq \theta_j |S_j| k_0$. Indeed, otherwise we could make S_j acyclic by deleting fewer than $\theta_j |S_j| k_0 + (|S_j| - k_0) \theta_j |S_j| = \theta_j |S_j|^2$ edges, which is impossible. Let $C = c_1 \cdots c_p$ be a directed cycle in R_0 of length $p \geq (\theta - 2d)k$. It can be found by considering a longest directed path and using the fact that the end of the path has at least $(\theta - 2d)k$ outneighbours, which all lie on the path. Recall that

$$\beta(S_j) = \theta_j |S_j|^2 \geq (\theta - 2d)k |S_j| \geq (\theta - 2d) |S_j|^2.$$

By Corollary 1.3, if S_j is r -free, then $(\theta - 2d) |S_j|^2 \leq \beta(S_j) \leq 25 |S_j|^2 / r^2$, so

$$r \leq 5(\theta - 2d)^{-1/2} < (5 + \delta)\theta^{-1/2},$$

where we use $d \ll \delta, \theta$. Therefore, there is a directed cycle $C' = c'_1 \cdots c'_r$ in S_j of length r for some $2 \leq r \leq (5 + \delta)\theta^{-1/2}$ (which may intersect C in an arbitrary fashion). Also, by strong connectivity of S_j we can find a directed path Q_1 from c_p to c'_r and a directed path Q_2 from c'_r to c_1 . Suppose that the lengths of these paths are q_1 and q_2 , respectively. We note that $q_1, q_2 \leq k$.

Let H denote the digraph with vertex set V_{S_j} and edge set $E_C \cup E_{C'} \cup E_{Q_1} \cup E_{Q_2}$. Note that the maximum total degree of H is at most 8, as each path and cycle has maximum total degree at most 2. By Lemma 4.6, for each vertex i of S_j there exists $U_i \subset V_i$ with $|U_i| = (1 - 8\epsilon) |V_i|$ such that, for each edge ij of H , the pair $(U_i, U_j)_G$ is $(2\epsilon, d - 8\epsilon)$ -super-regular.

Suppose $0 \leq m \leq (1 - \delta)\theta n$ is given. We give separate arguments depending on whether the cycles we seek in G are short or long. First consider the case $m < 3k$. Choose ℓ

divisible by r with $m \leq \ell < m + r$. Then we can find a cycle of length ℓ within the classes U_i corresponding to C' , as noted after Lemma 4.2. (This argument holds as long as $r \geq 4$ or $\ell \geq 2r$. If otherwise, then $\ell = r \in \{2, 3\}$ and we can find a cycle of length $2r$ in G . This $2r$ -cycle completes this case as $m \leq \ell = r \leq 2r \leq 6 < 5\theta^{-1/2}$, where we use $\theta < 1/2$.) Now suppose $m \geq 3k$ and write $m = q_1 + q_2 + sp + t$, with $0 \leq t < p$ and $1 \leq s < (1 - \delta/2)n/k$ (since $p \geq (\theta - 2d)k$). The integer t is indeed non-negative since $q_1, q_2, p \leq k$ and $m \geq 3k$. We can choose $\ell = q_1 + q_2 + sp + u$, where $u < p + r$ is a multiple of r and $m \leq \ell < m + r$. Say that a path $P = v_1 \cdots v_e$ in G corresponds to a walk $W = w_1 \cdots w_e$ in R if every edge $v_i v_{i+1}$, $1 \leq i \leq e - 1$ of P goes from U_{w_i} to $U_{w_{i+1}}$. For ij an edge of H , the pair $(U_i, U_j)_G$ is $(2\epsilon, d - 8\epsilon)$ -super-regular, so any vertex in U_i has at least $(d - 10\epsilon)|U_j|$ outneighbours in U_j . Therefore, we can greedily find:

- (1) a directed path P_1 in G corresponding to Q_1 in R , starting at some $y \in U_{c_p}$ and ending at some $z \in U_{c_r}$,
- (2) a directed path P_2 in G corresponding to u/r copies of C' in R , starting at z and ending at some other $z' \in U_{c_r}$, and avoiding P_1 ,
- (3) a directed path P_3 in G corresponding to Q_2 in R , starting at z' and ending at some $x \in U_{c_1}$, avoiding $P_1 \cup P_2$.

Let P be the path $P_1 P_2 P_3$. Note that P has at most $u/r + 2$ vertices in each U_i . As we next find a path from x to y disjoint from $P \setminus \{x, y\}$, we delete the vertices of $P \setminus \{x, y\}$ and also at most $u/r + 2$ vertices from each U_i so that they all still have the same size, letting U'_i be the resulting subset of U_i . Now

$$|U'_i| \geq |U_i| - (u/r + 2) \geq (1 - 8\epsilon)|V_i| - (u/r + 2) > (1 - d/2)|V_i| > (1 - \delta/2)(n/k) = s.$$

This also gives $|U'_i|/|U_i| > 1 - d/2$ for each vertex i of S_j . For each edge ij of H , (U_i, U_j) is $(2\epsilon, d - 8\epsilon)$ -super-regular. Hence, Lemma 4.5 with $\alpha = 1 - d/2$ implies that (U'_i, U'_j) is $(4\epsilon, d/4)$ super-regular, as $d - 8\epsilon - d/2 = d/2 - 8\epsilon \geq d/4$. Therefore, we can apply Lemma 4.2 with $U_i = U'_{c_i}$, $1 \leq i \leq p$ to obtain a directed path from x to y of length sp . Combining this with the path P already found from y to x gives a directed cycle of length ℓ , as required. □

For the proof of Theorem 1.5 we need the following two facts from elementary number theory.

Chinese Remainder Theorem. Suppose x_1, \dots, x_t are integers with greatest common factor 1. Then any integer n can be expressed as $n = a_1 x_1 + \dots + a_t x_t$ with integers a_1, \dots, a_t .

Sylvester’s ‘coin problem’. Suppose x and y are coprime positive integers. Then every integer $n \geq (x - 1)(y - 1)$ can be represented as $n = ax + by$ with a, b non-negative integers.

Proof of Theorem 1.5. It is straightforward to see that λ (similarly to β) is additive on strong components, i.e., if a digraph G has strong components T_1, \dots, T_g , then $\lambda(G) = \sum_{i=1}^g \lambda(T_i)$. Also, $\lambda(G) \leq \beta(G)$, since every acyclic digraph is pseudoperiodic. Therefore we start as in the proof of Theorem 1.4 by applying Lemma 4.1 to obtain a partition of the

vertices of G into V_0, V_1, \dots, V_k for some $M' \leq k \leq M$ and letting R be the reduced digraph on $[k]$ with parameters (ϵ, d) . As before, G has at most $2dn^2$ edges not corresponding to edges of the reduced digraph R , so we must have $\lambda(R) \geq (\theta - 2d)k^2$. Then, as in the proof of Theorem 1.4, we find a strong component S_j of R with $\beta(S_j) \geq \lambda(S_j) = \theta_j |S_j|^2$ and $\theta_j |S_j| \geq (\theta - 2d)k$, directed cycles $C = c_1 \cdots c_p$ and $C' = c'_1 \cdots c'_r$ in S_j with $p \geq (\theta - 2d)k$ and $2 \leq r \leq (5 + \delta)\theta^{-1/2}$, a directed path Q_1 from c_p to c'_r of length $q_1 \leq k$ and a directed path Q_2 from c'_r to c_1 of length $q_2 \leq k$.

Next we show how to construct a closed walk W in S_j starting and ending at c'_r with length $l(W) = w$ coprime to r . Since S_j is not f -periodic for any $f \geq 2$, for each prime factor f of r there is a directed cycle with length not divisible by f . Therefore we can choose cycles D_1, \dots, D_r so that $l(C'), l(D_1), \dots, l(D_r)$ have greatest common factor 1. Fix vertices $d_i \in D_i$, $1 \leq i \leq r$ and choose the following directed paths in S_j (which exist by strong connectivity): Q'_1 from c'_r to d_1 and Q''_1 from d_1 to c'_r , Q'_i from d_{i-1} to d_i and Q''_i from d_i to d_{i-1} for $2 \leq i \leq r$. Let W' be the walk $Q'_1 \cdots Q'_r Q''_r \cdots Q''_1$. By the Chinese Remainder Theorem we can find integers a_1, \dots, a_r such that $l(W') + a_1 l(D_1) + \cdots + a_r l(D_r) \equiv 1 \pmod r$. By reducing mod r we can assume that $0 \leq a_i \leq r - 1$ for $1 \leq i \leq r$. We let W be the walk obtained from W' by including a_i copies of D_i when d_i is first visited. That is, we obtain W by walking along W' , and, for $1 \leq i \leq r$, when we first reach d_i , before we continue onto the next vertex, we first walk a_i times around the cycle D_i . Then $l(W) \equiv 1 \pmod r$ is coprime to r . The walk W visits any vertex at most $2r^2$ times. Indeed, each of the $2r$ directed paths Q'_i and Q''_i visit each vertex at most once, and each time we go around cycle D_i adds at most one new visit to any vertex, so W visits each vertex at most $2r + a_1 + \cdots + a_r \leq 2r + r^2 \leq 2r^2$ times. As S_j has at most k vertices and visits each vertex at most $2r^2$ times, $w = l(W) \leq 2r^2 k$.

Let H be the digraph with vertex set V_{S_j} and edge set $E_C \cup E_{C'} \cup E_W$. Since W visits any vertex at most $2r^2$ times, each vertex in W is in at most $4r^2$ edges of H . Therefore, H has maximum total degree at most $4 + 4r^2 \leq 8r^2$. By Lemma 4.6, for each vertex i of S_j there exists $U_i \subset V_i$ with $|U_i| = (1 - 8r^2\epsilon)|V_i|$ such that, for each edge ij of H , the pair $(U_i, U_j)_G$ is $(2\epsilon, d - 8r^2\epsilon)$ -super-regular.

Fix any ℓ with $500\theta^{-3/2}M \leq \ell \leq (1 - \delta)\theta n$. We will show that G contains a directed cycle of length ℓ . As $2 \leq r < 6\theta^{-1/2}$, $p, q_1, q_2 \leq k \leq M$ and $w \leq 2r^2 k$, we have

$$\ell \geq 500\theta^{-3/2}M \geq 3k + 2r^3 k \geq q_1 + q_2 + p + rw.$$

Therefore, we can write $\ell = q_1 + q_2 + sp + u$, with $rw \leq u < rw + p$ and $1 \leq s < (1 - \delta/2)n/k$ (the last inequality uses $p \geq (\theta - 2d)k$). Since r, w are coprime, by the ‘coin problem’ result of Sylvester we can write $u = ar + bw$ with a, b non-negative integers. We have $a \leq u/r < w + p \leq 2r^2 k + k$ and $b \leq u/w < r + p \leq 2k$. For ij an edge of H , the pair $(U_i, U_j)_G$ is $(2\epsilon, d - 8r^2\epsilon)$ -super-regular, so any vertex in U_i has at least $(d - 10r^2\epsilon)|U_j|$ outneighbours in U_j . Therefore, we can greedily find:

- (1) a directed path P_1 in G corresponding to Q_1 in R , starting at some $y \in U_{c_p}$ and ending at some $z \in U_{c'_r}$,
- (2) a directed path P_2 in G corresponding to a copies of C' in R , starting at z and ending at some other $z' \in U_{c'_r}$, and avoiding P_1 ,

- (3) a directed path P_3 in G corresponding to b copies of W in R , starting at z' and ending at some other $z'' \in U_{c_r}$, and avoiding $P_1 \cup P_2$,
- (4) a directed path P_4 in G corresponding to Q_2 in R , starting at z'' and ending at some $x \in U_{c_1}$, avoiding $P_1 \cup P_2 \cup P_3$.

Let P be the path $P_1P_2P_3P_4$. As we walk along path P , for each i , the number of times U_i is visited is at most once for P_1 , at most a times for P_2 , at most $b \cdot 2r^2$ times for P_3 , and at most once for P_4 . Therefore, for each i ,

$$|P \cap U_i| \leq 1 + a + b \cdot 2r^2 + 1 \leq 1 + 2r^2k + k + 2k \cdot 2r^2 + 1 \leq 10r^2k.$$

We delete the vertices of $P \setminus \{x, y\}$ as we next find a directed path from x to y that is disjoint from $P \setminus \{x, y\}$. We further delete at most $10r^2k$ vertices from each U_i so that they all still have the same size, and let U'_i be the resulting subset of U_i . Now

$$|U'_i| \geq |U_i| - 10r^2k = (1 - 8r^2\epsilon)|V_i| - 10r^2k > (1 - d/2)|V_i| > (1 - \delta/2)(n/k) = s.$$

Then $|U'_i|/|U_i| > (1 - d/2)$, and Lemma 4.5 with $\alpha = 1 - d/2$ implies that each pair $(U'_i, U'_j)_G$ with ij an edge of H is $(4\epsilon, d/4)$ -super-regular, as $d - 8r^2\epsilon - d/2 \geq d/4$. Therefore we can apply Lemma 4.2 with $U_i = U'_{c_i}$, $1 \leq i \leq p$ to obtain a directed path from x to y of length sp . Combining this with the path P already found from y to x gives a directed cycle of length ℓ , as required. □

6. Concluding remarks

(a) We have not presented the best possible constants that come from our methods, opting to give reasonable constants that can be obtained with relatively clean proofs. With more work one can replace the constant 25 in Theorem 1.2, and so in Corollary 1.3, by a constant that approaches 8 as r becomes large. However, Sullivan [13] conjectures that the correct constant is 2, and it would be interesting to close this gap. The problems of estimating β and μ are roughly equivalent: we used the bound on μ from Theorem 1.2 to establish the bound on β in Theorem 1.3. Conversely, if we delete $\beta(G)$ edges from G to make it acyclic, order the vertices so that all remaining edges point in one direction and take S to be the first $n/2$ vertices in the ordering, we see that

$$\mu(G)(n/2) = \mu(G)|S| \leq \mu(S)|S| = \min(e(S, V_G \setminus S), e(V_G \setminus S, S)) \leq \beta(G),$$

so a bound on β gives a bound on μ . However, these arguments may be too crude to give the correct constants.

(b) Applying this better constant 8 (mentioned above) in Corollary 1.3, we can replace the constant 5 by 3 (say) in Theorem 1.4, so that the parameter K in Yuster's question (the length of the interval where we look for a cycle length) is determined up to a factor of 3. The parameter η (the maximum length of a cycle as a proportion of n) is determined up to a factor of about 4 if the question is posed for oriented graphs, or a factor of 2 if the question is posed for digraphs. Indeed, Yuster shows that $\eta \leq 4\theta$ for oriented graphs by taking $1/4\theta$ copies of a random regular tournament on $4\theta n$ vertices; for digraphs one can show $\eta \leq 2\theta$ by taking $1/2\theta$ copies of the complete digraph on $2\theta n$ vertices. We can

find longer cycles in a periodic digraph G on n vertices with $\beta(G) \geq \theta n^2$, but θn is still the correct bound up to a constant of about 2, as may be seen from the blow-up of a 2-cycle with parts of size $(1 + 2\theta)\theta n$ and $(1 - (1 + 2\theta)\theta)n$.

(c) If a digraph G is far from being acyclic but we can obtain a pseudoperiodic digraph G' by deleting few edges of G , then some strong component of G' has small period. More precisely, if $\beta(G) \geq \theta n^2$ and we can obtain a pseudoperiodic G' by deleting at most δn^2 edges from G , then some strong component of G' must have period at most $(\theta - \delta)^{-1/2}$. To see this, note that $\beta(G') \geq (\theta - \delta)n^2$, so some strong component H of G' satisfies $\beta(H) \geq (\theta - \delta)m^2$, where $m = |V_H|$. Since G' is pseudoperiodic, H is p -periodic, for some p , so is contained in the blow-up of a p -cycle, i.e., the vertex set of H can be partitioned as $V(H) = V_1 \cup \dots \cup V_p$, so that every edge goes from V_i to V_{i+1} , for some $1 \leq i \leq p$, writing $V_{p+1} = V_1$. (For a proof see Theorem 10.5.1 in [3].) Write $t_i = |V_i|/m$. Then there is some $1 \leq i \leq p$ for which $t_i t_{i+1} \leq 1/p^2$. This can be seen from the arithmetic–geometric mean inequality: we have

$$1 = \sum_{i=1}^p t_i \geq p \prod_{i=1}^p t_i^{1/p} = p \prod_{i=1}^p (t_i t_{i+1})^{1/2p},$$

so

$$\prod_{i=1}^p t_i t_{i+1} \leq (1/p^2)^p.$$

It follows that $\beta(H) \leq (m/p)^2$, i.e., $p \leq (\theta - \delta)^{-1/2}$, as required.

(d) The dependence of C on θ which we get in Theorem 1.5 is quite poor, since the proof uses Szemerédi's Regularity Lemma and the value of C depends on the number of parts in the regular partition. It would be interesting to determine the right dependence of C on θ . One should note that we obtained good constants in the proof of Theorem 1.4 despite using the Regularity Lemma, so it may not be necessary to avoid its use.

References

- [1] Alon, N. (2006) Ranking tournaments. *SIAM J. Discrete Math.* **20** 137–142.
- [2] Alon, N. and Shapira, A. (2004) Testing subgraphs in directed graphs. *J. Comput. System Sci.* **69** 354–382.
- [3] Bang-Jensen, J. and Gutin, G. (2001) *Digraphs: Theory, Algorithms and Applications*, Springer.
- [4] Caccetta, L. and Häggkvist, R. (1978) On minimal digraphs with given girth. In *Proc. 9th Southeastern Conference on Combinatorics, Graph Theory, and Computing* (Boca Raton 1978), *Congress. Numer.* **XXI** 181–187.
- [5] Charbit, P., Thomassé, S. and Yeo, A. (2007) The minimum feedback arc set problem is NP-hard for tournaments. *Combin. Probab. Comput.* **16** 1–4.
- [6] Christofides, D., Keevash, P., Kühn, D. and Osthus, D. Finding Hamilton cycles in robustly expanding digraphs. Submitted.
- [7] Chudnovsky, M., Seymour, P. and Sullivan, B. (2008) Cycles in dense digraphs. *Combinatorica* **28** 1–18.
- [8] Komlós, J., Sárközy, G. N. and Szemerédi, E. (1997) Blow-up lemma. *Combinatorica* **17** 109–123.

- [9] Komlós, J. and Simonovits, M. (1996) Szemerédi's regularity lemma and its applications in graph theory. In *Combinatorics: Paul Erdős is Eighty*, Vol. 2 (D. Miklós, V. T. Sós and T. Szőnyi, eds), Vol. 2 of *Bolyai Society Mathematical Studies*, Budapest, pp. 295–352.
- [10] Leiserson, C. E. and Saxe, J. B. (1991) Retiming synchronous circuitry. *Algorithmica* **6** 5–35.
- [11] Nathanson, M. B. The Caccetta–Häggkvist conjecture and additive number theory. AIM Preprint 2006-10: www.aimath.org/preprints.html.
- [12] Shaw, A. C. (1974) *The Logical Design of Operating Systems*, Prentice-Hall, Englewood Cliffs, NJ.
- [13] Sullivan, B. (2008) Extremal problems in digraphs. PhD thesis, Princeton University.
- [14] Sullivan, B. A summary of results and problems related to the Caccetta–Häggkvist conjecture. AIM Preprint 2006-13: www.aimath.org/preprints.html.
- [15] Yuster, R. (2008) Almost given length cycles in digraphs. *Graphs Combin.* **24** 59–65.