



ELSEVIER

Contents lists available at ScienceDirect

Linear Algebra and its Applications

www.elsevier.com/locate/laa



Flattening rank and its combinatorial applications



David Munhá Correia, Benny Sudakov*, István Tomon

ETH Zurich, Switzerland

ARTICLE INFO

Article history:

Received 8 March 2021

Accepted 5 May 2021

Available online 11 May 2021

Submitted by R. Brualdi

MSC:

05D40

05C50

05C70

05C35

15A69

Keywords:

Tensor

Rank

Algebraic methods

Extremal problems

ABSTRACT

Given a d -dimensional tensor $T : A_1 \times \cdots \times A_d \rightarrow \mathbb{F}$ (where \mathbb{F} is a field), the i -flattening rank of T is the rank of the matrix whose rows are indexed by A_i , columns are indexed by $B_i = A_1 \times \cdots \times A_{i-1} \times A_{i+1} \times \cdots \times A_d$ and whose entries are given by the corresponding values of T . The *max-flattening rank* of T is defined as $\text{mfrank}(T) = \max_{i \in [d]} \text{frank}_i(T)$. A tensor $T : A^d \rightarrow \mathbb{F}$ is called *semi-diagonal*, if $T(a, \dots, a) \neq 0$ for every $a \in A$, and $T(a_1, \dots, a_d) = 0$ for every $a_1, \dots, a_d \in A$ that are all distinct. In this paper we prove that if $T : A^d \rightarrow \mathbb{F}$ is semi-diagonal, then $\text{mfrank}(T) \geq \frac{|A|}{d-1}$, and this bound is the best possible.

We give several applications of this result, including a generalization of the celebrated Frankl-Wilson theorem on forbidden intersections. Also, addressing a conjecture of Aharoni and Berger, we show that if the edges of an r -uniform multi-hypergraph \mathcal{H} are colored with z colors such that each color class is a matching of size t , then \mathcal{H} contains a rainbow matching of size t provided $z > (t-1) \binom{r-t}{r}$. This improves previous results of Alon and Glebov, Sudakov, and Szabó.

© 2021 Elsevier Inc. All rights reserved.

* Corresponding author.

E-mail addresses: david.munhacanascorreia@math.ethz.ch (D. Munhá Correia), benjamin.sudakov@math.ethz.ch (B. Sudakov), istvan.tomon@math.ethz.ch (I. Tomon).

1. Introduction

A d -dimensional tensor over a field \mathbb{F} is a function $T : A_1 \times \cdots \times A_d \rightarrow \mathbb{F}$, where A_1, \dots, A_d are finite sets. For $i \in [d]$, the i -flattening rank of T , denoted by $\text{frank}_i(T)$, is defined as follows. Let $B_i = A_1 \times \cdots \times A_{i-1} \times A_{i+1} \times \cdots \times A_d$, and view T as the matrix T_i whose rows are indexed by A_i , and columns indexed by B_i . Then $\text{frank}_i(T) := \text{rank}(T_i)$. Note that $\text{frank}_i(T) = 1$ if and only if $T \neq 0$, and there exist two functions $f : A_i \rightarrow \mathbb{F}$ and $g : B_i \rightarrow \mathbb{F}$ such that $T(a_1, \dots, a_d) = f(a_i)g(a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_d)$. Also, the i -flattening rank of T is the minimum r such that T is the sum of r tensors of i -flattening rank 1. Equivalently, $\text{frank}_i(T)$ is the dimension of the vector space generated by the rows of T in the i -th dimension, which are also the columns of the matrix T_i . Define the *max-flattening rank* of T as

$$\text{mfrank}(T) = \max_{i \in [d]} \text{frank}_i(T).$$

It is easy to see that the max-flattening rank and i -flattening rank satisfy the usual properties of rank. More precisely, they are subadditive, and if T' is a subtensor of T , then $\text{frank}_i(T') \leq \text{frank}_i(T)$ and $\text{mfrank}(T') \leq \text{mfrank}(T)$. Here, $T' : A'_1 \times \cdots \times A'_r \rightarrow \mathbb{F}$ is a *subtensor* of $T : A_1 \times \cdots \times A_r \rightarrow \mathbb{F}$ if $A'_i \subset A_i$ for $i \in [r]$, and $T'(a_1, \dots, a_r) = T(a_1, \dots, a_r)$ for $(a_1, \dots, a_r) \in A'_1 \times \cdots \times A'_r$. Also, the usual notion of tensor rank is always an upper bound for the max-flattening rank. As a reminder, T has *tensor rank* 1 if there are d functions f_1, \dots, f_d such that $T(a_1, \dots, a_d) = f_1(a_1) \dots f_d(a_d)$, and the tensor rank $\text{trank}(T)$ is the minimal r such that T is the sum of r tensors of tensor rank 1.

In this paper, we are interested in combinatorial applications of the max-flattening rank. Note that one of the trivial, but important properties of the matrix rank is that diagonal matrices have full rank. The analogue of this is also trivially true for the flattening rank: if $T : A^d \rightarrow \mathbb{F}$ is a diagonal tensor, that is, $T(a_1, \dots, a_d) \neq 0$ if and only if $a_1 = \cdots = a_d$, then $\text{frank}_i(T) = \text{mfrank}(T) = |A|$ for $i \in [d]$. However, in certain applications this is not really what is needed, thus we would like to relax the notion of diagonality.

Say that d -dimensional tensor $T : A^d \rightarrow \mathbb{F}$ is *semi-diagonal* if the following holds: $T(a_1, \dots, a_d) = 0$ if a_1, \dots, a_d are all distinct, and $T(a_1, \dots, a_d) \neq 0$ if $a_1 = \cdots = a_d$. If a_1, \dots, a_d are neither all equal or all distinct, then there is no restriction on $T(a_1, \dots, a_d)$. Our main technical result is the following lower bound on the rank of semi-diagonal tensors.

Theorem 1. *Let $T : A^d \rightarrow \mathbb{F}$ be a d -dimensional semi-diagonal tensor. Then*

$$\text{mfrank}(T) \geq \frac{|A|}{d - 1}.$$

Let us make a few remarks about this theorem. The bound $\text{mfrank}(T) \geq \left\lceil \frac{|A|}{d-1} \right\rceil$ is the best possible for any positive integers $d \geq 2$ and $|A|$. Indeed, let A_1, \dots, A_m form a partition of A into $m = \left\lceil \frac{|A|}{d-1} \right\rceil$ parts of size at most $d - 1$, and define the tensor $T : A^d \rightarrow \mathbb{F}$ such that

$$T(a_1, \dots, a_d) = \begin{cases} 1 & \text{if } a_1, \dots, a_d \in A_i \text{ for some } i \in [m] \\ 0 & \text{otherwise.} \end{cases}$$

Then T is semi-diagonal, and the i -flattening rank of T is exactly m for $i \in [d]$. To see this, for $j \in [m]$, let $v_j : A \rightarrow \mathbb{F}$ be the vector defined as $v_j(a) = 1$ if $a \in A_j$, and $v_j(a) = 0$ otherwise. Then for $i \in [d]$, each column of the flattened matrix T_i is equal to one of the vectors v_j , and these vectors generate an m -dimensional space.

Moreover, the i -flattening rank of a semi-diagonal tensor need not be large for any fixed i . Indeed, if $T : A^d \rightarrow \mathbb{F}$ is defined as

$$T(a_1, \dots, a_d) = \begin{cases} 1 & \text{if } a_1 = \dots = a_{i-1} = a_{i+1} = \dots = a_d \\ 0 & \text{otherwise,} \end{cases}$$

then $\text{frank}_i(T) = 1$. This is true as each column of T_i is either the constant 0 or the constant 1 vector.

We will prove Theorem 1 in the next section and provide some of its combinatorial applications in Section 3. We would also like to mention that further application of Theorem 1 appears in [12], where it is used to establish certain Ramsey properties of algebraic hypergraphs.

2. Semi-diagonal tensors

In this section, we prove Theorem 1. More precisely, we prove the following theorem, which then immediately implies Theorem 1.

Theorem 2. *Let $T : A^d \rightarrow \mathbb{F}$ be a semi-diagonal tensor. Then*

$$\sum_{i=1}^d \text{frank}_i(T) \geq \frac{d}{d-1} |A|.$$

Proof. Let us introduce some notation. If $\mathbf{v} \in \mathbb{F}^A$, let $\text{supp}(\mathbf{v})$ be the *support* of \mathbf{v} , that is, the set of elements $b \in A$ such that $\mathbf{v}(b) \neq 0$. If $\mathbf{a} \in A^d$, let $\{\mathbf{a}\} = \{\mathbf{a}(1), \dots, \mathbf{a}(d)\} \subset A$ be the set of elements of A which appear as coordinates of the vector \mathbf{a} (no confusion will arise in this notation with the singleton set containing \mathbf{a}). Let $\phi(\mathbf{a}) \subset [d]$ be the set of indices $i \in [d]$ such that $\mathbf{a}(i)$ appears at least twice among $\mathbf{a}(1), \dots, \mathbf{a}(d)$. Also, for $i \in [d]$, let $\mathbf{a}[i] \in \mathbb{F}^A$ be the vector defined as

$$\mathbf{a}[i](b) = T(\mathbf{a}(1), \dots, \mathbf{a}(i - 1), b, \mathbf{a}(i + 1), \dots, \mathbf{a}(d))$$

for $b \in A$. Finally, let $U_i(T) = \{\mathbf{a}[i] : \mathbf{a} \in A^d\}$, and let $V_i(T)$ be the subspace of \mathbb{F}^A generated by the elements of $U_i(T)$. Note that $U_i(T)$ is the set of columns of the flattened matrix T_i , so, by definition,

$$\text{frank}_i(T) = \dim(V_i(T)).$$

We prove the theorem by induction on $|A|$. If $|A| \leq d - 1$, the statement is clearly true as the i -flattening is at least 1 for $i \in [d]$, so let us assume that $|A| \geq d$. Choose $\mathbf{a} \in A^d$ such that $T(\mathbf{a}) \neq 0$, and the set $\{\mathbf{a}\}$ has maximal size. Then $|\{\mathbf{a}\}| \leq d - 1$ as T is semi-diagonal.

Claim 3. *If $i \in \phi(\mathbf{a})$, then $\text{supp}(\mathbf{a}[i]) \subset \{\mathbf{a}\}$.*

Proof. Suppose this is not the case, and let $c \in \text{supp}(\mathbf{a}[i]) \setminus \{\mathbf{a}\}$. Let \mathbf{a}' be the d -tuple we get by replacing $\mathbf{a}(i)$ with c in \mathbf{a} . Then $T(\mathbf{a}') \neq 0$ and $\{\mathbf{a}'\} = \{\mathbf{a}\} \cup \{c\}$, contradicting the maximality of $|\{\mathbf{a}\}|$. \square

Let \mathcal{A} be the set of all d -tuples $\mathbf{b} \in A^d$ such that $T(\mathbf{b}) \neq 0$ and $\{\mathbf{b}\} = \{\mathbf{a}\}$. Define the graph G on \mathcal{A} as follows: connect \mathbf{b} and \mathbf{b}' by an edge if they differ in exactly one coordinate. Let $\mathcal{C} \subset \mathcal{A}$ be the connected component of G containing \mathbf{a} . Say that an index $j \in [d]$ is *good* if there exists $\mathbf{b} \in \mathcal{C}$ such that $j \in \phi(\mathbf{b})$, and let $J \subset [d]$ be the set of good indices.

Claim 4. *If j is not good, then $\mathbf{a}(j) = \mathbf{b}(j)$ for every $\mathbf{b} \in \mathcal{C}$.*

Proof. As \mathbf{a} and \mathbf{b} are in the same connected component, there exists a path from \mathbf{a} to \mathbf{b} in G , which means that there exists a sequence $\mathbf{a} = \mathbf{a}_0, \dots, \mathbf{a}_p = \mathbf{b}$ of elements of \mathcal{A} such that \mathbf{a}_k and \mathbf{a}_{k+1} differ in exactly one coordinate for $k = 0, \dots, p - 1$, say in coordinate j_k . But as $\{\mathbf{a}_k\} = \{\mathbf{a}_{k+1}\}$, we have that $\mathbf{a}_k(j_k)$ is equal to some other coordinate $\mathbf{a}_k(j')$. Therefore, j_k is good, so $j \neq j_k$ and the coordinate at j was never changed. \square

Let $X = \{\mathbf{a}(j) : j \in [d] \setminus J\}$. By definition of goodness, all $\mathbf{a}(j)$ are distinct elements of A and appear only once as a coordinate of \mathbf{a} . Thus $|X| = d - |J|$. Also, let $Y = \{\mathbf{a}\} \setminus X$, then

$$1 \leq |Y| = |\{\mathbf{a}\}| - |X| \leq d - 1 - |X| = |J| - 1,$$

where we used the inequality $|\{\mathbf{a}\}| \leq d - 1$, which holds by the semi-diagonality of T . For every $j \in J$, pick an element $\mathbf{b} \in \mathcal{C} \subset \mathcal{A}$ such that $j \in \phi(\mathbf{b})$, and let $\mathbf{v}_j = \mathbf{b}[j] \in V_j(T)$.

Claim 5. $\text{supp}(\mathbf{v}_j) \subset Y$.

Proof. As $j \in \phi(\mathbf{b})$, we have by Claim 3 that $\text{supp}(\mathbf{v}_j) \subset \{\mathbf{b}\} = \{\mathbf{a}\}$. Since $Y = \{\mathbf{a}\} \setminus X = \{\mathbf{b}\} \setminus X$, if $\text{supp}(\mathbf{v}_j) \not\subset Y$, then there exists $c \in X$ such that $c \in \text{supp}(\mathbf{v}_j)$. Then $c \neq \mathbf{b}(j)$ as $c \in X$. Let \mathbf{b}' be the d -tuple we get after replacing $\mathbf{b}(j)$ with c in \mathbf{b} . Since $c \in \{\mathbf{b}\}$, we have that $\{\mathbf{b}'\} = \{\mathbf{b}\} = \{\mathbf{a}\}$ and \mathbf{b}' differs from \mathbf{b} in exactly one coordinate. Then $\mathbf{b}' \in \mathcal{C}$ but $\mathbf{b}'(j) \neq \mathbf{b}(j)$, contradicting Claim 4. \square

Let $A' = A \setminus Y$ and let T' be the restriction of T to $(A')^d$. Then T' is an $(|A| - |Y|) \times \dots \times (|A| - |Y|)$ -sized semi-diagonal tensor, so by our induction hypothesis, we have

$$\sum_{i=1}^d \dim(V_i(T')) \geq \frac{d}{d-1}(|A| - |Y|).$$

However, by Claim 5, the support of the vector \mathbf{v}_j is disjoint from A' for $j \in J$. Let $\mathbf{e}_1, \dots, \mathbf{e}_r \in (A')^d$ such that the restriction of the vectors $\mathbf{e}_1[j], \dots, \mathbf{e}_r[j]$ to A' is a basis of $V_j(T')$ (so $r = \dim(V_j(T'))$), then the vectors $\mathbf{v}_j, \mathbf{e}_1[j], \dots, \mathbf{e}_r[j]$ are linearly independent in \mathbb{F}^A . Therefore,

$$\dim(V_j(T)) \geq \dim(V_j(T')) + 1.$$

But then

$$\sum_{i=1}^d \dim(V_i(T)) \geq |J| + \sum_{i=1}^d \dim(V_i(T')) \geq |J| + \frac{d}{d-1}(|A| - |Y|) \geq \frac{d}{d-1}|A|,$$

where the last inequality holds noting that $|Y| \leq |J| - 1 \leq d - 1$. \square

3. Applications

3.1. Oddtown

A family \mathcal{A} of subsets of an n -element set is an *Oddtown* if $|A|$ is odd for every $A \in \mathcal{A}$, and $|A \cap B|$ is even for every $A, B \in \mathcal{A}, A \neq B$. It was proved famously by Berkelamp [5] that the size of an Oddtown on an n element ground set is at most n .

The following generalization of this problem was considered by Vu [13]. A *d -wise Oddtown* is a family \mathcal{A} of subsets of some ground set such that $|A|$ is odd for every $A \in \mathcal{A}$, and $|A_1 \cap \dots \cap A_d|$ is even for every $A_1, \dots, A_d \in \mathcal{A}$ that are all distinct. Vu [13] proved that the size of a d -wise Oddtown on an n element ground set is at most $(d - 1)n$. If we allow repetitions in \mathcal{A} this bound is also the best possible.

As our first application, we show that the cross-version of this result also holds with the same bound. A *cross- d -wise Oddtown* is a family \mathcal{A} such that every $A \in \mathcal{A}$ is an ordered d -tuple $A(1), \dots, A(d)$ of subsets of the ground set (which do not need to be distinct) with the property that $|A(1) \cap \dots \cap A(d)|$ is odd for every $A \in \mathcal{A}$, and $|A_1(1) \cap \dots \cap A_d(d)|$

is even for every $A_1, \dots, A_d \in \mathcal{A}$ that are all distinct. Note that d -wise Oddtown is a special case of cross- d -wise Oddtown, where every d -tuple contains the same set d times.

Theorem 6. *If \mathcal{A} is a cross- d -wise Oddtown on an n element ground set, then $|\mathcal{A}| \leq (d - 1)n$.*

Proof. Let $m = |\mathcal{A}|$. Define the tensor $T : \mathcal{A}^d \rightarrow \mathbb{F}_2$ such that for $A_1, \dots, A_d \in \mathcal{A}$ we have $T(A_1, \dots, A_d) = |A_1(1) \cap \dots \cap A_d(d)|$. Then \mathcal{A} is a semi-diagonal tensor (as we are working over a field of characteristic 2), so we have $\text{mfrank}(T) \geq \frac{m}{d-1}$ by Theorem 1. On the other hand, we show that $\text{trank}(T) \leq n$, which then implies $\text{mfrank}(T) \leq n$. For $A \in \mathcal{A}$ and $j \in [d]$, let $v_{A(j)} : [n] \rightarrow \mathbb{F}_2$ be the characteristic function of $A(j)$. For $k \in [n]$, define the function $f_{j,k} : \mathcal{A} \rightarrow \mathbb{F}_2$ as $f_{j,k}(A) = v_{A(j)}(k)$. Then

$$T(A_1, \dots, A_d) = \sum_{k=1}^n f_{1,k}(A_1) \dots f_{d,k}(A_d).$$

Therefore, $\text{trank}(T) \leq n$. \square

3.2. Forbidden intersections

Let p be a prime, $L \subset \mathbb{F}_p$ be a set of residues mod p , and $\mathcal{F} \subset 2^{[n]}$ be a family such that $|A| \notin L$ for every $A \in \mathcal{F}$, but $|A \cap B| \in L$ for every distinct $A, B \in \mathcal{F}$. The celebrated Frankl-Wilson theorem [7] on forbidden intersections says that

$$|\mathcal{F}| \leq \sum_{s=0}^{|L|} \binom{n}{s}.$$

A natural extension of this was given by Grolmusz and Sudakov [10], who proved that if one requires that $|A| \notin L$ for every $A \in \mathcal{F}$, but every intersection of k distinct sets in \mathcal{F} has size in L , then $|\mathcal{F}| \leq (k - 1) \sum_{s=0}^{|L|} \binom{n}{s}$. This bound is tight if we allow \mathcal{F} to be a multiset. Here, we prove the following extension of these results for more general, forbidden configurations. A *configuration of order k modulo p* is a pair (\mathcal{C}, L) , where $\mathcal{C} \subset 2^{[k]}$ and $L \subset \mathbb{F}_p$. Say that a family $\mathcal{F} \subset 2^{[n]}$ is (\mathcal{C}, L) -satisfying, if $|A| \notin L$ for every $A \in \mathcal{F}$, but there exist no k distinct sets $A_1, \dots, A_k \in \mathcal{F}$ such that $|\bigcap_{i \in X} A_i| \notin L$ for every $X \in \mathcal{C}$.

Clearly, asking \mathcal{F} to be $(\{\{1, 2\}\}, L)$ -satisfying is equivalent to the condition of the Frankl-Wilson theorem and being $(\{\{1, \dots, k\}\}, L)$ -satisfying is equivalent to restricted k -wise intersections. We bound the size of the maximal (\mathcal{C}, L) -satisfying family by a function of n , $|L|$ and the maximum degree of \mathcal{C} , which we define next. Given a family \mathcal{C} , the *degree* of $a \in [k]$ in \mathcal{C} is the number of sets in \mathcal{C} containing a , and is denoted by $\text{deg}_{\mathcal{C}}(a)$. The *maximum degree* of \mathcal{C} is $\Delta(\mathcal{C}) = \max_{a \in [k]} \text{deg}_{\mathcal{C}}(a)$.

Theorem 7. *Let (C, L) be a configuration of order k modulo p , and let $\Delta = \Delta(C)$. If $\mathcal{F} \subset 2^{[n]}$ is (C, L) -satisfying, then*

$$|\mathcal{F}| \leq (k - 1) \sum_{s=0}^{\Delta|L|} \binom{n}{s}.$$

Proof. Let $h : \mathbb{F}_p \rightarrow \mathbb{F}_p$ be the polynomial defined as $h(x) = \prod_{\ell \in L} (x - \ell)$. Define the k -dimensional tensor $T : \mathcal{F}^k \rightarrow \mathbb{F}_p$ as follows. For A_1, \dots, A_k , let

$$T(A_1, \dots, A_k) = \prod_{X \in \mathcal{C}} h \left(\left| \bigcap_{i \in X} A_i \right| \right).$$

Then T is semi-diagonal as \mathcal{F} is (C, L) -satisfying. Therefore, by Theorem 1, we have

$$\text{mfrank}(T) \geq \frac{|\mathcal{F}|}{k - 1}.$$

We show that for $j \in [k]$, the j -flattening rank of T is at most $|\mathcal{F}| \leq \sum_{s=0}^{d_j} \binom{n}{s}$, where $d_j = \deg_{\mathcal{C}}(j)|L|$. For ease of notation, let us show this for $j = 1$, it follows for the other values of j by the same reasoning.

For $A \in \mathcal{F}$, let $v_A \in \mathbb{F}_p^n$ be the characteristic vector of A . Let $A_1, \dots, A_k \in \mathcal{F}$, then

$$T(A_1, \dots, A_k) = \prod_{X \in \mathcal{C}} h \left(\sum_{j=1}^n \prod_{i \in X} v_{A_i}(j) \right).$$

Let $p : \mathbb{F}_p^{kn} \rightarrow \mathbb{F}_p$ be the polynomial defined as

$$p(v_1, \dots, v_k) = \prod_{X \in \mathcal{C}} h \left(\sum_{j=1}^n \prod_{i \in X} v_i(j) \right),$$

where $v_1, \dots, v_k \in \mathbb{F}^n$. Write p as the sum of monomials and in each monomial replace $v_i(j)^a, a \geq 1$ by $v_i(j)$. Let q be the resulting polynomial and note that $q(v_1, \dots, v_k) = p(v_1, \dots, v_k)$ if v_1, \dots, v_k are characteristic vectors, having all their coordinates equal to 0 or 1.

The polynomial $q(v_1, \dots, v_k)$ can be written as the sum of polynomials of the form

$$\beta_J \left(\prod_{j=1}^n v_1(j)^{J(j)} \right) q_J(v_2, \dots, v_k),$$

where $J \in \{0, 1\}^n$, $\beta_J \in \mathbb{F}_p$ and $q_J : \mathbb{F}_p^{(k-1)n} \rightarrow \mathbb{F}_p$ is some polynomial. Note that $\beta_J = 0$ unless $|J| \leq |L|\deg_{\mathcal{C}}(1)$. Let $\mathcal{J} = \{J \in \{0, 1\}^n : |J| \leq |L|\deg_{\mathcal{C}}(1)\}$. For $J \in \mathcal{J}$, define the functions $f_J : \mathcal{F} \rightarrow \mathbb{F}_p$ and $g_J : \mathcal{F}^{k-1} \rightarrow \mathbb{F}_p$ as

$$f_J(A_1) = \beta_J \prod_{j=1}^k v_{A_1}(j)^{J(j)},$$

and

$$g_J(A_2, \dots, A_k) = q_J(v_{A_2}, \dots, v_{A_k}).$$

Then

$$T(A_1, \dots, A_k) = \sum_{J \in \mathcal{J}} f_J(A_1)g_J(A_2, \dots, A_k),$$

which proves that $\text{frank}_1(T) \leq |\mathcal{J}| \leq \sum_{s=0}^{d_1} \binom{n}{s}$. As the corresponding bound holds for the j -flattening as well for $j \in [k]$, we get

$$\text{mfrank}(T) \leq \sum_{s=0}^{\Delta(\mathcal{C})|L|} \binom{n}{s}.$$

Comparing the lower and upper bound on the max-flattening rank, we get the desired bound

$$|\mathcal{F}| \leq (k - 1) \sum_{s=0}^{\Delta(\mathcal{C})|L|} \binom{n}{s}. \quad \square$$

So far we showed that for a fixed configuration (\mathcal{C}, L) of order k modulo p , the maximal size of a (\mathcal{C}, L) -satisfying family of subsets of $[n]$ is of order at most $n^{\Delta|L|}$, where Δ is the maximum degree of the set family \mathcal{C} . For $\Delta = 1$ the exponent of n is clearly best possible, as the Frankl-Wilson bound is sharp. On the other hand, for $\Delta \geq 2$ we do not know how accurate our result is. Nevertheless, we can show that the exponent of n must depend on Δ .

Indeed, consider the case $p = 2$, $\mathcal{C}_k = [k]^{(2)}$ is a complete graph of order k and $L = \{0\}$. Then $\Delta = k - 1$ and we construct families in $2^{[n]}$ which are $(\mathcal{C}_k, \{0\})$ -satisfying and have size $n^{\Omega(\log k / \log \log k)}$. Our construction is a modification of an argument of Alon and Szegedy [3].

Theorem 8. *Let t, s be positive integers, $k = \lfloor \frac{2^{t+1}}{t-1} \rfloor$, and $n = t^s$. Then, there exists a family $\mathcal{F} \subseteq 2^{[n]}$ which is $(\mathcal{C}_k, \{0\})$ -satisfying and has size at least $2^{(t-1)s/4}$.*

Proof. Let $\mathcal{G} \subseteq 2^{[t]}$ be a family of odd-sized sets, all whose pairwise intersections have also odd size. By the well known variation of the Oddtown problem, we have $|\mathcal{G}| \leq 2^{(t-1)/2}$.

Now let $\mathcal{O}_t \subseteq 2^{[t]}$ denote the family of all odd-sized subsets of $[t]$, which clearly has size 2^{t-1} . Since $n = t^s$, we can identify $[n]$ with the set $[t]^s$. Let $\mathcal{O}_t^s \subseteq 2^{[n]}$ denote the family of sets of the form $A_1 \times \dots \times A_s$, where $A_i \in \mathcal{O}_t$ for all i . Note that all sets in \mathcal{O}_t^s

are odd-sized. If $\mathcal{F}_1, \dots, \mathcal{F}_s$ are subsets of \mathcal{O}_t with none of \mathcal{F}_i containing a pair of sets with even-sized intersection, then we call the collection

$$\mathcal{F}_1 \times \dots \times \mathcal{F}_s := \{A_1 \times \dots \times A_s : A_i \in \mathcal{F}_i\} \subseteq \mathcal{O}_t^s$$

a *bad box*. As we explained above, in this case $|\mathcal{F}_i| \leq 2^{(t-1)/2}$, and therefore every bad box contains at most $2^{s(t-1)/2}$ sets. Note that the intersection of $A_1 \times \dots \times A_s$ and $B_1 \times \dots \times B_s$ is $(A_1 \cap B_1) \times \dots \times (A_s \cap B_s)$. Therefore, $\mathcal{F}_1 \times \dots \times \mathcal{F}_s$ is a bad box if and only if it contains no two sets with even intersection.

Take \mathcal{F} to be a random family given by choosing uniformly and independently, with repetition, $\lceil 2^{(t-1)s/4} \rceil$ sets in \mathcal{O}_t^s . Then, for every bad box \mathcal{B} , the probability that at least k elements of \mathcal{F} are contained in \mathcal{B} is at most $\binom{|\mathcal{F}|}{k} \left(\frac{|\mathcal{B}|}{|\mathcal{O}_t^s|}\right)^k \leq 2^{-(t-1)sk/4}$. The number of bad boxes can be upper bounded by $2^{s2^{t-1}}$. Thus, using our choices of k and t , we conclude that the probability that some bad box contains k elements of \mathcal{F} is at most

$$2^{s2^{t-1}} \cdot 2^{-(t-1)sk/4} \leq 1.$$

To finish, note that this implies that $\mathcal{F} \subseteq 2^{[n]}$ is $(\mathcal{C}_k, \{0\})$ -satisfying. Indeed, suppose this is not the case. Since each member of \mathcal{O}_t^s has odd size, there exist k sets $S_1, \dots, S_k \in \mathcal{F}$ such that for all distinct p, q , we have $|S_p \cap S_q| = 1 \pmod{2}$. Let the collection $\{A_j^{(i)} : 1 \leq i \leq k, 1 \leq j \leq s\}$ be such that $S_i = A_1^{(i)} \times \dots \times A_s^{(i)}$ for all i and for each j , let $\mathcal{F}_j = \{A_j^{(i)} : 1 \leq i \leq k\} \subseteq \mathcal{O}_t$. Since the size of $S_p \cap S_q$ is the product of the sizes of the intersections $A_j^{(p)} \cap A_j^{(q)}$, we must have that each collection \mathcal{F}_j has only odd-sized pairwise intersections. Hence, $\mathcal{F}_1 \times \dots \times \mathcal{F}_s$ is a bad box and contains the sets S_1, \dots, S_k . This is a contradiction, since no bad box contains k members of \mathcal{F} . \square

As $\Delta = k - 1$ and $t = \Theta(\log k)$, the family provided by the previous theorem has size $n^{\Omega(\log \Delta / \log \log \Delta)}$. It would be interesting to improve this result and get a better understanding of how much the exponent of n should depend on Δ .

3.3. Rainbow matchings

Let \mathcal{H} be an r -uniform multi-hypergraph (that is, we allow repetitions of the edges). Given a coloring $c : E(\mathcal{H}) \rightarrow [z]$, a rainbow matching in \mathcal{H} is a matching in which no two edges have the same color. The hypergraph \mathcal{H} is (z, t) -colored if it is colored with z colors, and each color class is a matching of size t . Let $f(r, t)$ denote the maximal z such that there exists a (z, t) -colored r -partite r -uniform multi-hypergraph which contains no rainbow matching of size t . Also, let $F(r, t)$ denote the maximal z such that there exists a (z, t) -colored r -uniform multi-hypergraph which contains no rainbow matching of size t . Clearly, $f(r, t) \leq F(r, t)$. Aharoni and Berger [1] proved that $f(r, t) \geq (t - 1)2^r$, and equality holds if $r = 2$ or $t = 2$. They also conjectured that $f(r, t) = (t - 1)2^r$

holds in general. This was disproved by Alon [2], who showed that $f(r, 3) \geq 2.71^r$. More precisely, Alon discovered a connection between $f(r, t)$ and the following well studied function. Let $g(r, t)$ denote the smallest integer g such that any sequence of g elements of the Abelian group \mathbb{Z}_t^r contains a subsequence of length t , whose elements sum up to zero. Then $f(r, t) \geq g(r - 1, t) - 1$.

On the other hand, Glebov, Sudakov and Szabó [9] proved, using combinatorial techniques, that

$$F(r, t) \leq \min\{(r + 1)^{2r+1}t^{2r+1}, 8^{rt}\}.$$

Our next theorem improves this upper bound for every (r, t) satisfying $r, t \geq 3$, which also improves all known upper bounds for $f(r, t)$ as well.

Theorem 9. $F(r, t) \leq (t - 1) \binom{rt}{r}$.

The proof is based on the exterior algebra method, which was famously used by Lovász [11] to prove the skew version of Bollobás’s set pair inequality. See also the result of Füredi [8] about a t -intersecting generalization of the problem. The interested reader can find a detailed description of this method as well as various applications in Chapter 6 of the book by Babai and Frankl [4]. Here, let us only give a basic introduction to exterior algebras.

Let V be a vector space over some field \mathbb{F} . The *exterior algebra* $\bigwedge V$ is the associative algebra generated by the elements of V and the associative binary operation \wedge , called *wedge product (or exterior product)*. Subject to these, \wedge has the additional property that $v \wedge v = 0$ for all $v \in V$. Also, the k -th *exterior power of k* , denoted by $\bigwedge^k V$, is the vector space generated by the elements $v_1 \wedge \dots \wedge v_k$, where $v_1, \dots, v_k \in V$. Let us list some of the well known properties of the wedge product, $\bigwedge^k V$ and $\bigwedge V$.

1. ($\bigwedge V$ is an associative algebra.) If $a, b, c \in \bigwedge V$ and $\lambda \in \mathbb{F}$, then

$$\begin{aligned} (a \wedge b) \wedge c &= a \wedge (b \wedge c), \\ a \wedge (\lambda b) &= (\lambda a) \wedge b = \lambda(a \wedge b), \\ a \wedge (b + c) &= (a \wedge b) + (a \wedge c), \\ (a + b) \wedge c &= (a \wedge c) + (b \wedge c). \end{aligned}$$

2. If $v, w \in V$, then $v \wedge w = -w \wedge v$.
3. If v_1, \dots, v_k , then $v_1 \wedge \dots \wedge v_k \neq 0$ if and only if v_1, \dots, v_k are linearly independent.
4. If $\dim(V) = n$, then $\dim(\bigwedge^k V) = \binom{n}{k}$. Moreover, if e_1, \dots, e_n is a basis of V , then

$$\{e_{i_1} \wedge \dots \wedge e_{i_k}\}_{1 \leq i_1 < \dots < i_k \leq n}$$

is a basis of $\bigwedge^k V$.

If \mathbb{F} has characteristic 2, then \wedge is also commutative by property 2. Therefore, in this case, if $A = \{a_1, \dots, a_k\} \subset V$, we can write $\bigwedge_{a \in A} a$ instead of $a_1 \wedge \dots \wedge a_k$ without specifying the order of terms.

Proof of Theorem 9. Let \mathcal{H} be an r -uniform multi-hypergraph with (z, t) -coloring c . Let \mathbb{F} be an infinite field of characteristic 2, and let V be an n -dimensional vector space over \mathbb{F} , where $n = rt$. For every $x \in V(\mathcal{H})$, choose a vector $v_x \in V$ such that these vectors are in general position, i.e., any n of them are linearly independent. For every r -tuple $A \subset V(\mathcal{H})$, let $w(A) = \bigwedge_{x \in A} v_x \in \bigwedge^r V$. Also, let e_1, \dots, e_n be a basis of V .

For $i \in [z]$, let $A_{i,1}, \dots, A_{i,t}$ be the edges of color i . Define the t -dimensional tensor $T : [z]^t \rightarrow \mathbb{F}$ as follows. With slight abuse of notation, let

$$T(i_1, \dots, i_t) = w(A_{i_1,1}) \wedge \dots \wedge w(A_{i_t,t}).$$

To be more precise, $w(A_{i_1,1}) \wedge \dots \wedge w(A_{i_t,t}) \in \bigwedge^n V$. But each $a \in \bigwedge^n V$ can be written as $a = \lambda(e_1 \wedge \dots \wedge e_n)$ for some $\lambda \in \mathbb{F}$, so we can identify a with λ .

Note that as $\{v_x\}_{x \in V(\mathcal{H})}$ are in general position, $T(i_1, \dots, i_t) \neq 0$ if and only if $A_{i_1,1}, \dots, A_{i_t,t}$ are pairwise disjoint. But then, as \mathcal{H} contains no rainbow matching of size t , T is semi-diagonal. By Theorem 1, this gives

$$\text{mfrank}(T) \geq \frac{z}{t-1}.$$

We finish the proof by showing that $\text{frank}_\ell(T) \leq \binom{n}{r}$ for every $\ell \in [t]$. For ease of notation, we show this for $\ell = 1$, the other cases follow by the same argument. For $I \subset [n]$, let $e_I = \bigwedge_{i \in I} e_i$. Then $\{e_I\}_{I \in [n]^{(r)}}$ is a basis of $\bigwedge^r V$, so for every $A \subset V(\mathcal{H})$ and $I \in [n]^{(r)}$ there exists $\lambda(A, I) \in \mathbb{F}$ such that

$$w(A) = \sum_{I \in [n]^{(r)}} \lambda(A, I) e_I.$$

But then

$$T(i_1, \dots, i_t) = \bigwedge_{j=1}^t \left(\sum_{I \in [n]^{(r)}} \lambda(A_{i_j,j}, I) e_I \right) = \sum_{\substack{I_1, \dots, I_t \in [n]^{(r)} \\ I_1 \cup \dots \cup I_t = [n]}} \lambda(A_{i_1,1}, I_1) \dots \lambda(A_{i_t,t}, I_t).$$

For $I \in [n]^{(r)}$, define the functions $f_I : [z] \rightarrow \mathbb{F}$ and $g_I : [z]^{t-1} \rightarrow \mathbb{F}$ as follows. Let

$$f_I(i_1) = \lambda(A_{i_1,1}, I),$$

and

$$g_I(i_2, \dots, i_t) = \sum_{\substack{I_2, \dots, I_t \in [n]^{(r)} \\ I_2 \cup \dots \cup I_t = [n] \setminus I}} \lambda(A_{i_2,2}, I_2) \dots \lambda(A_{i_t,t}, I_t).$$

Then

$$T(i_1, \dots, i_t) = \sum_{I \in [n]^{(r)}} f_I(i_1) g_I(i_2, \dots, i_t),$$

which shows that $\text{frank}_1(T) \leq \binom{n}{r}$. As this holds for $\text{frank}_\ell(T)$ as well for every $\ell \in [t]$, we get

$$\text{mfrank}(T) \leq \binom{n}{r}.$$

Comparing the lower and upper bound on the max-flattening rank of T , we deduce that $z \leq (t-1) \binom{n}{t}$, finishing the proof. \square

In particular, one can slightly modify our proof to show the following extension of the Bollobás set pair inequality [6], which might be of independent interest.

Theorem 10. *Let \mathcal{A} be a family of t -tuples of subsets of some base set X such that $|A(i)| = r_i$ for every $A \in \mathcal{A}$ and $i \in [t]$. Suppose that $A(1), \dots, A(t)$ are pairwise disjoint for every $A \in \mathcal{A}$, but $A_1(1), \dots, A_t(t)$ are not pairwise disjoint if $A_1, \dots, A_t \in \mathcal{A}$ are all distinct. Then*

$$|\mathcal{A}| \leq (t-1) \max_{i \in [t]} \binom{r_1 + \dots + r_t}{r_i}.$$

Declaration of competing interest

None declared.

Acknowledgements

We would like to thank the anonymous referee for many useful comments and suggestions. All authors were supported by the SNSF grant 200021_196965. István Tomon also acknowledges the support of The Russian Government in the framework of MegaGrant no 075-15-2019-1926, and the support of MIPT Moscow.

References

- [1] R. Aharoni, E. Berger, Rainbow matchings in r -partite r -graphs, *Electron. J. Comb.* 16 (1) (2009), #R119.
- [2] N. Alon, Multicolored matchings in hypergraphs, *Mosc. J. Comb. Number Theory* 1 (2011) 3–10.
- [3] N. Alon, M. Szegedy, Large sets of nearly orthogonal vectors, *Graphs Comb.* 15 (1999) 1–4.
- [4] L. Babai, P. Frankl, *Linear Algebra Methods in Combinatorics*, University of Chicago, 1988.
- [5] E.R. Berlekamp, On subsets with intersections of even cardinality, *Can. Math. Bull.* 12 (1969) 471–477.
- [6] B. Bollobás, On generalized graphs, *Acta Math. Acad. Sci. Hung.* 16 (1965) 447–452.

- [7] P. Frankl, R.M. Wilson, Intersection theorems with geometric consequences, *Combinatorica* 1 (1981) 357–368.
- [8] Z. Füredi, Geometrical solution of an intersection problem for two hypergraphs, *Eur. J. Comb.* 5 (1984) 133–136.
- [9] R. Glebov, B. Sudakov, T. Szabó, How many colors guarantee a rainbow matching?, *Electron. J. Comb.* 21 (1) (2014) P1.27.
- [10] V. Grolmusz, B. Sudakov, On k -wise set intersections and k -wise hamming distances, *J. Comb. Theory, Ser. A* 99 (2002) 180–190.
- [11] L. Lovász, Flats in matroids and geometric graphs, in: *Combinatorial Surveys, Proc. Sixth British Combinatorial Conf.*, Egham, 1977, 1977, pp. 45–86.
- [12] B. Sudakov, I. Tomon, Ramsey properties of algebraic graphs and hypergraphs, preprint, arXiv: 2103.05618, 2021.
- [13] V.H. Vu, Set systems with weakly restricted intersections, *Combinatorica* 19 (1999) 567–587.