

Embedded surfaces of arbitrary genus minimizing the Willmore energy under isoperimetric constraint

Laura Gioia Andrea Keller ^{*}, Andrea Mondino [†], Tristan Rivière [‡]

May 23, 2013

Abstract

The isoperimetric ratio of an embedded surface in \mathbb{R}^3 is defined as the ratio of the area of the surface to power three over the squared enclosed volume. The aim of the present work is to study the minimization of the Willmore energy under fixed isoperimetric ratio when the underlying abstract surface has fixed genus $g \geq 0$. The corresponding problem in the case of spherical surfaces, i.e. $g = 0$, was recently solved by Schygulla [57] with different methods.

1 Introduction and motivation

The Willmore functional of an immersion $\vec{\Phi}$ of an abstract oriented surface Σ^2 is given by

$$W(\vec{\Phi}) = \int_{\Sigma^2} |\vec{H}|^2 dvol_g$$

where \vec{H} is the mean curvature of the immersion $\vec{\Phi}$ and $dvol_g$ the induced volume form. A surface which is a critical point of this functional is called Willmore surface. Such a class of surfaces has been introduced in the beginning of the XX's century in the framework of "conformal geometry" by Wilhelm Blaschke who called them at the time "conformal minimal surfaces" (see [8]). He observed that the Willmore energy has the property to be invariant under conformal transformations of the ambient spaces, the transformations which infinitesimally preserve angles. He introduced then the theory of Willmore surfaces as being the "natural" merging between minimal surface theory and conformal invariance.

Probably because of the richness of the symmetries preserving the class of Willmore surfaces and because of the simplicity and the universality of its definition, the Willmore functional shows up in various fields of sciences and technology. It appears for instance in biology in the study of lipid bilayer cell membranes under the name "Helfrich energy" (as we will see in more detail later in the introduction), in general relativity as being the main term in the so called "Hawking

^{*}Institut für Numerische und Angewandte Mathematik, Universität Münster, 48149 Münster, Germany

[†]Department of Mathematics, ETH Zürich, 8092 Zürich, Switzerland

[‡]Department of Mathematics, ETH Zürich, 8092 Zürich, Switzerland

Mass”, in string theory in high energy physics it appears in the definition of the Polyakov extrinsic action, in elasticity theory as free energy of the non-linear plate Birkhoff theory, in optics and lens design, . . . etc.

The theory of Willmore surfaces, named after the person who revisited the theory in an important work of the mid-sixties [70]-[71], has flourished in the last decades.

Over the last twenty years, existence of minimizers of the Willmore energy under various constraints has been obtained: in the class of smooth immersions for a given abstract surface Σ^2 (see [60] and [3]), within a fixed conformal class (see [21] and [28]) or more recently in the class of smooth embeddings of the sphere under fixed isoperimetric ratio (see [57])

$$iso(\vec{\Phi}) = \frac{(Area(\vec{\Phi}))^3}{(Vol(\vec{\Phi}))^2} \quad (1)$$

where $Vol(\vec{\Phi})$ denotes the volume enclosed by $\vec{\Phi}(\Sigma^2)$, i.e. the volume of the bounded connected component of $\mathbb{R}^3 \setminus \vec{\Phi}(\Sigma^2)$.

The aim of the present work is to study the minimization of the Willmore energy under fixed isoperimetric ratio when the underlying abstract surface has fixed genus $g \geq 0$.

Beside the obvious geometric intrinsic interest such a minimization under isoperimetric and genus constraint could have, a motivation to study this problem comes from the modelization of the free energy of elastic lipid bilayer membranes in cell biology. Indeed the Willmore functional is closely related to the Helfrich functional which describes the free energy of a closed lipid bilayer

$$F_{Helfrich} = \int_{\text{lipid bilayer}} \left(\frac{k_c}{2}(2H + c_0)^2 + \bar{k}K + \lambda \right) + p \cdot V$$

where k_c and \bar{k} denote bending rigidities, c_0 stands for the spontaneous curvature, λ is the surface tension, K and H denote as usual the Gauss curvature and the mean curvature, respectively, p denotes the osmotic pressure and V denotes the enclosed volume.

The shapes of such membranes at equilibrium are then given by the corresponding Euler-Lagrange equation. If $c_0 = 0$, $\lambda = 0$ and $p = 0$ the Willmore functional captures the leading terms in Helfrich’s functional (up to a topological constant). Whereas if these physical constants do not vanish, λ and p can be seen as Lagrange multipliers for area and volume constraints. Thus, thanks to the invariance under rescaling of both the Willmore functional and the isoperimetric ratio, we exactly face the problem of minimizing the Willmore functional under an isoperimetric constraint.

In the context of vesicles, imposing a fixed area and a fixed volume has perfect biological meaning: on one hand, it is observed that at experimental time scales the lipid bilayers exchange only few molecules with the ambient and the possible contribution to the elastic energy due to displacements within the membrane is negligible. Thus, the area of the vesicle can be treated as a fixed one. On the

other hand, a change in volume would be the result of a transfer of liquid into or out of the vesicle. But this would significantly change the osmotic pressure and thus would lead to an energy change of much bigger scale than the scale of bending energy.

At first glimpse one may think that biologically relevant vesicles should always be of spherical shape. But in fact also vesicles of higher genus, and in particular of toroidal shape are observed (see [41] and [58]). Further details about this can be found e.g. in [34].

In addition, often in biology the ratio of area (the place where a molecule is produced) to volume (how much of the produced molecule can be stored) is crucial.

Moreover, the Helfrich functional as well as the Willmore functional are widely used for modeling biological phenomena, e.g. red blood cells (see e.g. [36]), folds of the endoplasmic reticulum (see e.g. [59]) and morphologies (Cristae junction) of mitochondria (see e.g. [43]).

Thus, also from an applied, biological point of view, it is perfectly reasonable to look at the problem we propose to study in this article.

In order to describe the results of this paper, let us introduce the framework where we are going to work.

In the aforementioned article of Schygulla [57], the main analytical strategy was to study immersions from the point of view of the image $\vec{\Phi}(\Sigma^2)$. This strategy, which has been extensively used also in a series of works by Kuwert and Schätzle [22]-[23]-[24]-[25]-[27]-[28]-[29], was introduced by Simon [60] and it is also known under the denomination of *Simon's ambient approach*.

In a series of papers (see [45], [48], [47]), the third author established a new framework for the study of variational problems related to the Willmore functional in which he was favoring the study of the immersion $\vec{\Phi}$ itself instead of its image. This approach could be called *the parametric approach*. It provides a general framework in which not only the above mentioned existence results (free minimization and minimization in a fixed conformal class) could be extended to the class of Lipschitz immersions with L^2 -bounded second fundamental form, also called *weak immersions*, but it is also suitable for applying fundamental principles of the calculus of variations (such as the mountain pass lemma for instance, in order to produce saddle type critical points).

In the present paper we will adopt this parametric approach. For the reader's convenience, all the important concepts as e.g. this space of Lipschitz immersions that we will denote \mathcal{E}_Σ , are introduced and explained in Section 2.

From now on, the underlying abstract surface is closed of genus $g \geq 0$ and our goal is to look for minimizers of the Willmore energy in the class of Lipschitz immersions with second fundamental form in L^2 under the additional constraint that the isoperimetric ratio, defined in (1), is fixed (by definition, if $\vec{\Phi} \in \mathcal{E}_{\Sigma^2}$ is not an embedding, i.e. it has self intersections, we set $iso(\vec{\Phi}) := 0$).

Our first result is an alternative proof, using the parametric approach, of the aforementioned theorem of Schygulla [57], i.e. existence of smooth embedded

spheres minimizing W under isoperimetric constraint; actually we manage to prove a stronger result, namely our minimization is performed on the larger class of weak immersions $\mathcal{E}_{\mathbb{S}^2}$ rather than among smooth embeddings. Before stating it, recall that by the isoperimetric inequality in \mathbb{R}^3 , for every embedded surface $\vec{\Phi}$ one has $iso(\vec{\Phi}) \leq iso(\mathbb{S}^2) = 36\pi$ and equality occurs if and only if $\vec{\Phi}$ is actually a round sphere.

Theorem 1.1 (The genus 0 case) *For every $R \in (0, 36\pi]$ there exists a smooth embedded spherical surface, called later on Schygulla sphere and denoted with $\mathbb{S}_{S,R}$, which minimizes the Willmore functional W among weak immersions $\mathcal{E}_{\mathbb{S}^2}$ having constrained isoperimetric ratio equal to R .*

Since, in virtue of the theorem above, the genus 0 case is well understood, from now on we will assume that Σ^2 is a surface of genus $g \geq 1$. Before stating our main theorems let us introduce some notation: let β_g^3 denote the infimum (actually it is a minimum thanks to [60]-[3]-[48]) of the Willmore energy among surfaces of genus $g \geq 1$ immersed in \mathbb{R}^3

$$\beta_g^3 := \inf \left\{ W(\vec{\Phi}) \mid \vec{\Phi} \text{ is an immersion of the genus } g \text{ closed surface} \right\} \quad , \quad (2)$$

let $\vec{\Phi}_g$ be a smooth minimizer attaining the infimum β_g^3 (notice that all the compositions of $\vec{\Phi}_g$ with conformal maps of \mathbb{R}^3 are still minimizers thanks to the conformal invariance of W), and let ω_g^3 be defined by

$$\omega_g^3 := \min \left\{ 4\pi + \sum_{i=1}^p (\beta_{g_i}^3 - 4\pi) \mid g = g_1 + \dots + g_p, 1 \leq g_i < g \right\} \quad . \quad (3)$$

Now we can state the main results of the paper; the first one guarantees the existence of minimizers of our problem under certain hypothesis and the second one discusses when such assumptions are satisfied.

Theorem 1.2 (Isoperimetric-constrained minimizers of W of every genus)

Let Σ_g^2 be the abstract genus $g \geq 1$ closed (i.e. compact without boundary) orientable surface and consider the set

$$I_g := \left\{ R \in \mathbb{R} \mid \inf_{\substack{\vec{\Phi} \in \mathcal{E}_{\Sigma_g^2} \\ iso(\vec{\Phi})=R}} W(\vec{\Phi}) < \min \{ 8\pi, \omega_g^3, \beta_g^3 + W(\mathbb{S}_{S,R}) - 4\pi \} \right\} \quad . \quad (4)$$

Then for given $R \in I_g$ there exists an embedding $\vec{\Phi}$ of Σ_g^2 into \mathbb{R}^3 , with $iso(\vec{\Phi}) = R$, which minimizes the Willmore energy among all Lipschitz immersions of Σ_g^2 with second fundamental form bounded in L^2 (i.e. among $\mathcal{E}_{\Sigma_g^2}$) and fixed isoperimetric ratio equal to R .

We even know that the minimizer from Theorem 1.2 is a smooth embedding.

Corollary 1.3 (Smoothness of the minimizer) *The minimizing $\vec{\Phi}$ from Theorem 1.2 is a smooth embedded genus $g \geq 1$ surface which minimizes the Willmore energy among smooth genus g embeddings with given isoperimetric ratio R .*

As announced above, we will characterize further the set I_g for which there exist minimizers of the Willmore energy under the additional constraint of given isoperimetric ratio.

Theorem 1.4 ($I_g \neq \emptyset$ is open) $I_g \subset \mathbb{R}$ is a non empty open set containing $iso(\vec{\Phi}_g)$, the isoperimetric ratio of any free minimizer of W among smooth genus g immersed closed surfaces.

Remark 1.5 ($I_g \supset [R_g - \delta, 36\pi)$) Notice that from Theorem 1.4, I_g contains the whole interval $[iso(\vec{\Phi}_g), 36\pi)$, where $36\pi = iso(\mathbb{S}^2)$. Indeed, since for every minimizer $\vec{\Phi}_g$ of the free minimization problem one has $iso(\vec{\Phi}_g) \in I_g$ and since the embeddings $\vec{\Xi} \circ \vec{\Phi}_g$ are still minimizers of W , where $\vec{\Xi}$ is any conformal transformation (i.e. a Moebius map) of \mathbb{R}^3 , it follows that $iso(\vec{\Xi} \circ \vec{\Phi}_g) \in I_g$ for every $\vec{\Xi}$.

Let us now consider a special smooth one parameter family $\{\vec{\Xi}_r\}_{r \in (0, +\infty)}$ of Moebius maps given by the inversion with respect to a sphere of unit radius and center $p(r)$, where $p : (0, \infty) \rightarrow \mathbb{R}^3$ is a smooth curve such that, for every $r > 0$, $p(r)$ is at distance $\frac{1}{r}$ from the fixed minimizer $\vec{\Phi}_g(\Sigma_g^2)$.

From this construction, it is not difficult to check that $iso(\vec{\Xi}_r \circ \vec{\Phi}_g)$ varies smoothly on r and that

$$\lim_{r \rightarrow 0^+} iso(\vec{\Xi}_r \circ \vec{\Phi}_g) = iso(\vec{\Phi}_g) \quad \& \quad \lim_{r \rightarrow +\infty} iso(\vec{\Xi}_r \circ \vec{\Phi}_g) = iso(\mathbb{S}^2) = 36\pi.$$

A continuity argument then shows that $I_g \supset [iso(\vec{\Phi}_g), 36\pi)$.

On the other hand, observe that for $R \in [iso(\vec{\Phi}_g), 36\pi)$, the isoperimetric-constrained minimizer produced by Theorem 1.2 is exactly the free minimizer $\vec{\Xi}_{r(R)} \circ \vec{\Phi}_g$, for a suitable $r = r(R)$. Nevertheless, called

$$R_g := \min_{\vec{\Xi} \in Moeb(\mathbb{R}^3)} iso(\vec{\Xi} \circ \vec{\Phi}_g)$$

(notice that the minimum is attained since, as explained above, when the the Moebius map diverges the isoperimetric ratio converges to the maximum value, given by $36\pi = iso(\mathbb{S}^2)$), the point of Theorem 1.4 is that I_g is *open*, so it contains an interval of the type $(R_g - \delta, 36\pi)$ for some $\delta > 0$; and in the interval $(R_g - \delta, R_g)$ the constrained minimizer is a new surface (i.e. not free Willmore) which is interesting to investigate. Let us also stress that we expect that the conditions used to define I_g are satisfied by a larger class of isoperimetric ratios, for more details see Remark 1.7. \square

Since the genus one case, i.e. $\Sigma^2 = \mathbb{T}^2$ is the 2-d torus, is particularly important for the applications, let us discuss it in more detail. Thanks to the recent proof of the Willmore conjecture by Marques and Neves [35], we know that $\beta_1^3 = 2\pi^2$, the set of minimizers is made by the Clifford torus $\mathbb{T}_{Clifford}^2$ (i.e. the torus of revolution with radii ratio $1 : \sqrt{2}$) and its images under conformal mappings of \mathbb{R}^3 ; moreover, by definition, $\omega_1^3 = +\infty$. We summarize Theorem 1.2, Corollary 1.3 and Theorem 1.4 for the genus one case in the following theorem.

Theorem 1.6 (Isoperimetric-constrained minimizers of W among tori)

Let \mathbb{T}^2 be the abstract 2-dimensional torus and consider the set

$$I_1 := \left\{ R \in \mathbb{R} \mid \inf_{\substack{\vec{\Phi} \in \mathcal{E}_{\mathbb{T}^2} \\ iso(\vec{\Phi})=R}} W(\vec{\Phi}) < \min \{ 8\pi, 2\pi^2 + W(\mathbb{S}_{S,R}) - 4\pi \} \right\}. \quad (5)$$

Then

i) $I_1 \subset \mathbb{R}$ is a non empty open set satisfying

$$I_1 \supset \left[16\pi\sqrt{2}, 36\pi \right),$$

where the numbers above come from the fact that $iso(\mathbb{S}^2) = 36\pi$ and $iso(\mathbb{T}_{Clifford}^2) = 16\pi\sqrt{2}$.

- ii) For given $R \in I_1$ there exists an embedding $\vec{\Phi}$ of \mathbb{T}^2 into \mathbb{R}^3 , with $iso(\vec{\Phi}) = R$, which minimizes the Willmore energy among all Lipschitz immersions of \mathbb{T}^2 with second fundamental form bounded in L^2 (i.e. among $\mathcal{E}_{\mathbb{T}^2}$) and fixed isoperimetric ratio equal to R .
- iii) The minimizing $\vec{\Phi}$ given in ii) is a smooth embedded torus which minimizes the Willmore energy among smooth embedded tori with isoperimetric ratio R .

Remark 1.7 (About the assumptions of Theorem 1.2) i) *The 8π bound:*

the natural framework for studying the isoperimetric constraint is given by embedded surfaces (i.e. surfaces without self intersection) since for this class it is clear what the enclosed volume is. A celebrated inequality of Li and Yau (see Theorem 2.4) states that if $\vec{\Phi}$ has self intersections then $W(\vec{\Phi}) \geq 8\pi$; therefore, since we want to handle embedded surfaces, it is natural to work assuming an 8π bound on the Willmore energy.

- ii) *The ω_g^3 bound:* this assumption is technical and it is used, together with the 8π bound, to ensure that the conformal structures of the minimizing sequence do not degenerate in the moduli space, see Theorem 5.2 in the Appendix. Notice that in the genus one case this condition is not needed. An interesting open problem is to rule out this condition from Theorem 1.2 and study also the possible degeneration.

- iii) *The $\beta_g^3 + W(\mathbb{S}_{S,R}) - 4\pi$ bound:* notice that the condition

$$\inf_{\substack{\vec{\Phi} \in \mathcal{E}_{\Sigma_g^2} \\ iso(\vec{\Phi})=R}} W(\vec{\Phi}) < \beta_g^3 + W(\mathbb{S}_{S,R}) - 4\pi \quad (6)$$

is that the *strict* inequality holds. In fact, for every given isoperimetric ratio different from the one of the round sphere the weak inequality

$$\inf_{\substack{\vec{\Phi} \in \mathcal{E}_{\Sigma_g^2} \\ iso(\vec{\Phi})=R}} W(\vec{\Phi}) \leq \beta_g^3 + W(\mathbb{S}_{S,R}) - 4\pi$$

holds. This can be seen as follows.

Let r and $\varepsilon > 0$ be given. Then by gluing a Schygulla sphere corresponding to an isoperimetric ratio $\tilde{r} = r + \delta$, for some small δ , and an inverted minimizer $\vec{\Phi}_g$, i.e. the image of $\vec{\Phi}_g$ (recall that $\vec{\Phi}_g$ is a minimizer of the Willmore energy among genus g immersed surfaces in \mathbb{R}^3 , i.e. $W(\vec{\Phi}_g) = \beta_g^3$) under a sphere inversion based at a point of $\vec{\Phi}_g(\Sigma_g^2)$, one obtains a surface whose isoperimetric ratio equals the given r . This can be achieved by appropriately choosing δ and the way how the two parts are glued together. Moreover, the Willmore energy of this new surface can be bounded from above by $\beta_g^3 + W(\mathbb{S}_{S,r}) - 4\pi + \varepsilon$. It is an interesting open problem whether or not the strict inequality (6) is actually always satisfied. \square

Now we briefly outline the strategy to prove Theorem 1.2. As mentioned above, following the approach of [48], we will directly work with the immersions $\vec{\Phi}_k$ and not with the immersed surfaces, i.e. the images $\vec{\Phi}_k(\Sigma^2) \subset \mathbb{R}^m$.

At first glimpse, it may appear a drawback to work with the immersions itself due to the invariance of the problem under the action of the noncompact group of diffeomorphisms of Σ^2 but, as we will see, this difficulty can be handled by choosing an appropriate gauge, namely the *Coulomb gauge*. Once such a Coulomb gauge is at hand, one can construct *conformal coordinates*. Together with suitable *estimates for the conformal factors* this leads to a setting in which we have very powerful analytical tools at hand in order to solve our problem. For example, we can use the fact that the equation for the Willmore surfaces can be reformulated as a conservation law, which enables us to convert the initial supercritical problem into a *critical* one.

More precisely, fixed $g \geq 1$, we consider a minimizing sequence $\{\vec{\Phi}_k\}$ in $\mathcal{E}_{\Sigma_g^2}$ for the Willmore energy under the constraint that the isoperimetric ratio $iso(\vec{\Phi}_k)$ is fixed and equals R , for some $R \in I_g$ where I_g is defined in (5). Notice that, since our abstract surface is a surface of genus $g \geq 1$, the reference metric of constant scalar curvature is flat or hyperbolic, moreover up to a change of coordinates we can assume that the $\vec{\Phi}_k$ s are conformal immersions (see Theorem 2.7) and, since by assumption $W(\vec{\Phi}_k) \leq \min\{8\pi, \omega_g^3\} - \delta$ (k big enough, for some $\delta > 0$), the conformal classes do not degenerate (see Theorem 5.2 in the Appendix).

The proof then splits into two cases: either the conformal factors of the immersions $\vec{\Phi}_k$ remain bounded or they diverge to $-\infty$.

In the first case, we proceed as follows: in a first step, we show that $\vec{\Phi}_k$ converges weakly in $W^{2,2}$ possibly away from finitely many points of energy concentration. Then, in a second step, we show a point removability result for these possible points of energy concentration. Once we have this, together with the exclusion of possible bubbling, we can conclude convergence in $\mathcal{E}_{\Sigma_g^2}$ and finally show smoothness of the minimizer.

In the second case, our analysis reveals that the assumption of diverging conformal factors yields a dichotomy between one part of the surface carrying the topological information and another part carrying the isoperimetric ratio; at this point we will perform a “cut-and-replace” argument (which recently appeared in a paper of the second and third authors [38]) which will bring to a

contradiction of the hypothesis (5) on I_g . Thus the case of diverging conformal factors is excluded and the proof is complete.

Acknowledgment Part of this work was done when the first two authors were visiting ETH Zürich. They would like to thank the FIM (Forschungsinstitut für Mathematik) and ETH Zürich for the hospitality and the excellent working atmosphere.

2 Setup and preliminaries

In this section we will recall the necessary definitions and we will collect and present the most important result which will play a crucial role in our analysis.

2.1 Definition of the Willmore functional, its fundamental properties and the Willmore surface equation

First of all, let us recall the definition of the Willmore functional, also called Willmore energy:

Let Σ_g^2 be an abstract oriented closed surface of genus $g \geq 0$ (sometimes we will simply write Σ^2 in place of Σ_g^2), and let $\vec{\Phi}$ be a C^2 -immersion of the surface Σ^2 into \mathbb{R}^m , $m \geq 3$ (we will mainly work immersions into \mathbb{R}^3). We denote with $g = \vec{\Phi}^* g_{\mathbb{R}^m}$ the pullback of the canonical metric on \mathbb{R}^m on $T\Sigma^2$ via the immersion $\vec{\Phi}$, g is also called first fundamental form. There is also a second fundamental form associated to the immersion $\vec{\Phi}$ and it is given by the following map

$$\begin{aligned} \vec{\mathbb{I}}_p : T_p \Sigma^2 \times T_p \Sigma^2 &\rightarrow (\vec{\Phi}_* T_p \Sigma^2)^\perp \\ (X, Y) &\mapsto \vec{\mathbb{I}}_p(X, Y) := \pi_{\vec{n}}(d^2 \vec{\Phi}(X, Y)) = \bar{\nabla}_{\vec{X}} \vec{X} - \nabla_Y X \end{aligned}$$

where $\vec{Z} = d\vec{\Phi} \cdot Z$, V^\perp is the orthogonal complement of the subspace V of \mathbb{R}^m , $\bar{\nabla}$ is the Levi-Civita connection in \mathbb{R}^m for $g_{\mathbb{R}^m}$ and ∇ is the Levi-Civita connection on $T\Sigma^2$ induced by g .

Recall that $\vec{\Phi}$ is said *conformal immersion* if $|\vec{\Phi}_x| \equiv |\vec{\Phi}_y|$ and $g_{\mathbb{R}^m}(\vec{\Phi}_x, \vec{\Phi}_y) \equiv 0$; in this case $e^\lambda = |\vec{\Phi}_x| = |\vec{\Phi}_y|$ is called the *conformal factor* of $\vec{\Phi}$.

Now, we can define the Willmore functional

Definition 2.1 (Willmore Functional) *Let Σ^2 , $\vec{\Phi}$ and g be as above. Then the Willmore functional of the immersion $\vec{\Phi}$ is given by the following expression*

$$W(\vec{\Phi}) = \int_{\Sigma^2} |\vec{H}|^2 \, dvol_g \quad ,$$

where $\vec{H} := \frac{1}{2} \text{tr}(g^{-1} \vec{\mathbb{I}}) = \frac{1}{2} \sum_{i,j=1}^2 g^{ij} \vec{\mathbb{I}}(\partial_{x_i}, \partial_{x_j})$ is the mean curvature, and $dvol_g$ is the volume form associated to g .

Note that in dimension three, we have $\vec{H} = H\vec{n}$, where \vec{n} is a unit vector orthogonal to $\vec{\Phi}_*(T\Sigma^2)$ in \mathbb{R}^3 .

Recall that the above Willmore functional is *conformally invariant* (see [11] and [50]), i.e. it is invariant under isometries, dilations and sphere inversions of the ambient space \mathbb{R}^n . Moreover, thanks to the Gauss-Bonnet theorem, we have

$$\begin{aligned} W(\vec{\Phi}) &= \frac{1}{4} \int_{\Sigma^2} |\Delta_g \vec{\Phi}|^2 dvol_g = \frac{1}{4} \int_{\Sigma^2} |d\vec{n}|_g^2 dvol_g + \pi\chi(\Sigma^2) \\ &= \frac{1}{4} \int_{\Sigma^2} |\vec{\mathbb{I}}|^2 dvol_g + \pi\chi(\Sigma^2) = \frac{1}{4} \int_{\Sigma^2} (\kappa_1 - \kappa_2)^2 dvol_g + 2\pi\chi(\Sigma^2) \quad , \end{aligned} \quad (7)$$

where $\chi(\Sigma^2) = 2g - 2$ denotes the Euler Characteristic of the surface Σ^2 , κ_i are the principal curvatures, and Δ_g denotes the intrinsic negative Laplace-Beltrami operator.

Next, recall the notion of Willmore immersion (or Willmore surface).

Definition 2.2 (Willmore immersion) *Let Σ^2 be a closed oriented surface and let $\vec{\Phi}$ be a smooth immersion of Σ^2 into \mathbb{R}^m . Then $\vec{\Phi}$ is called Willmore immersion if it is critical point of the Willmore functional W , i.e. if*

$$\forall \vec{\xi} \in C_0^\infty(\Sigma^2, \mathbb{R}^m) \quad \frac{d}{dt} W(\vec{\Phi} + t\vec{\xi})_{t=0} = 0.$$

The corresponding Euler-Lagrange equation is due to Weiner [68] and, for a smooth immersion into \mathbb{R}^3 , takes the form

$$\Delta_g H + 2H(H^2 - K) = 0 \quad (8)$$

where K is the Gauss curvature.

Unfortunately, there is a huge drawback in this equation: it is supercritical; i.e. while the functional can be defined for a weak immersion (see later for more details) with L^2 -bounded second fundamental form, the corresponding Euler-Lagrange equation written as in (8) needs that H is in L^3 to make sense, even in a distributional way. But thanks to the significant breakthrough in [45], the equation can be reformulated as a conservation law that makes sense for weak immersions with L^2 -bounded second fundamental form, i.e. the problem becomes critical. Namely it holds (see [45]; again we limit ourselves to the case of an immersion into \mathbb{R}^3)

Corollary 2.3 *A conformal immersion $\vec{\Phi}$ is Willmore if and only if*

$$\operatorname{div}(2\nabla\vec{H} - 3H\nabla\vec{n} - \nabla^\perp\vec{n} \times \vec{H}) = 0 \quad (9)$$

where $\nabla^\perp f = (-\partial_{x_2} f, \partial_{x_1} f)$ denotes the rotated gradient.

The fact that in the above corollary we assume the immersion to be conformal is not restrictive as we will see later on.

Before passing to the description of the variational framework in which we will work, let us recall the following lower bound on the Willmore energy due to Li and Yau [33].

Theorem 2.4 *Let Σ^2 be a closed surface and let $\vec{\Phi}$ be a smooth immersion of Σ^2 into \mathbb{R}^m . Assume that there exists a point $p \in \mathbb{R}^m$ with at least k pre-images under $\vec{\Phi}$. Then the following estimate holds*

$$W(\vec{\Phi}) \geq 4\pi k.$$

This estimate has a generalization to the context of varifolds which reads as follows (see e.g. Appendix of the article of Kuwert and Schätzle [24])

$$\inf_{\Sigma^2 \text{ smooth}} W(\Sigma^2) = 4\pi = \inf_{\mu \neq 0} W(\mu). \quad (10)$$

The above Theorem 2.4 has the following useful corollary.

Corollary 2.5 *Let Σ^2 be a closed surface and let $\vec{\Phi}$ be a smooth immersion of Σ^2 into \mathbb{R}^m . If*

$$W(\vec{\Phi}) < 8\pi$$

then $\vec{\Phi}$ is an embedding.

2.2 The variational framework

The variational framework in which we will study the minimization problem is the one of Lipschitz immersions with L^2 -bounded second fundamental form that we are now going to discuss.

We start by briefly recalling the definition of the Sobolev spaces $W^{k,p}(\Sigma^2, \mathbb{R}^m)$. Let Σ^2 be a smooth closed oriented 2-dimensional manifold and let g_0 be a smooth reference metric on it.

Then the Sobolev spaces $W^{k,p}(\Sigma^2, \mathbb{R}^m)$ are defined as

$$W^{k,p}(\Sigma^2, \mathbb{R}^m) := \left\{ f : \Sigma^2 \rightarrow \mathbb{R}^m \mid \sum_{j=1}^k \int_{\Sigma^2} |\nabla^j f|_{g_0}^p \, d\text{vol}_{g_0} < \infty \right\}.$$

Note that due to the fact that Σ^2 is assumed to be compact, the above spaces are independent of the metric g_0 .

One particular case is of great interest for us: the Lipschitz immersions $\vec{\Phi} \in W^{1,\infty}(\Sigma^2, \mathbb{R}^m)$ such that

$$\exists C > 0 \text{ such that } |d\vec{\Phi} \wedge d\vec{\Phi}|_{g_0} \geq C > 0. \quad (11)$$

Notice that by the compactness of Σ , $g := \vec{\Phi}^* g_{\mathbb{R}^m}$ defines a bounded metric comparable to g_0 ; i.e there exists $C > 0$ such that for all $X \in T\Sigma^2$ one has

$$C^{-1}g_0(X, X) \leq \vec{\Phi}^* g_{\mathbb{R}^m}(X, X) \leq Cg_0(X, X).$$

Next, observe that for a Lipschitz immersion satisfying (11), we have the following expression for the normal vector (called also Gauss map)

$$\vec{n}_{\vec{\Phi}} := * \frac{\partial_{x_i} \vec{\Phi} \wedge \partial_{x_j} \vec{\Phi}}{|\partial_{x_i} \vec{\Phi} \wedge \partial_{x_j} \vec{\Phi}|}$$

for an arbitrary choice of local positive coordinates (x_1, x_2) , where $*$ denotes the Hodge duality in \mathbb{R}^m . Notice that, from the definition, \vec{n} is an L^∞ -map on Σ^2 with values in \mathbb{R}^m .

The space in which we will study our minimization problem is given by the Lipschitz immersions of Σ^2 with Gauss map in $W^{1,2}$ (or, equivalently, with L^2 -bounded second fundamental form) defined as

$$\mathcal{E}_{\Sigma^2} := \left\{ \begin{array}{l} \vec{\Phi} \in W^{1,\infty}(\Sigma^2, \mathbb{R}^m) \text{ such that} \\ (11) \text{ holds for some } C > 0 \text{ and } \int_{\Sigma^2} |d\vec{n}|_g^2 dvol_g < \infty \end{array} \right\} .$$

In particular, we will consider immersions in \mathbb{R}^3 .

Observe that the Willmore functional is well defined on the space \mathcal{E}_{Σ^2} , see (7). Moreover, thanks to the (local) existence of a smooth conformal structure (see Theorem 2.7 below), we can extend the notion of *Willmore immersion* to the larger class \mathcal{E}_{Σ^2} as follows (in the case of immersions into \mathbb{R}^3).

Definition 2.6 (Weak Willmore immersion) *Let $\vec{\Phi}$ belong to \mathcal{E}_{Σ^2} .*

*$\vec{\Phi}$ is called a weak Willmore immersion if in any Lipschitz conformal chart Ψ from the two-dimensional disk D^2 into $(\Sigma^2, \vec{\Phi}^*g_{\mathbb{R}^3})$ the following holds*

$$\operatorname{div}(2\nabla\vec{H} - 3H\nabla\vec{n} - \nabla^\perp\vec{n} \times \vec{H}) = 0 \text{ in } \mathcal{D}'(D^2).$$

2.3 The isoperimetric constraint

As already mentioned in the introduction we want to study sequences of immersions of a closed surface Σ^2 of any genus minimizing the Willmore energy under the constraint that the isoperimetric ratio is fixed. More precisely, we minimize the Willmore energy among the weak immersions $\vec{\Phi} \in \mathcal{E}_{\Sigma^2}$ satisfying

$$\operatorname{iso}(\vec{\Phi}) := \frac{(\operatorname{Area}(\vec{\Phi}))^3}{(\operatorname{Vol}(\vec{\Phi}))^2} = R \quad ,$$

where R is a given constant and $\operatorname{Vol}(\vec{\Phi})$ denotes the volume enclosed by $\vec{\Phi}$, i.e. the volume of the bounded connected component of $\mathbb{R}^3 \setminus \vec{\Phi}(\Sigma^2)$.

Since by definition of I_g as in (5) we work below the energy threshold of 8π , thanks to Corollary 2.5 (or more precisely to its generalization to integer varifolds by Kuwert-Schätzle in [24, Appendix]), the weak immersions we will handle do not have any self intersections. Thus there are no difficulties in defining the isoperimetric ratio.

Observe that due to the scaling invariance of the Willmore functional, fixing the isoperimetric ratio is equivalent to fix both area and enclosed volume.

At this point, it may be of interest to ask which equation is satisfied by a critical point of the Willmore functional under the isoperimetric constraint. By scaling, we can assume that the enclosed volume equals to one, so fixing the isoperimetric ratio is equivalent to fix the area; therefore we are looking for critical points of the Willmore functional under area constraint. Applying the Lagrange multiplier principle, we find the following characterization - for conformal immersions which are area-constraint Willmore-

$$\Delta_g H \vec{n} + 2H(H^2 - K)\vec{n} = \mu H \vec{n} = \mu \vec{H} = \frac{\mu}{2} \Delta_g \vec{\Phi} = \frac{\mu}{2} e^{-2\lambda} \Delta \vec{\Phi} \quad ,$$

where we used $\vec{H} = \frac{1}{2}\Delta_g\vec{\Phi} = \frac{1}{2}e^{-2\lambda}\Delta\vec{\Phi}$. Equivalently, we have

$$\operatorname{div}(2\nabla\vec{H} - 3H\nabla\vec{n} - \nabla^\perp\vec{n} \times \vec{H} - \mu\nabla\vec{\Phi}) = 0 \quad (12)$$

where we used the description of Willmore surfaces given by equation (9). More details about this equation will be given later when studying the regularity of the minimizer.

2.4 Coulomb frames, conformal coordinates and estimates for the conformal factor

As mentioned above, in addition to the formulation of the Willmore surface equation as a conservation law, we have to break the symmetry group, i.e. the invariance under diffeomorphisms of the surface Σ^2 .

In fact, not only the right choice of coordinates, or gauge, is crucial, but in addition we need an appropriate control of the conformal factor.

A detailed presentation of the relation between conformal coordinates, Coulomb gauges and the Chern moving frame method is obviously beyond the scope of this article. Nevertheless, let us shortly summarize the most important results. First of all, the existence of conformal coordinates and a smooth conformal structure is asserted by the following theorem (see [47], [17])

Theorem 2.7 *Let Σ^2 be a closed smooth 2-dimensional manifold. Let $\vec{\Phi}$ be an element of \mathcal{E}_{Σ^2} . Then there exists a finite covering of Σ^2 by discs $(U_i)_{i \in I}$ and Lipschitz diffeomorphisms Ψ_i from the 2-dimensional unit disc D^2 into U_i such that $\vec{\Phi} \circ \Psi_i$ realizes a Lipschitz conformal immersion of D^2 . Since $\Psi_j^{-1} \circ \Psi_i$ is conformal and positive (i.e. holomorphic) on $\Psi_i^{-1}(U_i \cap U_j)$ the system of charts (U_i, Ψ_i) defines a smooth conformal structure c on Σ^2 and in particular there exists a constant scalar curvature metric g_c on Σ^2 and a Lipschitz diffeomorphism Ψ of Σ^2 such that $\vec{\Phi} \circ \Psi$ realizes a conformal immersion of the Riemann surface (Σ^2, g_c) .*

Once we know about the existence of a conformal immersion, we want to address the question whether the conformal factor can be estimated appropriately. Concerning this latter question, we have the following answer (see [50] and [48]).

Theorem 2.8 *Let $\vec{\Phi}$ be a Lipschitz conformal immersion from the disc D^2 into \mathbb{R}^m . Assume*

$$\int_{D^2} |\nabla\vec{n}_{\vec{\Phi}}|^2 < 8\pi/3. \quad (13)$$

Then for any $0 < \rho < 1$ there exists a constant C_ρ independent of $\vec{\Phi}$ such that for the conformal factor e^λ the following estimate holds

$$\sup_{p \in B_\rho^2(0)} e^\lambda(p) \leq C_\rho \left[\operatorname{Area}(\vec{\Phi}(D^2)) \right]^{1/2} \exp \left(C \int_{D^2} |\nabla\vec{n}_{\vec{\Phi}}|^2 \right).$$

Moreover, for two given distinct points p_1 and p_2 in the interior of D^2 and again for $0 < \rho < 1$ there exists a constant $C_\rho > 0$ independent of $\vec{\Phi}$ such that

$$\begin{aligned} \|\lambda\|_{L^\infty(B_\rho^2(0))} &\leq C_\rho \int_{D^2} |\nabla\vec{n}_{\vec{\Phi}}|^2 \\ &+ C_\rho \left| \log \frac{|\vec{\Phi}(p_1) - \vec{\Phi}(p_2)|}{|p_2 - p_1|} \right| + C_\rho \log^+ [C_\rho \operatorname{Area}(\vec{\Phi}(D^2))] \end{aligned}$$

where \log^+ denotes the positive part of the logarithm.

In our particular case, we will have a variant of such an estimate of the conformal factor.

For a precise statement of this modified estimate, we refer to Lemma 4.1 below.

Note that the (Willmore) energy assumption (13) is also crucial in Hélein's lifting theorem which asserts the existence of a moving frame with energy estimates provided that one starts with a map whose second fundamental form, measured in L^2 , is below the critical threshold.

Theorem 2.9 *Let \vec{n} be a $W^{1,2}$ -map from the 2-dimensional disc D^2 into the Grassmannian manifold $Gr_{m-2}(\mathbb{R}^m)$ of oriented $m-2$ -planes in \mathbb{R}^m . Then there exists a constant $C > 0$ such that there exist \vec{e}_1 and \vec{e}_2 in $W^{1,2}(D^2, S^{m-1})$ such that*

$$\vec{n} = *(\vec{e}_1 \wedge \vec{e}_2)$$

with

$$\int_{D^2} \sum_{i=1}^2 |\nabla \vec{e}_i|^2 < C \int_{D^2} |\nabla \vec{n}|^2$$

provided that

$$\int_{D^2} |\nabla \vec{n}|^2 < \frac{8\pi}{3}.$$

A proof of this theorem can be found in [17].

3 Proof of Theorem 1.1

First of all, let us state the following useful result proved by Schygulla [57].

Proposition 3.1 *For every $R \in (0, 36\pi]$ there exists a smooth embedded spherical surface with isoperimetric ratio equal to R and having Willmore energy strictly less than 8π .*

Let us recall the main lines of the nice proof: inverting a Catenoid in the origin and desingularizing it in 0, one obtains a spherical surface with energy strictly less than 8π and arbitrarily small isoperimetric ratio; then, make this surface evolve under Willmore flow. Thanks to the results of Kuwert-Schätzle [22]-[23]-[24], the flow will converge smoothly (as $t \uparrow +\infty$) to a round sphere (whose isoperimetric ratio is 36π); therefore, since the flow is smooth and does not increase the Willmore energy, and since the isoperimetric ratio depend continuously on the parameter of the flow, for every $R \in (0, 36\pi]$ one has produced a spherical surface with isoperimetric ratio equal to R and Willmore energy strictly less than 8π as desired.

The second ingredient for the proof of Theorem 1.1 is a compactness result for weak immersions of spheres proved by the second and third authors [38, Theorem I.2]. Since by Proposition 3.1 we can work under an $8\pi - \delta$ assumption, let us state the compactness result under this simplifying hypothesis (which prevents the bubbling phenomenon and the presence of branch points, thanks to Theorem 2.4).

Proposition 3.2 *Let $\{\vec{\Phi}_k\} \subset \mathcal{E}_{\mathbb{S}^2}$ be a sequence of weak conformal immersions of the 2-sphere \mathbb{S}^2 into \mathbb{R}^3 such that*

$$\sup_k \text{Area}(\vec{\Phi}_k) < \infty, \quad \inf_k \text{diam}(\vec{\Phi}_k(\mathbb{S}^2)) > 0 \quad \text{and} \quad W(\vec{\Phi}_k) < 8\pi - \delta \quad , \quad (14)$$

where $\text{diam}(E)$ is the diameter of the subset $E \subset \mathbb{R}^3$ and $\delta > 0$ is some positive constant.

Then there exist a subsequence that we still denote $\{\vec{\Phi}_k\}$, a sequence $\{f_k\}$ of elements in $\mathcal{M}^+(\mathbb{S}^2)$ (the positive Moebius group of \mathbb{S}^2) and finitely many points $\{a_1, \dots, a_n\}$ such that

$$\vec{\Phi}_k \circ f_k \rightharpoonup \vec{\Phi}_\infty \quad \text{weakly in } W_{loc}^{2,2}(\mathbb{S}^2 \setminus \{a_1 \cdots a_n\}) \quad , \quad (15)$$

where $\vec{\Phi}_\infty \in \mathcal{E}_{\mathbb{S}^2}$ is conformal. In addition, by lower semicontinuity of W under weak $W^{2,2}$ convergence, we have

$$W(\vec{\Phi}_\infty) \leq \liminf_k W(\vec{\Phi}_k) < 8\pi \quad ,$$

so $\vec{\Phi}_\infty$ is a (weak) embedding, and moreover

$$\text{Area}(\vec{\Phi}_k) \rightarrow \text{Area}(\vec{\Phi}_\infty) \quad \text{and} \quad \text{Vol}(\overline{\vec{\Phi}_k}) \rightarrow \text{Vol}(\overline{\vec{\Phi}_\infty}) \quad . \quad (16)$$

The proof Theorem 1.1 now follows quite easily by the two propositions above: thanks to Proposition 3.1, for any $R \in (0, 36\pi]$, the infimum of W among weak immersions in $\mathcal{E}_{\mathbb{S}^2}$ under the constraint of fixed isoperimetric ratio equal to R is strictly less than then 8π ; therefore, for any minimizing sequence $\{\vec{\Phi}_k\} \subset \mathcal{E}_{\mathbb{S}^2}$ of the constrained problem we have

$$W(\vec{\Phi}_k) \leq 8\pi - \delta \quad ,$$

for some $\delta > 0$. Therefore $\vec{\Phi}_k$ are (weak) embeddings and, as already observed by the scale invariance of W , we can assume that the enclosed volume of $\vec{\Phi}_k$ is constantly equal to 1 and the area of $\vec{\Phi}_k$ is constantly equal to $R^{1/3}$. From Simon's Lemma 5.5 recalled in the Appendix, we also have a strictly positive lower bound on the diameters of $\vec{\Phi}_k(\mathbb{S}^2)$ as subsets of \mathbb{R}^3 . Collecting the above informations, we conclude that $\vec{\Phi}_k$ satisfies the assumptions of Proposition 3.2; it follows that there exists a weak embedding $\vec{\Phi}_\infty \in \mathcal{E}_{\mathbb{S}^2}$, with $\text{iso}(\vec{\Phi}_\infty) = R$, which realizes the infimum of W under the isoperimetric constraint, as desired. For the proof of the regularity see Subsection 4.2.6. \square

In the next section we will analyze the more delicate case of genus $g \geq 1$.

4 Proof of Theorem 1.2

Let Σ^2 be a 2-dimensional surface of genus $g \geq 1$, and $\{\vec{\Phi}_k\} \subset \mathcal{E}_{\Sigma^2}$ be a minimizing sequence for the minimization problem of Theorem 1.2, i.e. minimization of the Willmore energy among weak immersions of Σ^2 under the constraint that the isoperimetric ratio $\frac{(\text{Area}(\vec{\Phi}_k))^3}{\left(\text{Vol}(\overline{\vec{\Phi}_k})\right)^2}$ is fixed and equals R , for some $R \in I_g$

where I_g is defined in (5). Our goal is to show that the minimizing sequence is compact, the limit belongs to \mathcal{E}_{Σ^2} and satisfies the required geometrical properties. The procedure we apply here is inspired by the one used in [48] (see also [6], Lemma II.1).

4.1 Normalization of the minimizing sequence

4.1.1 Conformality of $\vec{\Phi}_k \in \mathcal{E}_{\Sigma^2}$ in \mathcal{E}_{Σ^2}

Note that a priori, the elements of our sequence are not necessarily conformal. But thanks to theorem 2.7 we may replace the original $\vec{\Phi}_k$ s by $\vec{\Phi}_k \circ \Psi_k$ which are weakly conformal immersions.

In a few words, the idea behind the quoted theorem is that one starts with a frame with controlled energy, improves it into a Coulomb frame on each coordinate patch, applies the improvement of the well known Wente estimate due to Chanillo and Li (see [10]) and finally concludes by using the Riemann mapping theorem.

This new sequence will still be denoted - by abuse of the notation - $\{\vec{\Phi}_k\}$. Note that this new sequence consists of conformal immersions of Σ^2 . Rephrased, we have exploited the *invariance in the domain*.

Note that this procedure does not affect the isoperimetric ratio since of course this latter quantity is intrinsic, i.e. it does not depend on the choice of coordinates.

Remark At this stage, we would briefly comment on the difference between the present procedure and the one used in the unconstrained case presented in [48]. In this latter case the *invariance in the target under Möbius transformations* is used as well (see [48], Lemma A.4). In particular, inversions play a crucial role. Since such inversions do not preserve the isoperimetric ratio we will not exploit this invariance.

4.1.2 Points of energy concentration

Roughly speaking we will have only finitely many points $\{a_i\}$ where energy can accumulate with a critical energy threshold equal to $\frac{8\pi}{3}$. This is done as follows. First of all, recall that $g_k := \vec{\Phi}_k^* g_{\mathbb{R}^3}$ is the pull back metric, and h_k denotes the metric of constant scalar curvature associated to the conformal class c_k of the metric g_k .

Next, the hypothesis that the Willmore energy of our immersions stays below $\min\{8\pi, \omega_g^3\}$ (for sufficiently large k) implies that the conformal structures are contained in a compact subset of the moduli space of Σ^2 (see Theorem 5.2 in the Appendix). Thus the metrics h_k converge to h_∞ , the metric of constant scalar curvature associated to the limiting conformal structure c_∞ (about the existence of such a metric we refer to [19]). This convergence holds in $C^s(\Sigma^2)$ for all s .

Now, to each point x in Σ^2 we assign a critical radius which “cuts out $\frac{8\pi}{3}$ Willmore energy”, more precisely, this radius ρ_x^k is defined as follows

$$\rho_x^k := \inf \left\{ \rho \mid \int_{B_{\rho_k}(x)} |\nabla n_{\vec{\Phi}_k}|_{g_k}^2 dvol_{g_k} = \int_{B_{\rho_k}(x)} |\nabla n_{\vec{\Phi}_k}|_{h_k}^2 dvol_{h_k} = \frac{8\pi}{3} \right\}$$

where $B_{\rho_k}(x)$ denotes the geodesic ball in (Σ^2, g_k) centered at x with radius ρ_k . Obviously, $\{B_{\rho_k^x/2}(x)\}_{x \in \Sigma^2}$ is a covering of Σ^2 .

Next we extract a finite Besicovitch covering: each point $x \in \Sigma^2$ is covered by at most $N = N(\Sigma^2, g_\infty) \in \mathbb{N}$ such balls. This extracted covering is denoted $\{B_{\rho_k^i/2}(x_k^i)\}_{i \in I}$. Then we pass to a subsequence such that the following properties are satisfied

i)

I is independent on k

ii)

$$x_k^i \rightarrow x_\infty^i$$

iii)

$$\rho_k^i \rightarrow \rho_\infty^i.$$

And we set

$$I_0 := \{i \in I \text{ such that } \rho_\infty^i = 0\} \quad \text{and} \quad I_1 := I \setminus I_0.$$

Obviously,

$$\bigcup_{i \in I_1} \bar{B}_{\rho_\infty^i/2}(x_\infty^i) \text{ covers } \Sigma^2$$

(where the balls are measured in the metric g_∞) and thanks to the strict convexity of our balls - with respect to the flat or hyperbolic metric - there are only isolated, finitely many point in Σ^2 which are not covered by the union of the open balls. Thus we denote these exceptional point by $\{a_1, \dots, a_n\}$, or more precisely

$$\{a_1, \dots, a_n\} := \Sigma^2 \setminus \bigcup_{i \in I_1} B_{\rho_\infty^i/2}(x_\infty^i).$$

Rephrased, we have identified the points of energy concentration.

Note that the final covering of $\Sigma^2 \setminus \{a_1, \dots, a_n\}$,

$$\Sigma^2 \setminus \{a_1, \dots, a_n\} \subset \bigcup_{i \in I_1} B_{\rho_\infty^i/2}(x_\infty^i) \tag{17}$$

satisfies

$$\int_{B_{\rho_\infty^i}(x_\infty^i)} |\nabla \vec{n}_{\vec{\Phi}_k}|^2 \leq \frac{8\pi}{3}$$

for all k and all $i \in I_1$.

4.1.3 Analysis away from the energy-concentration-points: Control of the conformal factor

In this subsection we will perform the analysis of our sequence $\{\vec{\Phi}_k\}$ away from the energy-concentration points $\{a_1, \dots, a_n\}$. Let us start with an estimate for the conformal factors; as above, denote the conformal factor of $\vec{\Phi}_k$ by λ_k .

Lemma 4.1 *Let $\varepsilon > 0$. Then there exist constants C_k and a constant C_ε such that - up to passing to a subsequence - we have*

$$\|\lambda_k - C_k\|_{L^\infty(\Sigma^2 \setminus \cup_{i=1}^n B_\varepsilon(a_i))} \leq C_\varepsilon.$$

Note that C_ε does not depend on k , but only on ε . Whereas the C_k may depend on k .

A result of the same spirit is used in [48]. Although the proof is quite similar to the one in [48], let us recall the main arguments for the reader's convenience.

Proof of lemma 4.1:

First of all, since Σ^2 is a surface of genus at least one, the reference metric g_0 of constant scalar curvature is flat or hyperbolic. Therefore, recalling that the immersions $\vec{\Phi}_k$ are conformal so that $g_k := \vec{\Phi}_k^* g_{\mathbb{R}^3} = e^{2\lambda_k} g_0$, the conformal factors λ_k satisfy the Gauss-Liouville equation

$$-\Delta_{g_0} \lambda_k = K_{g_k} e^{2\lambda_k} + K_{g_0} \quad .$$

Now, since the second fundamental form is in L^2 , observe that $K_{g_k} e^{2\lambda_k}$ belongs to L^1 - with respect to g_0 . This fact implies by standard elliptic estimates that for a constant depending only on the surface Σ^2 and the metric g_0 we have

$$\begin{aligned} \|d\lambda_k\|_{L^{2,\infty}(\Sigma^2, g_0)} &\leq C \|\Delta_{g_0} \lambda_k\|_{L^1(\Sigma^2, g_0)} \leq C \left[\int_{\Sigma^2} |K_{g_k}| e^{2\lambda_k} d\text{vol}_{g_0} + |K_{g_0}| \text{Area}_{g_0}(\Sigma^2) \right] \\ &\leq C' \left[\int_{\Sigma^2} |K_{g_k}| d\text{vol}_{g_k} + 1 \right] \leq C' \left[\int_{\Sigma^2} |d\vec{n}_{\vec{\Phi}_k}|_{g_k}^2 d\text{vol}_{g_k} + 1 \right] \\ &\leq C'' \left[W(\vec{\Phi}_k) + 1 \right] < C''' . \end{aligned}$$

Now, let ε be given. Starting from the covering $\bigcup_{i \in I_1} B_{\rho_\infty^i/2}(x_\infty^i)$ (cf. (17)) we obtain a covering of $\Sigma^2 \setminus \cup_{i=1}^n B_\varepsilon(a_i)$ of the following form

$$\Sigma^2 \setminus \cup_{i=1}^n B_\varepsilon(a_i) \subset \bigcup_{i \in I_1} B_{r^i/2}(x_\infty^i)$$

where $r^i < \rho_\infty^i$ (the balls here and in the following steps are measured in the metric g_∞).

Note that the connectedness of Σ allows us, up to a relabeling of our balls, to assume that two consecutive balls have non-empty intersection. Recall now that by the properties of the covering (17) and the conformal invariance of the integrand, we have

$$\int_{B_{r^i}(x_\infty^i)} |\nabla \vec{n}_{\vec{\Phi}_k}|^2 < \frac{8\pi}{3} \quad .$$

Next we use Héleins moving frame method [17]:

Thanks to our construction, on the balls $B_{r^i}(x_\infty^i) \subset B_{\rho_\infty^i}(x_\infty^i)$ we have strictly less than the critical energy $\frac{8\pi}{3}$. Using Theorem 2.9 (for more details see [17] or [50]), upon identifying $B_{r^i}(x_\infty^i)$ with the 2-dimensional Euclidean unit disk D^2 , for each k there exists a moving frame with controlled energy, i.e. $(\vec{e}_k^1, \vec{e}_k^2) \in S^2 \times S^2$ with the following properties

i)

$$\vec{e}_k^1 \cdot \vec{e}_k^2 = 0 \text{ and } \vec{n}_{\vec{\Phi}_k} = \vec{e}_k^1 \wedge \vec{e}_k^2$$

ii)

$$\int_{D^2} (|\nabla \bar{e}_k^1|^2 + |\nabla \bar{e}_k^2|^2) \leq 2 \int_{D^2} |\nabla \bar{n}_{\bar{\Phi}_k}|^2 \leq \frac{16\pi}{3}$$

iii)

$$\begin{cases} \operatorname{div}(\bar{e}_k^1, \nabla \bar{e}_k^2) = 0 & \text{in } D^2 \\ \left(\bar{e}_k^1, \frac{\partial \bar{e}_k^2}{\partial \nu} \right) = 0 & \text{on } \partial D^2. \end{cases}$$

In these frames we can express the conformal factors λ_k as follows

$$-\Delta \lambda_k = (\nabla^\perp \bar{e}_k^1, \nabla \bar{e}_k^2).$$

Now, consider the solution μ_k of the following problem

$$\begin{cases} -\Delta \mu_k = (\nabla^\perp \bar{e}_k^1, \nabla \bar{e}_k^2) & \text{in } D^2 \\ \mu_k = 0 & \text{on } \partial D^2. \end{cases}$$

For this solution, Wente's theorem (see theorem 5.3 in the Appendix) gives the estimate

$$\begin{aligned} & \|\mu_k\|_{L^\infty(B_{r^i}(x_\infty^i))} + \|\nabla \mu_k\|_{L^{2,1}(B_{r^i}(x_\infty^i))} + \|\nabla^2 \mu_k\|_{L^1(B_{r^i}(x_\infty^i))} \\ & \leq C \int_{B_{r^i}(x_\infty^i)} (|\nabla \bar{e}_k^1|^2 + |\nabla \bar{e}_k^2|^2) \leq 2 \int_{B_{r^i}(x_\infty^i)} |\bar{n}_{\bar{\Phi}_k}|^2 \leq C \frac{16\pi}{3}. \end{aligned}$$

Next, we look at $v_k := \lambda_k - \mu_k$. Since v_k is harmonic, then we have (see e.g. [16])

$$\|v_k - \bar{v}_k\|_{L^\infty(B_{r^i/2}(x_\infty^i))} \leq C$$

where \bar{v}_k is the average of v_k over the ball $B_{r^i}(x_\infty^i)$. Putting together all the information we have so far, we conclude that there exist constants \bar{C}_k

$$\|\lambda_k - \bar{C}_k\|_{L^\infty(B_{r^i/2}(x_\infty^i))} \leq C.$$

In a last step, we combine the fact that the above arguments apply for all balls in our finite covering with the fact that two consecutive balls have non-empty intersection in order to conclude that the constant \bar{C}_k depends only on k but not on the ball we look at. This completes the proof of the lemma. \square

Now, a priori there are three possibilities: either C_k remain bounded, or they tend to $-\infty$ or diverge to $+\infty$.

Observe that The latter case is excluded, since our hypothesis imply that the area of $\bar{\Phi}_k$ remain bounded. Indeed, since the Willmore functional is scaling invariant, the isoperimetric constraint is equivalent to fix both area and enclosed volume. Thus, recalling that the area form is exactly $e^{2\lambda_k}$, the assumption that $C_k \rightarrow +\infty$ would imply that the area of $\bar{\Phi}_k(\Sigma^2)$, which is of course at least the area of $\Sigma^2 \setminus \cup_{i=1}^n B_\varepsilon(a_i)$, would become arbitrarily large. But this contradicts the assumption that the areas are fixed.

On the other hand, the fact that if C_k tend to $-\infty$ the area of $\Sigma^2 \setminus \cup_{i=1}^n B_\varepsilon(a_i)$ tends to zero does not a priori lead to a contradiction to the boundedness of the areas. In fact, due to a bubbling phenomenon, it could happen that we have large area on the exceptional ball $B_\varepsilon(a_i)$. So, a priori we can exclude only the possibility that $C_k \rightarrow +\infty$.

Below we will examine the remaining two possibilities.

4.2 The case of bounded conformal factors

In the present subsection, we assume that the C_k remain bounded. The case of diverging conformal factors will be discussed (and excluded) in Subsection 4.3.

4.2.1 Weak convergence in $W^{2,2}$

The assumption that the C_k remain bounded, i.e. $\sup_{k \in \mathbb{N}} |C_k| < \infty$, together with Lemma 4.1 immediately implies that

$$\limsup_{k \rightarrow \infty} \|\lambda_k\|_{L^\infty(\Sigma^2 \setminus \cup_{i=1}^n B_\varepsilon(a_i))} < \infty.$$

Therefore, recalling that $\Delta \vec{\Phi}_k = 2e^{2\lambda_k} \vec{H}_k$, we infer that the following estimates hold

$$\begin{cases} \|\Delta \vec{\Phi}_k\|_{L^2(\Sigma^2 \setminus \cup_{i=1}^n B_\varepsilon(a_i))} < C(\varepsilon) \\ \|\log |\nabla \vec{\Phi}_k|\|_{L^\infty(\Sigma^2 \setminus \cup_{i=1}^n B_\varepsilon(a_i))} < C(\varepsilon) \end{cases}$$

From that, we deduce that there exists a subsequence which converges weakly to a limit $\vec{\Phi}_\infty$ in $W^{2,2}(\Sigma^2 \setminus \cup_{i=1}^n B_\varepsilon(a_i))$; in particular, by Rellich Theorem, we have that

$$\nabla \vec{\Phi}_k \rightarrow \nabla \vec{\Phi}_\infty \text{ strongly in } L^p(\Sigma^2 \setminus \cup_{i=1}^n B_\varepsilon(a_i)) \quad \forall p < \infty$$

and hence upon passing to a subsequence, the gradients converge even almost everywhere pointwise.

Remark 4.2 Note that the above strong convergence of the gradients implies the convergence of the areas - away from the points a_i .

Moreover, by Sobolev embeddings we know that the sequence $\vec{\Phi}_k$ converges in $C^{0,\alpha}$ for $\alpha < 1$ and thus, roughly said, the enclosed volume converges as well away the points a_i and hence “the isoperimetric ratio is preserved in the limit away the points a_i ”. We will make this statement more precise later in Subsubsection 4.2.5. \square

4.2.2 Conformality of the limit

Recall that the sequence we have after all the preceding steps consists of conformal embeddings, i.e. we have for all k

$$\begin{cases} \partial_{x_1} \vec{\Phi}_k \cdot \partial_{x_2} \vec{\Phi}_k = 0 \\ |\partial_{x_1} \vec{\Phi}_k|^2 - |\partial_{x_2} \vec{\Phi}_k|^2 = 0. \end{cases}$$

Due to the a.e. pointwise convergence of the gradients $\nabla \vec{\Phi}_k$ (see above) we immediately can conclude that the conformality condition passes to the limit, i.e.

$$\begin{cases} \partial_{x_1} \vec{\Phi}_\infty \cdot \partial_{x_2} \vec{\Phi}_\infty = 0 \\ |\partial_{x_1} \vec{\Phi}_\infty|^2 - |\partial_{x_2} \vec{\Phi}_\infty|^2 = 0. \end{cases}$$

Together with the L^∞ -control of the conformal factors, this implies that

$$\vec{\Phi}_\infty \in \mathcal{E}_{\Sigma^2 \setminus \cup_{i=1}^n B_\varepsilon(a_i)}.$$

4.2.3 Control of $\vec{\Phi}_\infty$ over the whole Σ^2

Thanks to the results of the analysis performed so far, we can apply the following lemma due to Rivière (see [48], see also the article of Kuwert and Li [21]).

Lemma 4.3 *Let $\vec{\xi}$ be a conformal immersion of $D^2 \setminus \{0\}$ into \mathbb{R}^m in $W^{2,2}(D^2 \setminus \{0\}, \mathbb{R}^m)$ and such that $\log |\nabla \vec{\xi}| \in L_{loc}^\infty(D^2 \setminus \{0\})$. Assume $\vec{\xi}$ extends to a map in $W^{1,2}(D^2)$ and that the corresponding Gauss map $\vec{n}_{\vec{\xi}}$ also extends to a map in $W^{1,2}(D^2, Gr_{m-2}(\mathbb{R}^m))$. Then $\vec{\xi}$ realizes a Lipschitz conformal immersion of the whole disc D^2 and there exists a positive integer n and a constant C such that*

$$(C - o(1))|z|^{n-1} \leq \left| \frac{\partial \vec{\xi}}{\partial z} \right| \leq (C + o(1))|z|^{n-1}.$$

More precisely, we apply this lemma to $\vec{\Phi}_\infty$ around each exceptional point a_i , $i = 1, \dots, n$.

Claim: In our situation, the assertion of the above lemma holds with $n = 1$, i.e.

$$(C - o(1))|z| \leq \left| \frac{\partial \vec{\xi}}{\partial z} \right| \leq (C + o(1))|z|.$$

In other words, there is no branching.

Proof of the claim:

First of all, observe that the above chain of inequalities implies that for any $\delta > 0$ there exists a radius $r_\delta > 0$ such that for all $r < r_\delta$ we have

$$\vec{\Phi}_\infty(B_r(a_i)) \subset B_\rho(\vec{\Phi}_\infty(a_i)) \quad \text{and} \quad |\partial_z \vec{\Phi}_\infty| = e^{\lambda_\infty} \geq C \frac{1-\delta}{\sqrt{2}} |z|^{n-1}.$$

where $\rho = C2^{-1/2}n^{-1}(1+\delta)r^n$.

Using these facts, we then can estimate the mass of $\vec{\Phi}_\infty(\Sigma^2)$ inside the ball $B_\rho(\vec{\Phi}_\infty(a_i))$ as follows (note that here $\vec{\Phi}_\infty$ is seen as a varifold)

$$\begin{aligned} \mu(\vec{\Phi}_\infty \cap B_\rho(\vec{\Phi}_\infty(a_i))) &\geq C^2 \frac{(1-\delta)^2}{2} \int_{B_r(a_i)} |z|^{2n-2} \\ &\geq \frac{\pi C^2 (1-\delta)^2}{2n} r^{2n} \\ &\geq n\pi \left(\frac{1-\delta}{1+\delta} \right)^2 \rho^2. \end{aligned}$$

From this estimate we deduce that the 2-dimensional lower density of $\vec{\Phi}_\infty(\Sigma^2)$ at the point $\vec{\Phi}_\infty(a_i)$, $\theta_*^2(\vec{\Phi}_\infty(\Sigma^2), \vec{\Phi}_\infty(a_i))$ is bigger or equal to n .

On the other hand, the Li-Yau inequality (see [33]) - and in particular the extension of this inequality to the setting of varifolds with mean curvature in L^2 (see [24] (Appendix))-

$$\theta_*^2(\mu, x_*) \leq \frac{1}{4\pi} W(\mu)$$

implies that

$$n \leq \theta_*^2(\vec{\Phi}_\infty(\Sigma^2), \vec{\Phi}_\infty(a_i)) \leq \frac{W(\vec{\Phi}_\infty(\Sigma^2))}{4\pi} < \frac{8\pi - \delta}{4\pi} < 2$$

where we used also the lower semicontinuity of the Willmore functional W and the assumption that $W(\vec{\Phi}_k) < 8\pi - \delta$ for some $\delta > 0$.

Finally, this leads to the conclusion that $n = 1$. \square

In other words, we have a point removability phenomenon (cf. also proof of the above cited lemma).

Note that this together with the fact that away from the points a_i we have convergence in $C^{0,\alpha}$ implies that $\vec{\Phi}_k(\Sigma^2)$ remain bounded.

4.2.4 Limit of $\{\vec{\Phi}_k\}$ in \mathcal{E}_{Σ^2}

The results from the previous section immediately lead to the conclusion that $\vec{\Phi}_\infty$ is a Lipschitz immersion, and thus together with the $W^{2,2}$ convergence established earlier we find that $\vec{\Phi}_\infty \in \mathcal{E}_{\Sigma^2}$.

4.2.5 The isoperimetric constraint in the limit

It remains to show that the limit $\vec{\Phi}_\infty$ satisfies the isoperimetric constraint.

First of all, observe that in all the modifications of our initial minimizing sequence, we did nothing that could affect the isoperimetric constraint. So, we always have a sequence of immersions respecting the isoperimetric constraint. Recall that the requirement of fixed isoperimetric ratio can be rephrased as fixing the area as well as the enclosed volume.

At first glimpse, one might have the idea that the information we have at hand - more precisely that our sequence $\{\vec{\Phi}_k\}$ converges in $W^{2,2}$ - away from points of energy concentration - might be enough in order to show that the limit $\vec{\Phi}_\infty$ satisfies the required isoperimetric constraint. But this is not the case. We will explain this by looking at the area.

Convergence of the area would be a consequence of

$$e^{\lambda_k} \rightarrow e^{\lambda_\infty}.$$

But this convergence does not need to hold, since we have only the local control of the conformal factors

$$\|\lambda_k - C_k\|_{L^\infty(\Sigma^2 \setminus \cup_{i=1}^n B_\varepsilon(a_i))} \leq C_\varepsilon$$

from Lemma 4.1. Thus, the closer we get to the points a_i the bigger the L^∞ -norms of the conformal factors may become.

Our strategy in order to show that the limit $\vec{\Phi}_\infty$ satisfies the isoperimetric constraint is to exclude a bubbling phenomenon.

Roughly speaking, we will detect regions of positive area which carry an energy contribution of at least 4π . This will lead to a contradiction to our initial hypothesis that $W(\vec{\Phi}_k) < 8\pi - \delta$. More precisely, we prove the following lemma.

Lemma 4.4 *Let $\{\vec{\Phi}_k\}$ be as above and denote by $a_i, i = 1, \dots, n$, the points of energy concentration. Then we have*

i) Assume that there exists an index i such that

$$\liminf_{\varepsilon \rightarrow 0} \liminf_{k \rightarrow \infty} d(\vec{\Phi}_k(B_\varepsilon(a_i)), \vec{\Phi}_k(\partial B_\varepsilon(a_i))) > 0 \quad ,$$

where d denotes the usual distance between two sets $d(A, B) := \sup_{p \in B} \inf_{q \in A} |p - q|$, then

$$\liminf_{\varepsilon \rightarrow 0} \liminf_{k \rightarrow \infty} W(\vec{\Phi}_k(B_\varepsilon(a_i))) \geq 4\pi.$$

ii) Assume that there exists an i such that

$$\liminf_{\varepsilon \rightarrow 0} \liminf_{k \rightarrow \infty} \text{Area}(\vec{\Phi}_k(B_\varepsilon(a_i))) > 0$$

then

$$\liminf_{\varepsilon \rightarrow 0} \liminf_{k \rightarrow \infty} W(\vec{\Phi}_k(B_\varepsilon(a_i))) \geq 4\pi.$$

Proof of Lemma 4.4:

Before passing to the proof of *i)* let us start with a general observation (which holds independently of *i)* or *ii)* above): we claim that, up to passing to a subsequence in k , we have

$$\lim_{\varepsilon \rightarrow 0} \lim_{k \rightarrow \infty} \mathcal{H}^1(\vec{\Phi}_k(\partial B_\varepsilon(a_i))) \rightarrow 0. \quad (18)$$

Indeed, recall that we have

$$\vec{\Phi}_k \rightarrow \vec{\Phi}_\infty \text{ weakly in } W^{2,2}(\Sigma^2 \setminus \cup_i B_\varepsilon(a_i)) \quad .$$

Then from the classical trace theorem (see for instance [54]), we know that $\nabla \vec{\Phi}_k(\partial B_\varepsilon(a_i))$ converges weakly in $F_{2,2}^{\frac{1}{2}}(\partial B_\varepsilon(a_i)) = H^{\frac{1}{2}}(\partial B_\varepsilon(a_i))$. Here, $F_{p,q}^s$ denotes the standard Triebel-Lizorkin space. Therefore, from the standard compactness part of the Sobolev embedding theorem, we find that $\nabla \vec{\Phi}_k(\partial B_\varepsilon(a_i))$ converges even strongly in $L^2(\partial B_\varepsilon(a_i))$ and in particular in $L^1(\partial B_\varepsilon(a_i))$.

Thus we have

$$\mathcal{H}^1(\vec{\Phi}_k(\partial B_\varepsilon(a_i))) = \int_{\partial B_\varepsilon(a_i)} |\dot{\vec{\Phi}}_k| dl \rightarrow \mathcal{H}^1(\vec{\Phi}_\infty(\partial B_\varepsilon(a_i))) = \int_{\partial B_\varepsilon(a_i)} |\dot{\vec{\Phi}}_\infty| dl.$$

Finally, we recall that the limit immersion $\vec{\Phi}_\infty$ is Lipschitz, so

$$\lim_{\varepsilon \rightarrow 0} \int_{\partial B_\varepsilon(a_i)} |\dot{\vec{\Phi}}_\infty| dl \rightarrow 0 \quad ,$$

since $|\partial B_\varepsilon(a_i)| \rightarrow 0$ as $\varepsilon \rightarrow 0$.

Proof of part i):

In order to prove statement *i)*, we will use a result of Rivière (see [47]) giving an estimate on the Willmore energy of a compact surface with boundary; for

the precise statement see Lemma 5.4 in the Appendix. Applying the mentioned estimate to the immersions $\vec{\Phi}_k$ restricted to $B_\varepsilon(a_i)$, we find

$$\begin{aligned} 4\pi &\leq \liminf_{\varepsilon \rightarrow 0} \liminf_{k \rightarrow \infty} \left(W(\vec{\Phi}_k(B_\varepsilon(a_i))) + 2 \frac{\mathcal{H}^1(\vec{\Phi}_k(\partial B_\varepsilon(a_i)))}{d(\vec{\Phi}_k(\partial B_\varepsilon(a_i)), \vec{\Phi}_k(B_\varepsilon(a_i)))} \right) \\ &= \liminf_{\varepsilon \rightarrow 0} \liminf_{k \rightarrow \infty} W(\vec{\Phi}_k(B_\varepsilon(a_i))) \quad , \end{aligned} \quad (19)$$

where in the last equality we used (18) together the assumption of i). Thus, assertion i) is proved.

Proof of part ii): Without loss of generality we might assume that the hypothesis holds for the point a_1 . In the sequel it is enough to study the situation around this particular point.

First of all let us recall the following assertion which can be seen as a version of Green's theorem for immersed surfaces:

Let Σ^2 be a smooth compact surface with boundary, let $\vec{\Phi} \in \mathcal{E}_{\Sigma^2}$ be an immersion of Σ^2 into \mathbb{R}^n and let \vec{X} be a smooth vector field in \mathbb{R}^m . Then it holds

$$\int_{\vec{\Phi}(\Sigma^2)} \operatorname{div}_{\vec{\Phi}(\Sigma^2)} \vec{X} \, d\operatorname{vol}_{g_{\vec{\Phi}}} = \int_{\vec{\Phi}(\partial\Sigma^2)} \langle \vec{X}, \vec{\nu} \rangle \, dl_{\vec{\Phi}(\partial\Sigma^2)} - 2 \int_{\vec{\Phi}(\Sigma^2)} \langle \vec{X}, \vec{H} \rangle \, d\operatorname{vol}_{g_{\vec{\Phi}}} \quad (20)$$

where

$$\operatorname{div}_{\vec{\Phi}(\Sigma^2)} \vec{X} := \sum_{k=1}^2 \langle d\vec{X} \cdot \vec{e}_k, \vec{e}_k \rangle$$

for any local orthonormal frame (\vec{e}_1, \vec{e}_2) on $\vec{\Phi}(\Sigma^2)$ and $\vec{\nu} := e^{-\lambda} \partial_r \vec{\Phi}$ (where as usual λ denotes the conformal factor of the immersion $\vec{\Phi}$).

A proof of this classical formula can be found for instance in [47].

Now, we continue the proof of part ii) of the Lemma 4.4.

In a first step, we will apply formula (20) to $\vec{\Phi}_k$, restricted to the ball $B_\varepsilon(a_1)$, and to the vector field $\vec{X}(x) = x - \vec{\Phi}_k(a_1)$; observing that $\operatorname{div} X \equiv 2$, we get

$$\begin{aligned} 2\operatorname{Area}(\vec{\Phi}_k(B_\varepsilon(a_1))) &= \int_{\vec{\Phi}(B_\varepsilon(a_1))} \operatorname{div} \vec{X} \, d\operatorname{vol}_{g_k} \\ &= \int_{\vec{\Phi}(\partial B_\varepsilon(a_1))} \langle \vec{X}, \vec{\nu} \rangle \, dl - 2 \int_{\vec{\Phi}(B_\varepsilon(a_1))} \langle \vec{X}, \vec{H} \rangle \, d\operatorname{vol}_{g_k} \\ &\leq \operatorname{diam}(\vec{\Phi}_k(B_\varepsilon(a_1))) \mathcal{H}^1(\partial\vec{\Phi}_k(B_\varepsilon(a_1))) - 2 \int_{\vec{\Phi}(B_\varepsilon(a_1))} \langle \vec{X}, \vec{H} \rangle \, d\operatorname{vol}_{g_k} \\ &\leq \operatorname{diam}(\vec{\Phi}_k(B_\varepsilon(a_1))) \mathcal{H}^1(\partial\vec{\Phi}_k(B_\varepsilon(a_1))) \\ &\quad + 2\operatorname{diam}(\vec{\Phi}_k(B_\varepsilon(a_1))) W(\vec{\Phi}_k(B_\varepsilon(a_1)))^{\frac{1}{2}} \operatorname{Area}(\vec{\Phi}_k(B_\varepsilon(a_1)))^{\frac{1}{2}}. \end{aligned}$$

Rearranging terms, we find

$$\begin{aligned} \operatorname{Area}(\vec{\Phi}_k(B_\varepsilon(a_1)))^{\frac{1}{2}} &- \frac{\operatorname{diam}(\vec{\Phi}_k(B_\varepsilon(a_1))) \mathcal{H}^1(\partial\vec{\Phi}_k(B_\varepsilon(a_1)))}{2\operatorname{Area}(\vec{\Phi}_k(B_\varepsilon(a_1)))^{\frac{1}{2}}} \\ &\leq \operatorname{diam}(\vec{\Phi}_k(B_\varepsilon(a_1))) W(\vec{\Phi}_k(B_\varepsilon(a_1)))^{\frac{1}{2}}. \end{aligned}$$

Next, we recall that the diameters of the whole surfaces $\vec{\Phi}_k(\Sigma^2)$ as well as the diameter of the limit $\vec{\Phi}_\infty(\Sigma^2)$ are uniformly bounded thanks to Lemma 5.5; thus we estimate

$$\begin{aligned}
8\pi \operatorname{diam}(\vec{\Phi}_k(B_\varepsilon(a_1))) &> W(\vec{\Phi}_k) \operatorname{diam}(\vec{\Phi}_k(B_\varepsilon(a_1))) \\
&\geq W(\vec{\Phi}_k(B_\varepsilon(a_1))) \operatorname{diam}(\vec{\Phi}_k(B_\varepsilon(a_1))) \\
&\geq \operatorname{Area}(\vec{\Phi}_k(B_\varepsilon(a_1)))^{\frac{1}{2}} \\
&\quad - \frac{\operatorname{diam}(\vec{\Phi}_k(B_\varepsilon(a_1))) \mathcal{H}^1(\partial\vec{\Phi}_k(B_\varepsilon(a_1)))}{2\operatorname{Area}(\vec{\Phi}_k(B_\varepsilon(a_1)))^{\frac{1}{2}}} \\
&\geq \delta^{\frac{1}{2}} - \frac{\operatorname{diam}(\vec{\Phi}_k(B_\varepsilon(a_1))) \mathcal{H}^1(\partial\vec{\Phi}_k(B_\varepsilon(a_1)))}{2\operatorname{Area}(\vec{\Phi}_k(B_\varepsilon(a_1)))^{\frac{1}{2}}} \\
&\geq \delta^{\frac{1}{2}} - \frac{C \mathcal{H}^1(\partial\vec{\Phi}_k(B_\varepsilon(a_1)))}{2\operatorname{Area}(\vec{\Phi}_k(B_\varepsilon(a_1)))^{\frac{1}{2}}}, \quad \text{for some } C > 0 \\
&> \delta_1 > 0, \quad \text{since } \mathcal{H}^1(\partial\vec{\Phi}_k(B_\varepsilon(a_1))) \rightarrow 0
\end{aligned}$$

Therefore we found that $\operatorname{diam}(\vec{\Phi}_k(B_\varepsilon(a_1))) \geq \delta_2 > 0$ for some δ_2 . But this together with the fact that $\mathcal{H}^1(\partial\vec{\Phi}_k(B_\varepsilon(a_1))) \rightarrow 0$ implies that

$$d(\vec{\Phi}_k(\partial B_\varepsilon(a_1)), \vec{\Phi}_k(B_\varepsilon(a_1))) > \delta^*$$

for some positive δ^* . We can now conclude the proof of part *ii*) by applying the (already proved) part *i*). \square

Now, let us explain how Lemma 4.4 implies that the limit satisfies the isoperimetric constraint. We claim that, for every concentration point a_i ,

$$\liminf_{\varepsilon \rightarrow 0} \liminf_{k \rightarrow \infty} d(\vec{\Phi}_k(B_\varepsilon(a_i)), \vec{\Phi}_k(\partial B_\varepsilon(a_i))) = 0 \quad (21)$$

and

$$\liminf_{\varepsilon \rightarrow 0} \liminf_{k \rightarrow \infty} \operatorname{Area}(\vec{\Phi}_k(B_\varepsilon(a_i))) = 0. \quad (22)$$

Indeed, if by contradiction there exists a concentration point, say a_1 , where one of the two statements above fail then

$$\begin{aligned}
\liminf_{k \rightarrow \infty} W(\vec{\Phi}_k) &= \lim_{\varepsilon \rightarrow 0} \liminf_{k \rightarrow \infty} \left[W(\vec{\Phi}_k(\Sigma^2 \setminus \cup_i B_\varepsilon(a_i))) + W(\vec{\Phi}_k(\cup_i B_\varepsilon(a_i))) \right] \\
&\geq W(\vec{\Phi}_\infty) + 4\pi \geq 8\pi, \quad (23)
\end{aligned}$$

where the first inequality comes from the lower semicontinuity of W under weak- $W^{2,2}$ convergence together with the fact that $\vec{\Phi}_\infty$ is an element of \mathcal{E}_{Σ^2} and Lemma 10 applied to a_1 ; the last inequality follows from $W(\vec{\Phi}_\infty) \geq 4\pi$.

But this last estimate is in contradiction with our hypothesis that the Willmore energy stays strictly below 8π . Thus, our claim (21)-(22) holds.

Now, combining Remark 4.2 and the second claim (22) we deduce that the areas converge, i.e.

$$\operatorname{Area}(\vec{\Phi}_k) \rightarrow \operatorname{Area}(\vec{\Phi}_\infty).$$

Moreover, combining (18) and the first claim (21), we infer that also the diameters have to vanish:

$$\text{diam}(\vec{\Phi}_k(B_\varepsilon(a_i))) \rightarrow 0 \text{ for all } i.$$

But this last fact together with Remark 4.2 immediately leads to the conclusion that the enclosed volumes converge as well

$$\text{Vol}(\vec{\Phi}_k) \rightarrow \text{Vol}(\vec{\Phi}_\infty).$$

We conclude that the isoperimetric constraint passes to the limit.

4.2.6 Regularity of the limit, proof of Corollary 1.3

First of all, thanks to Corollary 2.5 observe that the uniform energy estimate

$$W(\vec{\Phi}_k) < 8\pi - \delta$$

together with the lower semicontinuity of the Willmore functional under weak $W^{2,2}$ convergence immediately implies that the limit $\vec{\Phi}_\infty$ is an embedding, i.e. $\vec{\Phi}_\infty$ has no self intersection. Moreover, as we have seen in the preceding subsection, the limit satisfies the isoperimetric constraint.

Now, we study the regularity question. In a first step we will recall that the Willmore functional W is Fréchet differentiable and we will determine dW . Then we will establish the equation which is satisfied by the limit $\vec{\Phi}_\infty$ and finally show the regularity.

Lemma 4.5 *Let $\vec{\Phi}$ belong to \mathcal{E}_{Σ^2} . Then the Willmore functional W is Fréchet differentiable with respect to variations $\vec{w} \in W^{1,\infty} \cap W^{2,2}$ with compact support. Moreover, for the differential we have the following formula*

$$d_{\vec{\Phi}}W[\vec{w}] = \int \nabla \vec{w} \cdot \left(2\nabla \vec{H} - 3H\nabla \vec{n} - \nabla^\perp \vec{n} \times \vec{H} \right).$$

This lemma is a straightforward adaptation of the corresponding observation made in [48] (see also the Appendix of [39]). Note that the differential exactly corresponds to the reformulation of the Willmore equation in the form of a conservation law as in (9).

Next, we give the equation which is satisfied by our limiting object $\vec{\Phi}_\infty$.

Lemma 4.6 *Let $\vec{\Phi}_\infty$ be as above. Then it satisfies the following equation*

$$\text{div}(2\nabla \vec{H} - 3H\nabla \vec{n} - \nabla^\perp \vec{n} \times \vec{H} - \mu \nabla \vec{\Phi}) = 0.$$

Proof: The assertion is an immediate consequence of the lower semicontinuity of the Willmore functional W and its differentiability, the classical fact that the area functional A is Fréchet differentiable as well with differential

$$d_{\vec{\Phi}}A[\vec{w}] = - \int 2\vec{H} \cdot \vec{w}$$

and the principle of Lagrange multipliers. To get the final formula recall also that $\vec{H} = \frac{1}{2}\Delta_g \vec{\Phi} = \frac{1}{2}e^{-2\lambda}\Delta \vec{\Phi}$. \square

Finally we have the following regularity assertion, whose proof is an easy adaptation of the regularity of Willmore immersions established in [45] (for the regularity of area-constrained Willmore immersions see also [39], and for a comprehensive explanation see [50]).

Lemma 4.7 $\vec{\Phi}_\infty$ is smooth.

The proof of Corollary 1.3 is now complete. \square

4.3 The case of diverging conformal factors

So far, we have studied the case when C_k remain bounded. Now, we will analyze the case where this is no longer true. As we have seen above, the only remaining possibility is that - up to extraction of a subsequence - the conformal factors tend to $-\infty$.

4.3.1 Existence of at most one bubble

As above, the point where Willmore energy concentrates are denoted by a_i , and again we may assume that we have

$$\text{Area}(\vec{\Phi}_k) = 1 \quad \forall k.$$

Thus, the fact that now the conformal factors diverge, i.e.

$$\lambda_k \rightarrow -\infty \text{ on } \Sigma^2 \setminus \cup_i B_\varepsilon(a_i)$$

implies that there is at least one point a_i where area concentrates. More precisely, we have that there exists a point a^* such that

$$\liminf_{\varepsilon \rightarrow 0} \liminf_{k \rightarrow \infty} \text{Area}(\vec{\Phi}_k(B_\varepsilon(a_i))) > 0. \quad (24)$$

Our next goal is to show that there is exactly one such point; this is the content of the following lemma, whose proof follows by the $8\pi - \delta$ bound on the Willmore energy of $\vec{\Phi}_k$ and the second part of Lemma 4.4.

Lemma 4.8 Assume that the conformal factors tend to $-\infty$, i.e.

$$\lambda_k \rightarrow -\infty \text{ as } k \rightarrow \infty.$$

Then there exists exactly one point $a^* \in \Sigma^2$ of concentration of the area, i.e. where (24) holds. Of course a^* is also a point of concentration of the energy, i.e. $a^* \in \{a_i\}$.

Directly from the lemma it follows that

$$\liminf_{\varepsilon \rightarrow 0} \liminf_{k \rightarrow \infty} \text{Area}(\vec{\Phi}_k(B_\varepsilon(a^*))) = 1 \quad \text{and} \quad (25)$$

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{k \rightarrow \infty} \text{Area}(\vec{\Phi}_k(\Sigma^2 \setminus B_\varepsilon(a^*))) = 0 \quad . \quad (26)$$

Analogously, again using the $8\pi - \delta$ bound on the Willmore energy of $\vec{\Phi}_k$ and the first part of Lemma 4.4, we also have

$$\liminf_{\varepsilon \rightarrow 0} \liminf_{k \rightarrow \infty} d(\vec{\Phi}_k(B_\varepsilon(a^*)), \vec{\Phi}_k(\partial B_\varepsilon(a_i))) > 0 \quad \text{and} \quad (27)$$

$$\liminf_{\varepsilon \rightarrow 0} \liminf_{k \rightarrow \infty} d(\vec{\Phi}_k(\Sigma^2 \setminus B_\varepsilon(a^*)), \vec{\Phi}_k(\partial B_\varepsilon(a_i))) = 0. \quad (28)$$

On the other hand, recalling (18), also the lengths of the boundaries converge to zero

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{k \rightarrow \infty} \mathcal{H}^1(\vec{\Phi}_k(\partial B_\varepsilon(a^*))) \rightarrow 0,$$

and combining this last observation with (28) we obtain that also the diameter of the complementary of the bubble has to vanish in the limit:

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{k \rightarrow \infty} \text{diam}(\vec{\Phi}_k(\Sigma^2 \setminus B_\varepsilon(a^*))) = 0 \quad . \quad (29)$$

Observe that (26), respectively (29), implies that the portion of the surface outside the bubble in a^* is not contributing in the limit to the area, respectively to the enclosed volume; therefore it does not contribute to the isoperimetric ratio in the limit. Nevertheless notice that the bubble forming in a^* is topologically (a bigger and bigger portion of) a sphere, therefore the topological information is carried by the shrinking part made by the complementary of the bubble. Summarizing, we are facing a *dichotomy* between a (bigger) portion of the surface (namely, the bubble) carrying the isoperimetric ratio, and a (smaller) portion (the complementary of the bubble) carrying the topological information. In the next subsection we isolate the two parts by performing a “cut and fill” procedure.

4.3.2 Cut and fill

As described above the geometric situation can be described as follows (for k sufficiently large): there is a dichotomy between the topological information and the additional constraint of prescribed isoperimetric ratio. More precisely, $\vec{\Phi}_k(B_\varepsilon(a^*))$ forms a spherical bubble carrying the isoperimetric information and $\vec{\Phi}_k(\Sigma^2 \setminus B_\varepsilon(a^*))$ keeps the topological information, i.e. it has the topological type of Σ^2 , i.e. a genus $g \geq 1$ surface.

The strategy now is to find estimates for the Willmore energy for the two parts $\vec{\Phi}_k(\Sigma^2 \setminus B_\varepsilon(a^*))$ and $\vec{\Phi}_k(B_\varepsilon(a^*))$ and bring these estimates to a contradiction with our additional hypothesis relating the Willmore energy of Schygulla-spheres to the Willmore energy of our embeddings $\vec{\Phi}_k$.

In doing so, we will exploit the existence of a genus $g \geq 1$ minimizers of the Willmore energy among all genus g embedded surfaces (free minimization), the existence of Schygulla-spheres $\mathbb{S}_{S,r}$, i.e. smoothly embedded surfaces of type \mathbb{S}^2 minimizing the Willmore energy for the given isoperimetric ratio r , and the fact that the function which assigns to each given isoperimetric ratio the Willmore energy of the Schygulla-sphere for this given isoperimetric ratio is strictly monotone and continuous. These latter fact are proved in [57].

In order to perform the above strategy, we will apply a cut-and-fill-procedure in order to close each of the parts $\vec{\Phi}_k(\Sigma^2 \setminus B_\varepsilon(a^*))$ and $\vec{\Phi}_k(B_\varepsilon(a^*))$, complete

them separately to new closed surfaces and finally estimate appropriately their Willmore energy. For this procedure, we will use the following adaptation of a lemma which can be found (with proof) in [38].

Lemma 4.9 *Let $\vec{\Phi}_k$ be a sequence of conformal weak, immersions $\{\vec{\Phi}_k\} \subset \mathcal{E}_{\Sigma^2}$ into \mathbb{R}^3 . Assume that*

$$\limsup_{k \rightarrow \infty} \int_{\Sigma^2} [1 + |D\vec{n}_{\vec{\Phi}_k}|^2] dvol_{g_k} < \infty.$$

Let $a \in \Sigma^2$ and $s_k, t_k \rightarrow 0$ such that

$$\frac{t_k}{s_k} \rightarrow 0$$

and

$$\lim_{k \rightarrow \infty} \int_{B_{s_k}(a) \setminus B_{t_k}(a)} [1 + |\mathbb{I}_{\vec{\Phi}_k}|^2] dvol_{g_k} = 0.$$

Then there exist conformal immersions $\vec{\xi}_k$ from Σ^2 into \mathbb{R}^3 and a sequence of quasi conformal bilipschitz homeomorphisms ψ_k of Σ^2 , converging in C^0 -norm over Σ^2 to the identity map, such that

$$\vec{\xi}_k \circ \psi_k = \vec{\Phi}_k \text{ in } \Sigma^2 \setminus B_{s_k}(a)$$

and

$$\lim_{k \rightarrow \infty} \text{diam}(\vec{\xi}_k \circ \psi_k(B_{s_k}(a))) = 0, \quad \lim_{k \rightarrow \infty} \text{Area}(\vec{\xi}_k \circ \psi_k(B_{s_k}(a))) = 0.$$

Moreover

$$\lim_{k \rightarrow \infty} \int_{B_{s_k}(a)} |\mathbb{I}_{\vec{\xi}_k \circ \psi_k}^0|^2 dvol_{g_{\vec{\xi}_k \circ \psi_k}} = 0$$

where $\mathbb{I}_{\vec{\xi}_k \circ \psi_k}^0$ is the trace free second fundamental form.

Remark 4.10 • Note that due to the assumption of bounded Willmore energy

$$\inf_k W(\vec{\Phi}_k) < 8\pi$$

the hypothesis of the above lemma are satisfied, in particular it is possible to find radii s_k and t_k such that

$$\lim_{k \rightarrow \infty} \int_{B_{s_k}(a^*) \setminus B_{t_k}(a^*)} [1 + |\mathbb{I}_{\vec{\Phi}_k}|^2] dvol_{g_k} = 0.$$

- At first glimpse, one might have the impression that by the cut-and-fill-procedure guaranteed by the previous lemma we can close just one of the two parts of our surface, either $\vec{\Phi}_k(\Sigma^2 \setminus B_\varepsilon(a^*))$ or $\vec{\Phi}_k(B_\varepsilon(a^*))$. But the proof of the above cited result actually reveals that the procedure allows to close both parts. More precisely, depending on the side from which we approach the curve along which we cut (from “inside” or from “outside”), we have two possibilities of closing the surface: either we glue almost an entire shrinking sphere Σ_{1,s_k} with Willmore energy $W(\Sigma_{1,s_k}) \rightarrow 4\pi$ as $s_k \rightarrow 0$ or we glue a shrinking almost flat disk Σ_{2,s_k} with Willmore energy $W(\Sigma_{2,s_k}) \rightarrow 0$ as $s_k \rightarrow 0$.

- The only remaining risk would be that we create branch point. From the proof of the original version of the above lemma in [38] we see that this can be excluded once we can show that in

$$\Delta \hat{\lambda}_\infty = c_0 \delta_0$$

it holds $c_0 = 0$.

In the above equation, $\hat{\lambda}_\infty$ is the limit of the conformal factors $\hat{\lambda}_k = \log |\partial_{x_1} \hat{\Phi}_k| = \log |\partial_{x_2} \hat{\Phi}_k|$ where

$$\hat{\Phi}_k := e^{-C_k} (\vec{\Phi}_k - \vec{\Phi}_k(x_0))$$

for a suitable x_0 , and δ_0 denotes the Dirac delta distribution centered at the origin. We claim that, in the present framework, actually we have $c_0 = 0$. Indeed, by assumption, recall that there exists $\delta > 0$ such that

$$W(\vec{\Phi}_k) \leq 8\pi - \delta \text{ for all } k. \quad (30)$$

If we had $c_0 \neq 0$, $\hat{\Phi}_\infty$ had a branch point of order $\frac{c_0}{2\pi}$. In other words, we had that $\hat{\Phi}_\infty$ covers $\frac{c_0}{2\pi} + 1$ times the plane P_0^2 . Due to the fact that for any given $\varepsilon > 0$, on $\Sigma^2 \setminus B_\varepsilon(a^*)$ we have weak $W^{2,2}$ -convergence of $\hat{\Phi}_k$ to $\hat{\Phi}_\infty$, then for any choice of $0 < 2\alpha < \beta < \varepsilon$

$$\hat{\Phi}_k \rightarrow \hat{\Phi}_\infty \text{ in } C^{0,\gamma}(B_\beta(a^*) \setminus B_{2\alpha}(a^*))$$

by the classical Sobolev embedding theorem.

In this situation, we select a point $p \in B_\beta(a^*) \setminus B_{2\alpha}(a^*)$ and a radius η small enough such that

$$B_\eta(p) \subset B_\beta(a^*) \setminus B_{2\alpha}(a^*)$$

and we apply Lemma 5.6 in the Appendix to $\vec{\Phi}_k(\Sigma^2)$ and a ball $B_\rho \subset \mathbb{R}^3$ such that $\vec{\Phi}_k(B_\eta(p)) \subset B_\rho$ in order to conclude that - for η and ρ small enough -

$$\limsup_k W(\vec{\Phi}_k) \geq 8\pi - \frac{\delta}{2}.$$

Since this contradicts our hypothesis (30), we must have $c_0 = 0$ and therefore the immersions we created via Lemma 4.9 are unbranched. \square

In a first step, applying Lemma 4.9, we get the following equalities for k large enough

$$\begin{aligned} W(\vec{\Phi}_k) &= W(\vec{\Phi}_k(\Sigma^2 \setminus B_{s_k}(a^*))) + W(\vec{\Phi}_k(B_{s_k}(a^*))) \\ &= W(\vec{\xi}_k \circ \psi_k(\Sigma^2 \setminus B_{s_k}(a^*))) + W(\vec{\Phi}_k(B_{s_k}(a^*))) \end{aligned}$$

Now, in the present dichotomy situation we have the following two possibilities of filling: either we glue an almost entire sphere Σ_{1,s_k} to $\vec{\xi}_k \circ \psi_k(\Sigma^2 \setminus B_{s_k}(a^*))$ and an almost flat disc Σ_{2,s_k} to $\vec{\Phi}_k(B_{s_k}(a^*))$ or viceversa.

In the first case we can estimate

$$\begin{aligned}
W(\vec{\Phi}_k) &= W(\vec{\xi}_k \circ \psi_k(\Sigma^2 \setminus B_{s_k}(a^*))) + W(\vec{\Phi}_k(B_{s_k}(a^*))) \\
&= W(\vec{\xi}_k \circ \psi_k(\Sigma^2 \setminus B_{s_k}(a^*)) \cup \Sigma_{1,s_k}) - W(\Sigma_{1,s_k}) \\
&\quad + W(\vec{\Phi}_k(B_{s_k}(a^*)) \cup \Sigma_{2,s_k}) - W(\Sigma_{2,s_k}) \\
&\geq \beta_g^3 - W(\Sigma_{1,s_k}) + W(\vec{\Phi}_k(B_{s_k}(a^*)) \cup \Sigma_{2,s_k}) - W(\Sigma_{2,s_k}) \\
&\quad \text{since, by construction, } \vec{\xi}_k \circ \psi_k(\Sigma^2 \setminus B_{s_k}(a^*)) \cup \Sigma_{1,s_k} \text{ is a genus } g \text{ surface} \\
&\geq \beta_g^3 - 4\pi - \varepsilon_1(s_k) + W(\vec{\Phi}_k(B_{s_k}(a^*)) \cup \Sigma_{2,s_k}) - W(\Sigma_{2,s_k}) \\
&\quad \text{by the energy estimate for } \Sigma_{1,s_k} \text{ of Remark 4.10} \\
&\geq \beta_g^3 - 4\pi - \varepsilon_1(s_k) + W(\mathbb{S}_{S,iso}(\vec{\Phi}_k(B_{s_k}(a^*)) \cup \Sigma_{2,s_k})) - W(\Sigma_{2,s_k}) \\
&\quad \text{since } \vec{\Phi}_k(B_{s_k}(a^*)) \cup \Sigma_{2,s_k} \text{ has the topology of a sphere} \\
&\geq \beta_g^3 - 4\pi - \varepsilon_1(s_k) + W(\mathbb{S}_{S,iso}(\vec{\Phi}_k(B_{s_k}(a^*)) \cup \Sigma_{2,s_k})) - \varepsilon_2(s_k) \\
&\quad \text{by the energy estimate for } \Sigma_{2,s_k} \text{ of Remark 4.10} \quad .
\end{aligned}$$

Observe that as s_k tend to 0, the two terms $\varepsilon_1(s_k)$ and $\varepsilon_2(s_k)$ vanish. Moreover, exploiting the fact that

$$t \mapsto \beta(t) := W(\mathbb{S}_{S,t})$$

is continuous in t (see [57]) together with the observation that (here we use that $\vec{\Phi}_k(B_{s_k}(a^*))$ carries the isoperimetric information of $\vec{\Phi}_k$, and the isoperimetric contribution of Σ_{2,s_k} is negligible in the limit)

$$\lim_{k \rightarrow \infty} iso(\vec{\Phi}_k(B_{s_k}(a^*)) \cup \Sigma_{2,s_k}) = R \quad ,$$

we conclude that

$$W(\mathbb{S}_{S,iso}(\vec{\Phi}_k(B_{s_k}(a^*)) \cup \Sigma_{2,s_k})) \rightarrow W(\mathbb{S}_{S,R}).$$

In the second case (i.e. the shrinking almost entire sphere Σ_{1,s_k} is glued to the bubble $\vec{\Phi}_k(B_{s_k}(a^*))$, and the almost flat disc Σ_{2,s_k} is glued to $\vec{\xi}_k \circ \psi_k(\Sigma^2 \setminus B_{s_k}(a^*))$) the analogous estimate bring to the same conclusions.

So in the limit we find

$$\liminf W(\vec{\Phi}_k) \geq W(T_{Clifford}) + W(\mathbb{S}_{S,R}) - 4\pi,$$

which contradicts that $R \in I_g$, and I_g is defined as in (5); therefore the case $C_k \rightarrow -\infty$ cannot occur. Since the only remaining case is when $\{C_k\}$ is bounded, and in Subsection 4.2 we already proved the existence of a minimizer under this assumption, the proof of Theorem 1.2 is now complete. \square

5 Proof of Theorem 1.4

First of all, we will show that the set I_g is not empty. To this aim recall that for every genus $g \geq 1$ the infimum β_g^3 of the Willmore energy among genus g immersed surfaces in \mathbb{R}^3 is attained by a smooth embedding $\vec{\Phi}_g$ (for genus one

see [60], for higher genus see [3]; for a different proof of the general case see [48]). We claim that $iso(\vec{\Phi}_g) \in I_g$, where I_g is defined in (5).

By a direct comparison with the stereographic projections of the Lawson minimal surfaces of genus g in \mathbb{S}^3 (see e.g. the introduction of [60]), for every $g \geq 1$ one has

$$\beta_g^3 < 8\pi \quad ; \quad (31)$$

moreover, by the work of Bauer and Kuwert [3] we know that

$$\beta_g^3 < \omega_g^3 \quad ; \quad (32)$$

Finally, by the isoperimetric inequality in \mathbb{R}^3 , the isoperimetric ratio of any embedded surface in \mathbb{R}^3 different from a round sphere is strictly larger than $iso(\mathbb{S}^2)$, then

$$iso(\vec{\Phi}_g) > iso(\mathbb{S}^2) \quad . \quad (33)$$

By the fact that for any Schygulla sphere $\mathbb{S}_{S,t}$ different from the round sphere we have $W(\mathbb{S}_{S,t}) > 4\pi$ (see [57]), we then deduce

$$W(\vec{\Phi}_g) = \beta_g^3 < \beta_g^3 - 4\pi + W(\mathbb{S}_{S,r=iso(\vec{\Phi}_g)}) \quad ,$$

which, combined with (31) and (32) yields

$$W(\vec{\Phi}_g) < \min\{8\pi, \omega_g^3, \beta_g^3 - 4\pi + W(\mathbb{S}_{S,r=iso(\vec{\Phi}_g)})\} \quad .$$

It follows that $iso(\vec{\Phi}_g) \in I_g$ as desired.

Before we continue the proof of Theorem 1.4, let us state and prove an easy lemma which we will use later on.

Lemma 5.1 *Look at the following map*

$$\begin{aligned} \Psi : \{ \Sigma^2 : \Sigma^2 \text{ is a smoothly embedded surface in } \mathbb{R}^3 \} &\rightarrow \mathbb{R} \\ \Sigma^2 &\mapsto iso(\Sigma^2). \end{aligned}$$

Then Σ^2 is a critical point of Ψ if and only if Σ^2 is a round sphere.

Proof. Due to the fact that the isoperimetric ratio is invariant under rescaling, we may - without loss of generality - assume that the volume enclosed by Σ^2 is equal to 1.

Thus, Σ^2 is a critical point of Ψ if and only if it is a critical point of the area functional under the constraint of fixed volume.

But it is well known that this leads to the conclusion that Σ^2 has to be surface of constant mean curvature.

And finally by a famous theorem due to H. Hopf and Alexandrov (see for instance [20]), it follows that Σ^2 is a round sphere. Thus the proof of Lemma 5.1 is complete. \square

Now we come back to the proof of Theorem 1.4: given $r \in I_g$, we have to show that there exists $\delta > 0$ such that $(r - \delta, r + \delta) \subset I_g$.

By Theorem 1.2 we know that there exists a smooth embedding $\vec{\Phi}_r : \Sigma^2 \rightarrow \mathbb{R}^3$

minimizing the Willmore energy and respecting the given isoperimetric constraint $iso(\vec{\Phi}_r) = r$.

By Lemma 5.1 we deduce that $\vec{\Phi}_r$ is not a critical point of Ψ , in other words, the differential of Ψ at the point $\vec{\Phi}_r$ is surjective. Therefore, there exists a smooth 1-parameter family $\left\{ \vec{\Phi}_s \right\}_{s \in (r-\varepsilon_0, r+\varepsilon_0)}$ of embeddings - obtained by perturbing $\vec{\Phi}_r$ - such that for every $\varepsilon \in (0, \varepsilon_0)$ there exists $\delta > 0$ such that

$$\Psi \left(\left\{ \vec{\Phi}_s \right\}_{s \in (r-\varepsilon, r+\varepsilon)} \right) \supset (r - \delta, r + \delta). \quad (34)$$

Now, recall that for the embedding $\vec{\Phi}_r$ we started with, we have the following estimate

$$W(\vec{\Phi}_r) < \min \{ 8\pi, \omega_g^3, \beta_g^3 + W(\mathbb{S}_{S,r}) - 4\pi \}.$$

Thus, there exists $\eta > 0$ such that

$$W(\vec{\Phi}_r) \leq \min \{ 8\pi, \omega_g^3, \beta_g^3 + W(\mathbb{S}_{S,r}) - 4\pi \} - \eta.$$

Hence, from the smoothness in s of the family $\left\{ \vec{\Phi}_s \right\}_{s \in (r-\varepsilon, r+\varepsilon)}$, for s close enough to r we have

$$W(\vec{\Phi}_s) \leq \min \{ 8\pi, \omega_g^3, \beta_g^3 + W(\mathbb{S}_{S,r}) - 4\pi \} - \frac{\eta}{2}. \quad (35)$$

In a last step, we exploit the continuity of the function which to a given isoperimetric ratio assigns the Willmore energy of the corresponding Schygulla sphere, more precisely the function $t \mapsto \beta(t) := W(\mathbb{S}_{S,t})$ is continuous in t (see [57] for the proof). We deduce that for s close enough to r we have

$$|W(\mathbb{S}_{S,\Psi(\vec{\Phi}_s)}) - W(\mathbb{S}_{S,r})| \leq \frac{\eta}{4}. \quad (36)$$

Combining (35) and (36) we get

$$\begin{aligned} W(\vec{\Phi}_s) &\leq \beta_g^3 + W(\mathbb{S}_{S,\Psi(\vec{\Phi}_s)}) - 4\pi - \frac{\eta}{4} \\ &< \beta_g^3 + W(\mathbb{S}_{S,\Psi(\vec{\Phi}_s)}) - 4\pi. \end{aligned}$$

This last inequality together with (35) and (34) conclude the proof of Theorem 1.4. \square

Appendix

In this section we recall some useful (technical) results used in the main text.

We start with a result which relates the non-compactness of the conformal classes to an estimate for the Willmore energy from below.

This result can be found in [47] and in [21] (or in [27] for $m = 3, 4$).

Theorem 5.2 *Let (Σ^2, c_k) be a sequence of closed Riemannian surfaces of genus g and conformal classes c_k . Assume that $[c_k]$ is diverging to the boundary of the Moduli Space of the conformal classes on Σ^2 . Let $\{\vec{\Phi}_k\} \subset \mathcal{E}_{\Sigma^2}$ be a sequence of weak immersions, conformal with respect to c_k ; then*

$$\liminf_{k \rightarrow \infty} \int_{\Sigma^2} |\vec{H}_{\vec{\Phi}_k}|^2 d\text{vol}_{\vec{\Phi}_k^* g_{\mathbb{R}^m}} \geq \min\{8\pi, \omega_g^m\}$$

where

$$\omega_g^m := \min \left\{ 4\pi + \sum_{i=1}^p (\beta_{g_i}^m - 4\pi) \mid g = g_1 + \cdots + g_p, 1 \leq g_i < g \right\}$$

and

$$\beta_g^m := \inf \left\{ W(\vec{\Phi}) \mid \vec{\Phi} \text{ is an immersion of the genus } g \text{ closed surface} \right\}.$$

We continue with the famous Wente estimate.

Theorem 5.3 *Let a and b be two function in $W^{1,2}(D^2, \mathbb{R})$. Moreover, let ϕ be the unique solution of*

$$\begin{cases} -\Delta\phi = \nabla a \cdot \nabla^\perp b = \partial_x a \partial_y b - \partial_y a \partial_x b & \text{in } B_1^2(0) \\ \phi = 0 & \text{on } \partial B_1^2(0), \end{cases}$$

Then the following estimates hold

$$\|\phi\|_\infty + \|\nabla\phi\|_{2,1} + \|\nabla^2\phi\|_1 \leq C\|\nabla a\|_2 \|\nabla b\|_2$$

Here, $\|\cdot\|_{2,1}$ denotes the norm of the Lorentz space $L^{2,1}$.

A proof of this result can be found in [15] (see also [69] and [63]).

The next result - which can be found in [47] - gives an estimate of the Willmore energy of a compact surface with boundary. It is a consequence of Simon's monotonicity formula with boundary.

Lemma 5.4 *Let Σ^2 be a compact surface with boundary and let $\vec{\Phi}$ belong to \mathcal{E}_{Σ^2} . Then the following inequality holds*

$$4\pi \leq W(\vec{\Phi}) + 2 \frac{\mathcal{H}^1(\vec{\Phi}(\partial\Sigma^2))}{d(\vec{\Phi}(\partial\Sigma^2), \vec{\Phi}(\Sigma^2))} \quad (37)$$

where $\mathcal{H}^1(\vec{\Phi}(\partial\Sigma^2))$ denotes the 1-dimensional Hausdorff measure of the boundary of the immersion $\vec{\Phi}(\partial\Sigma^2)$ and where $d(\cdot, \cdot)$ denotes the usual distance between two sets in \mathbb{R}^m .

A useful lemma relating area, diameter and Willmore energy of a connected, compact surface without boundary is given by the following lemma due to Simon (see [60]).

Lemma 5.5 *Let $\Sigma^2 \subset \mathbb{R}^m$ be an immersed connected, compact surface without boundary.*

Then it holds

$$\frac{\text{Area}(\Sigma^2)}{W(\Sigma^2)} \leq \text{diam}^2(\Sigma^2) \leq C \text{Area}(\Sigma^2) W(\Sigma^2)$$

for some constant $C > 0$.

Finally, let us recall the following lemma due to Simon (see [60]) giving an estimate for the Willmore energy from below.

Lemma 5.6 *Assume that $\Sigma^2 \subset \mathbb{R}^m$ is an immersed compact surface without boundary, that ∂B_ρ intersects Σ^2 transversely and that $\Sigma^2 \cap B_\rho$ contains disjoint subsets Σ_1, Σ_2 with $\Sigma_j \cap B_{\theta\rho} \neq \emptyset$, $\partial\Sigma_j \subset \partial B_\rho$, and $|\partial\Sigma_j| \leq \beta\rho$ for $j = 1, 2$, $\theta \in (0, \frac{1}{2})$ and $\beta > 0$. Then*

$$W(\Sigma^2) \geq 8\pi - C\beta\theta$$

where C does not depend on Σ , β or θ .

References

- [1] David R. Adams, A note on Riesz potentials, *Duke Math. J.* 42 (1975), no. 4, 765-778
- [2] Robert A. Adams, John J. F. Fournier, *Sobolev spaces*, Elsevier, 2003
- [3] Matthias Bauer, Ernst Kuwert, Existence of Minimizing Willmore Surfaces of Prescribed Genus, *Int. Math. Res. Not.* 10 (2003), 553-576
- [4] Giovanni Bellettini, Luca Mugnai, Approximation of Helfrich's functional via diffuse interfaces, *SIAM J. Math. Anal.* 42 (2010), no. 6, 2402-2433
- [5] Yann Bernard, Tristan Rivière, Local Palais-Smale sequences for the Willmore functional, *Comm. Anal. Geom.* 19 (2011), no. 3, 563-599
- [6] Yann Bernard, Tristan Rivière, Energy Quantization for Willmore Surfaces and Applications, preprint (2011)
- [7] Yann Bernard, Tristan Rivière, Asymptotic Analysis of Branched Willmore Surfaces, to appear in *Pacific J. Math.* (2013)
- [8] Wilhelm Blaschke, "Vorlesungen über Differentialgeometrie und geometrische Grundlagen von Einsteins Relativitätstheorie" III, Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen, Bd. XXIX Differentialgeometrie der Kreise und Kugeln, bearbeitet von Gerhard Thomsen, Springer, 1929
- [9] Robert Bryant, A duality theorem for Willmore surfaces, *J. Differential Geom.* 20 (1984), 23-53

- [10] Sagun Chanillo, Yan Yan Li, Continuity of solutions of uniformly elliptic equations in \mathbb{R}^2 , *Manuscripta Math.* 77 (1992), no. 4, 415-433
- [11] Bang-Yen Chen, Some conformal Invariants of Submanifolds and Their Application, *Boll. Unione Mat. Ital.* 10 (1974), 380-385
- [12] Bang-yen Chen, On an inequality of T. J. Willmore, *Proc. Amer. Math. Soc.* 26 1970 473-479.
- [13] Ralph Chill, Eva Fařangová, Reiner Schätzle, Willmore blow-ups are never compact, *Duke Math. J.* 147 (2009), no. 2, 345-376
- [14] Rustum Choksi, Marco Veroni, Global minimizers for the doubly-constrained Helfrich energy: the axisymmetric case, preprint (2012)
- [15] R. Coifman, P. L. Lions, Y. Meyer, S. Semmes, Compensated compactness and Hardy spaces, *J. Math. Pures Appl.* 72 (1993), 247-286
- [16] David Gilbarg, Neil S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer, 1998
- [17] Frédéric Hélein, *Harmonic maps, conservation laws and moving frames*, Cambridge Tracts in Mathematics, 2002
- [18] Wolfgang Helfrich, Elastic properties of lipid bilayers - theory and possible experiments, *Zeitschrift Für Naturforschung C - A Journal Of Biosciences.* 28. (1973), 693-703
- [19] Jürgen Jost, *Compact Riemann Surfaces*, Springer, 2006
- [20] Wilhelm P. A. Klingenberg, *Riemannian Geometry*, De Gruyter, 1995
- [21] Ernst Kuwert, Yuxiang Li, $W^{2,2}$ -conformal immersions of a closed Riemann Surface into \mathbb{R}^n , *Comm. Anal. Geom.* 20 (2012), no. 2, 313340
- [22] Ernst Kuwert, Reiner Schätzle, The Willmore flow with small initial energy, *J. Differential Geom.* 57 (2001), 409-441
- [23] Ernst Kuwert, Reiner Schätzle, Gradient Flow for the Willmore Functional, *Communications in Analysis and Geometry*, 10 (2002), 307-339
- [24] Ernst Kuwert, Reiner Schätzle, Removability of point singularities of Willmore surfaces, *Ann. of Math.* 160(2004), 315-357
- [25] Ernst Kuwert, Reiner Schätzle, Branch points of Willmore surfaces, *Duke Math. J.* 138 (2007), no. 2, 179-201
- [26] Ernst Kuwert, Yuxiang Li, Reiner Schätzle, The large genus limit of the infimum of the Willmore energy. *Amer. J. Math.* 132 (2010), no. 1, 37-51
- [27] Ernst Kuwert, Reiner Schätzle, Closed surfaces with bound on their Willmore energy, to appear in *Annali della Scuola Normale Superiore di Pisa*, 2011, arXiv:math.DG/1009.5286.
- [28] Ernst Kuwert, Reiner Schätzle, Minimizers of the Willmore functional under fixed conformal class, *J. Differential Geom.* 93 (2013), no. 3, 471-530

- [29] Ernst Kuwert, Reiner Schätzle, The Willmore functional, Topics in modern regularity theory, 1–115, CRM Series, Scuola Normale Superiore Pisa, 2012
- [30] Ernst Kuwert, Andrea Mondino, Johannes Schygulla, Existence of immersed spheres minimizing curvature functionals in compact 3-manifolds, preprint, 2011
- [31] Tobias Lamm, Jan Metzger, Felix Schulze, Foliations of asymptotically flat manifolds by surfaces of Willmore type, Math. Ann. (2011), 350-378
- [32] L. D. Landau, E. M. Lifshitz, Theory of elasticity - Course of theoretical physics, volume 7 - third edition revised and enlarged by E.M. Lifshitz, A.M. Kosevich and L.P. Pitaevskii, Butterwoth-Heinemann 1986
- [33] P. Li, S.-T.Yau, A New Conformal Invariant and its Applications to the Willmore Conjecture and the First Eigenvalue on Compact Surfaces, Inventiones Math. 69 (1982), 269-291
- [34] Reinhard Lipowsky, Erich Sackman, Structure and Dynamics of Membranes, Elsevier, 1995
- [35] Fernando C. Marques, André Neves, Min-Max theory and the Willmore conjecture, preprint (2012)
- [36] Thomas Milcent, Shape derivative of the Willmore functional and applications to equilibrium shapes of vesicles, Rapport de Recherche, 2011
- [37] Andrea Mondino, Some results about the existence of critical points for the Willmore functional, Math. Z. 266 (2010), no. 3, 583-62
- [38] Andrea Mondino, Tristan Rivière, Immersed Spheres of Finite Total Curvature into Manifolds, preprint (2012)
- [39] Andrea Mondino, Tristan Rivière, Willmore spheres in compact Riemannian manifolds, Advances Math. 232 (2013), no. 1, 608-676
- [40] S. Müller, V. Sverak, On surfaces of finite total curvature, J. Differential Geom. 42 (1995), 229-258
- [41] M. Mutz, D. Benisom, Observation of toroidal vesicles, Phys. Rev. Letter A 43 (1991), 4525-4528
- [42] Mark A. Peletier, Matthias Röger, Partial Localization, Lipid Bilayers, and the Elastica Functional, Arch. Rational Mech. Anal. 193 (2009), 475-537
- [43] Christian Renken, Gino Siragusa, Guy Perkins, Lance Washington, Jim Nulton, Peter Salamon, Terrence G. Frey, A thermodynamic model describing the nature of the crista junction: a structural motif in the mitochondrion, Journal of Structural Biology 138 (2002), 137-144
- [44] Tristan Rivière, Conservation laws for conformal invariant variational problems, Invent. Math., 168 (2007), 1-22
- [45] Tristan Rivière, Analysis aspects of Willmore surfaces, Inventiones Math., 174 (2008), no.1, 1-45

- [46] Tristan Rivière, Sequences of smooth global isothermic immersions, *Comm. P. D. E.* 38 (2013), no. 2, 276-303
- [47] Tristan Rivière, Lipschitz conformal immersions from degenerating Riemann surfaces with L^2 -bounded second fundamental forms, *Adv. Calc. Var.* 6 (2013), no. 1, 1-31
- [48] Tristan Rivière, Variational Principles for immersed Surfaces with L^2 -bounded Second Fundamental Form, *J. für die reine und angewandte Mathematik (Crelles journal)* 2013
- [49] Tristan Rivière, The role of conservation laws in the analysis of conformally invariant problems, in *Topics in Modern Regularity Theory*, G. Mingione Ed., CRM series 13, Edizioni Della Normale, 2012
- [50] Tristan Rivière, Conformally Invariant Variational Problems, in preparation
- [51] Tristan Rivière, Error analysis for the Willmore-Helfrich Functional, Mini-workshop: Mathematics of Biological Membranes, Oberwolfach report 41/2008, 2305-2309
- [52] Tristan Rivière, Michael Struwe, Partial regularity for harmonic maps, and relates problems, *Comm. Pure Appl. Math.* 61 (2008), no. 4, 451-463
- [53] Matthias Röger, Reiner Schätzle, Control of the isoperimetric deficit by the Willmore deficit, *Analysis (Munich)* 32 (2012), no. 1, 17
- [54] Thomas Runst, Winfried Sickel, Sobolev spaces of fractional order, Nemytskij operators, and nonlinear partial differential equations, De Gruyter, 1996
- [55] Reiner Schätzle, Lower semicontinuity of the Willmore functional for currents. *J. Differential Geom.* 81 (2009), no. 2, 437-456
- [56] Reiner Schätzle, The Willmore boundary problem. *Calc. Var. Partial Differential Equations* 37 (2010), no. 3-4, 275-302
- [57] Johannes Schygulla, Willmore minimizers with prescribed isoperimetric ratio, *Arch. Ration. Mech. Anal.* 203 (2012), no. 3, 901-941
- [58] U. Seifert, R. Lipowsky, Morphology of vesicles, *Handbook of Biological Physics*, Elsevier, 1995
- [59] Yoko Shibata, Tom Shemesh, William A. Prinz, Alexander F. Palazzo, Michael M. Kozlov, Tom A. Rapoport, Mechanisms Determining the Morphology of the Peripheral ER, *Cell* 143 (2010), 775-788
- [60] Leon Simon, Existence of surfaces minimizing the Willmore functional, *Comm. Anal. Geom.* 1 (1993), 281-326
- [61] Elias M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, 1970
- [62] Elias M. Stein, *Harmonic analysis*, Princeton University Press, 1993

- [63] Luc Tartar, Remarks on oscillations and Stokes equation, in: "Macroscopic modeling of turbulent flows", Lect. Notes in Physics 230, Springer 1985, 24-31
- [64] Michael E. Taylor, Partial Differential Equations I-III, Springer, 1996
- [65] Peter Topping, Towards the Willmore conjecture, Calc. Var., 11 (2000), 361-393
- [66] Peter Topping, An Approach to the Willmore Conjecture, in Global theory of minimal surfaces, ed. David Hoffman. Clay math. proc., vol. 2, AMS (2005), 769-77
- [67] Karen K. Uhlenbeck, Connections with L^p bounds on curvature, Comm. Math. Phys. 83 (1982), no. 1, 31-42
- [68] Joel L. Weiner, On a problem of Chen, Willmore et al., Indiana University mathematics journal, 27 (1978), 19-35
- [69] Henry Wente, An existence theorem for surfaces of constant mean curvature, J. Math. Anal. Appl. 26 (1969), 318-344
- [70] Thomas J. Willmore, Riemannian Geometry, Oxford University Press, 1993
- [71] Thomas K. Willmore, Total curvature in Riemannian geometry, 1982
- [72] Ou-Yang Zhong-can, Wolfgang Helfrich, Instability and Deformation of a Spherical Vesicle by Pressure, Phys. Rev. Lett, 59 no. 21, 1987