

Finding Clusters of Primes, II

Progress Report 2006 - 2015

Jörg Waldvogel and Peter Leikauf
Seminar for Applied Mathematics SAM
Swiss Federal Institute of Technology ETH, CH-8092 Zürich

February 2007, May 2013, April 2015

1 Review of Earlier Results.

This project was initiated in the year 2000 with the development of the software system PMP, “Poor Man’s Parallelizer” by Peter Leikauf. PMP is a simple but robust system for parallelizing large computational efforts for workstations connected to the internet and/or for cluster computers. It is particularly well suited for easily parallelizable tasks with low data traffic. For details see the link [Projects/pmp.pdf](#) on the website [10].

In the current application of our parallelization software we are using an algorithm involving sieving techniques for locating and counting clusters of prime numbers. Whereas the distribution of primes seems to be fairly regular, the distribution of twin primes and longer clusters is largely unknown and is characterized by large-scale anomalies. Collecting experimental data on these anomalies is one of the reasons for the interest in clusters of primes. Research in the theory of prime numbers has again become fashionable with the invention of the RSA coding scheme [9] in 1978, and a world-wide competition for prime number records is still going on [7].

Another challenge of finding clusters of prime numbers is the unproven prime k -tuple hypothesis, which is concerned with patterns of natural numbers that occur repeatedly with all elements being prime. The hypothesis states that any pattern that is not forbidden by simple divisibility consider-

ations occurs infinitely often in the sequence of primes. Collecting data in support of this hypothesis is our current goal.

The discovery of a dense cluster of 18 primes among 71 consecutive integers in the range of $3 \cdot 10^{24}$ on November 13, 2000 was the first success of the project. This computation is about 50 times harder than finding the three 23-digit clusters of 17 primes among 67 consecutive integers, discovered by Tony Forbes [3], and independently by J. Waldvogel [10], both in 1998. Our discovery was immediately announced in the number theory press [6], and also got coverage in the web journal of ETH (ETHLife, December 6, 2000, [2]). For previous reports and more details see the links

[Projects/clprimes01.pdf](#)

[Projects/cl18.pdf](#)

[Projects/clprimes03.pdf](#)

[Projects/paricode.gp](#)

on the website [10]. Dense clusters of primes receive particular attention on the website [4], continuously actualized by Tony Forbes. The successes of our implementation bear the danger of monopolizing this site and taking away all the fun!

2 Some Particular Patterns

In this section we pick a few “exotic” admissible patterns with no obvious prime instances. If the Hardy-Littlewood (HL) estimates indicate any chance of reoccurrences in the range accessible with the current hardware and software, we tried to locate a few prime instances of the patterns.

We will consider four patterns of 16 or 18 elements, all of them being tight repetitions of well-known short patterns, as well as a remarkable “quadratic” pattern of 21 primes. In all cases the search for prime instances was successful, thus accumulating more experimental data on the prime k -tuple hypothesis. To begin, we briefly state this famous conjecture in its strong form put forth by Hardy and Littlewood [5].

Let $\mathbf{c} = [c_1, c_2, \dots, c_k]$ be a monotonically increasing sequence of $k \in \mathbb{N}$ integers, exclusively odd or exclusively even, called a *pattern*. The counting function $\pi_{\mathbf{c}}(x)$ of the pattern \mathbf{c} is defined as the number of shifts s , $0 \leq s \leq x$, such that every element of $\mathbf{c}_s := [s + c_1, s + c_2, \dots, s + c_k]$ is a prime (called a *prime instance* of \mathbf{c}). The conjecture states the asymptotic relationship

$$\pi_{\mathbf{c}}(x) \sim h_{\mathbf{c}} \cdot R_k(x) \text{ as } x \rightarrow \infty.$$

Instead of the conventional logarithmic integrals we prefer to use the generalized Riemann functions

$$R_k(x) := \sum_{j=0}^{\infty} \frac{(\log x)^j}{(k-1+j)! j \zeta(1+j)}.$$

The number $h_{\mathbf{c}}$ is the Hardy-Littlewood constant of the pattern \mathbf{c} ,

$$h_{\mathbf{c}} := 2^{k-1} \cdot \prod_q \frac{1 - q^{-1} r_{\mathbf{c}}(q)}{(1 - q^{-1})^k},$$

where the product is taken over all odd primes q , and $r_{\mathbf{c}}(q)$ is the number of distinct residue classes mod q in the pattern \mathbf{c} . Note that for $k = 1$ we have $h_{[0]} = 1$, and non-admissible patterns \mathbf{c} yield $h_{\mathbf{c}} = 0$. Despite the slow convergence of the above infinite product Hardy-Littlewood constants may be efficiently computed to arbitrary precision by means of the prime zeta function, see, e.g. [8].

2.1 Nine Close Prime Twins

Except for the singular situation of the overlapping twins (3,5) and (5,7), two twin primes may only occur at a distance of $6n - 2$, $n = 1, 2, \dots$, as is seen by a simple consideration of residue classes mod 3. The closest admissible arrangement of 9 prime twins spans an interval of length 104 and comes in 4 types represented by the patterns

$$\begin{aligned} \mathbf{c}_1 &= [-31, -29, -19, -17, -13, -11, 11, 13, 17, 19, 29, 31, 41, 43, 59, 61, 71, 73] \\ \mathbf{c}_2 &= [-43, -41, -31, -29, -19, -17, -13, -11, -1, 1, 11, 13, 17, 19, 41, 43, 59, 61] \end{aligned}$$

and their mirror images \mathbf{c}_1' and \mathbf{c}_2' . The Hardy-Littlewood constants are found to be

$$h_1 = 23041978.71272, \quad h_2 = 18433582.97017 = \frac{4}{5} \cdot h_1;$$

the corresponding estimated frequencies $\text{HL}_1(x)$, $\text{HL}_2(x)$ for the patterns $\mathbf{c}_1, \mathbf{c}_2$, respectively, are

| x | 10^{23} | $5 \cdot 10^{23}$ | 10^{24} | $2 \cdot 10^{24}$ |
|------------------|-----------|-------------------|-----------|-------------------|
| $\text{HL}_1(x)$ | 0.33 | 0.94 | 1.50 | 2.38 |
| $\text{HL}_2(x)$ | 0.26 | 0.75 | 1.20 | 1.90 |

We extended the search up to 10^{24} , except for the case \mathbf{c}_2' , where the range had to be doubled. The initial primes of the existing 9 instances are given in the table below; the total HL count is 6.1. We also indicate the differences in the patterns; the occurrence of an additional prime between two twins is indicated by a plus sign +.

| | Initial Prime | Differences in the Pattern | | | | | | | | | | | | | | | | |
|-----------------|---------------------------|----------------------------|-----|---|------|---|----|---|----|---|----|---|----|---|----|---|------|---|
| \mathbf{c}_1 | | (to 1e24) | | | | | | | | | | | | | | | | |
| 1 | 39582971901830749382519 | 2 | 10 | 2 | 4 | 2 | 22 | 2 | 4 | 2 | 10 | 2 | 10 | 2 | 16 | 2 | 10 | 2 |
| 2 | 85002283332977673203069 | 2 | 6+4 | 2 | 4 | 2 | 22 | 2 | 4 | 2 | 10 | 2 | 10 | 2 | 16 | 2 | 10 | 2 |
| 3 | 111871787983521806791079 | 2 | 10 | 2 | 4 | 2 | 22 | 2 | 4 | 2 | 10 | 2 | 10 | 2 | 16 | 2 | 10 | 2 |
| 4 | 237848512839161942948759 | 2 | 10 | 2 | 4 | 2 | 22 | 2 | 4 | 2 | 10 | 2 | 10 | 2 | 16 | 2 | 10 | 2 |
| \mathbf{c}_1' | | (to 1e24) | | | | | | | | | | | | | | | | |
| 1 | 35902987875008630158997 | 2 | 10 | 2 | 16 | 2 | 10 | 2 | 10 | 2 | 4 | 2 | 22 | 2 | 4 | 2 | 10 | 2 |
| 2 | 681954129502801901959427 | 2 | 10 | 2 | 16 | 2 | 10 | 2 | 10 | 2 | 4 | 2 | 22 | 2 | 4 | 2 | 10 | 2 |
| \mathbf{c}_2 | | (to 1e24) | | | | | | | | | | | | | | | | |
| 1 | 670962238726376003928317 | 2 | 10 | 2 | 10 | 2 | 4 | 2 | 10 | 2 | 10 | 2 | 4 | 2 | 22 | 2 | 10+6 | 2 |
| 2 | 785878473089354651160797 | 2 | 10 | 2 | 10 | 2 | 4 | 2 | 10 | 2 | 10 | 2 | 4 | 2 | 22 | 2 | 16 | 2 |
| \mathbf{c}_2' | | (to 2e24) | | | | | | | | | | | | | | | | |
| 1 | 1063660630652819772482009 | 2 | 16 | 2 | 18+4 | 2 | 4 | 2 | 10 | 2 | 10 | 2 | 4 | 2 | 10 | 2 | 10 | 2 |

2.2 Four Close Quadruplets

A striking short pattern is the arrangement of two twins in the same decade, such as $[11, 13, 17, 19]$, referred to as a *prime quadruplet*. Its HL constant is $h_4 = 4.1511808632$; in a large interval Δ near a much larger x we expect to find about $h_4 \cdot \Delta / (\log x)^4$ quadruplets.

Prime quadruplets may accumulate as closely as in the pattern $\mathbf{c}_8 = [-19, -17, -13, -11, 11, 13, 17, 19]$, since \mathbf{c}_8 is admissible. With its HL constant $h_8 = 288.1001283176$, \mathbf{c}_8 is expected to reoccur in terms of positive primes in the range of 4 000 000. In fact, the first pair of quadruplets occurs somewhat prematurely at 1 006 301, and up to $x = 2 \cdot 10^8$ there are 10 instances, whereas the HL estimate is 6.

Two pairs \mathbf{c}_8 of quadruplets in their closest arrangement are separated by a gap of length 382, i.e. the pattern

$$\mathbf{c}_{16} = [-229, -227, -223, -221, -199, -197, -193, -191, 191, 193, 197, 199, 221, 223, 227, 229]$$

is admissible. In this case, however, it is not possible to represent the pattern by a sequence of small primes: $221 = 13 \cdot 17!$ The HL constant is found to be

$h_{16} = 9350784.9154303126$. A search up to $x = \frac{4}{3} \cdot 10^{19}$ turned up two fairly close occurrences near $1.1 \cdot 10^{19}$ and $1.2 \cdot 10^{19}$, whereas the Hardy-Littlewood count yields $HL(x) = h_{16} \cdot R_{16}(x) = 1.003$. Below we show the initial primes of these instances; we again indicate the sequences of differences in the pattern; Q stands for [2,4,2], and a plus sign + indicates additional primes in the large gaps of the pattern.

| Initial Prime | | | Differences in the Pattern | | |
|----------------------|---|----|----------------------------|----------------------------|--------|
| 11281963036964038421 | Q | 22 | Q | 42+42+196+12+2+70+18 | Q 22 Q |
| 12114914563464663491 | Q | 22 | Q | 22+32+28+8+34+14+54+162+28 | Q 22 Q |

Four quadruplets in their *closest* arrangement span an interval of only 218 and occur in the pattern

$$\mathbf{C}_{16} = [-19, -17, -13, -11, 11, 13, 17, 19, 101, 103, 107, 109, 191, 193, 197, 199]$$

or in its mirror image \mathbf{C}_{16}' ; the common HL constant is $H_{16} = 6042367.03907$. For $x = \frac{4}{3} \cdot 10^{19}$ the Hardy-Littlewood count yields 0.648 for each pattern. Both patterns have early first occurrences, namely \mathbf{C}_{16} at the HL count of 0.262, \mathbf{C}_{16}' even at 0.066. In the table below we use the same coding scheme as above.

| Patt. | Initial Prime | | | Differences in the Pattern | | | | |
|-------|---------------------|---|-----------------|----------------------------|---------|----|----------------|---|
| C16 | 3051450534439926131 | Q | | 22 | Q | 82 | Q 12+16+6+8+40 | Q |
| C16' | 300000224101777931 | Q | 10+8+4+26+12+22 | Q | 4+54+24 | Q | 12+10 | Q |

2.3 Three Close Sixtuplets and Two Close 9-Tuplets

Being able to detect occurrences of clusters of 18 primes, we have a chance of displaying instances of three sixtuplets [97, 101, 103, 107, 109, 113] as well as two 9-tuplets [13, 17, 19, 23, 29, 31, 37, 41, 43] in their closest arrangements. The patterns are \mathbf{c}_6 and \mathbf{c}_9 below, spanning intervals of length 436 and 90, having the HL constants $h_6 = 92680700.00031$ and $h_9 = 19311372.63543$, respectively.

$$\begin{aligned} c_6 &= [-113, -109, -107, -103, -101, -97, 97, 101, 103, 107, 109, 113, 307, 311, 313, 317, 319, 323] \\ c_9 &= [-47, -43, -41, -37, -31, -29, -23, -19, -17, 13, 17, 19, 23, 29, 31, 37, 41, 43]. \end{aligned}$$

As it happened in the pattern \mathbf{c}_{16} of Section 2.2, \mathbf{c}_6 cannot be represented in terms of small primes: $319 = 11 \cdot 29$, $323 = 17 \cdot 19$! The two patterns were searched up to 10^{23} , corresponding to HL counts of 1.33 and 0.277; at the first occurrences given below the actual HL counts are 0.85 and 0.0026, respectively. We were lucky to capture the extremely early occurrence of \mathbf{c}_9 !

| Patt. | Initial Prime | Differences in the Pattern | |
|-------|-------------------------|----------------------------|--|
| c6 | 50038627250687303646277 | 4 2 4 2 4 | 56+138 4 2 4 2 4 38+90+10+56 4 2 4 2 4 |
| c9 | 54014646858393564377 | 4 2 4 6 2 6 4 2 | 30 4 2 4 6 2 6 4 2 |

2.4 21 Primes in a Quadratic Pattern

Leonhard Euler (1707-1783), the mathematical giant of the eighteenth century, made the stunning discovery that the quadratic polynomial $f(n) := n^2 + n + c$ with $c = 41$ has only prime values for $n = 0, 1, 2, \dots, 39$. Also for $n \geq 40$ prime values of $f(n)$ are extraordinarily abundant. One of the reasons for this is the obvious symmetry relation

$$f(n) \bmod p = f(p - n - 1) \bmod p \quad \text{for every prime } p.$$

This implies that a finite pattern $[f(n), n = 0, 1, \dots, N-1]$ of arbitrary length N occupies at most $\text{ceil}(p/2)$ residue classes modulo any prime p . Therefore at least $\text{floor}(p/2)$ residue classes mod p are empty, hence the pattern is admissible.

As a consequence, *quadratic patterns*

$$[n^2 + n + c, \quad n = 0, 1, \dots, N - 1]$$

of N elements are candidates to yield prime instances of rather long patterns of this preassigned structure. In fact, it was established that the first reoccurrence of the quadratic pattern

$$\mathbf{C} = [41, 43, 47, 53, 61, 71, 83, 97, 113, 131, 151, 173, 197, 223, 251, 281, 313, 347, 383, 421, 461]$$

of $N = 21$ (non-consecutive) primes begins at

$$x = 2\,34505\,01594\,32353\,29417.$$

The HL constant of \mathbf{C} is found to be $H = 3916679971309.28168$, and for the HL count we obtain $H \cdot R_{21}(x) = 0.0139$, hence we again profited from a rather early occurrence.

It is unlikely that the linearly growing sequence $[2, 4, 6, 8, \dots]$ persists throughout the entire pattern as the sequence of differences between consecutive primes. Rather, it will occasionally be broken by additional primes in the ever longer gaps of the pattern. The above instance has the difference pattern

$$2\ 4\ 6\ 8\ 10\ 12\ 14\ 16\ 18\ 20\ 4+18\ 14+6+4\ 26\ 10+18\ 30\ 32\ 22+12\ 36\ 24+14\ 40$$

where plus signs + again denote additional primes interrupting the gaps of the preassigned pattern.

If the prime k-tuple hypothesis is true, instances of \mathbf{C} in terms of *consecutive* primes must exist. To locate one of these, however, is beyond the possibilities of our current technique.

3 Migrating from Asgard to Hreidar

After almost five years of continuous operation on Asgard the more powerful hardware Hreidar became available in Spring 2005. Migration from the ailing Asgard to Hreidar, including tests of the adapted software, took place from May 11 to May 18, 2005. The authors thank the steering committee for again generously allotting idle computing power to this project. We also express our sincere thanks to George Sigut for his competent help during the migration process.

Our system consisting of the parallelizing software PMP and the searching algorithm (coded in PARI [1], the innermost loop in C++) proved to be very easy to migrate. E.g., the doubling of the word length from 32 to 64 bits could be handled by merely redefining a parameter called `bits`.

Migration took place while a long run was under way: the first block of length 10^{24} for the pattern C_2 , see Section 4 below. No data were lost; after migration, and also after a subsequent change of configuration, the search resumed exactly where it was stopped. The only losses (besides the week of migration) were the 337 tasks (or 0.07 %) that were interrupted. More details of the long run are given in the table below.

| Dates 2005 | Days | % | Tasks | % | Tsk/Day | Machine | Nodes |
|---------------|-------|-------|--------|--------|---------|---------|-------|
| 01/06 - 05/11 | 124.9 | 62.0 | 250425 | 50.17 | 2005 | Asgard | 100 |
| 05/18 - 07/19 | 61.9 | 30.7 | 95686 | 19.17 | 1545 | Hreidar | 32 |
| 07/19 - 08/03 | 14.8 | 7.3 | 153426 | 30.73 | 10362 | Hreidar | 32+32 |
| Total | 201.6 | 100.0 | 499537 | 100.07 | 2476 | | |

4 The Quest for a Maximally Dense 19-Tuple

Both of the patterns

$$\mathbf{c}_1 = [13, 17, 19, \dots, 79, 83, 89], \quad \mathbf{c}_2 = [37, 41, 43, \dots, 107, 109, 113],$$

consisting of 19 consecutive primes and encompassing 77 consecutive natural numbers each, are admissible, i.e. their elements leave at least one residue class unoccupied modulo every prime. Together with their mirror images $\mathbf{c}'_1 = [-89.. -13]$ and $\mathbf{c}'_2 = [-113.. -37]$ they constitute the four different patterns in which 19 primes can be arranged as densely as possible. The Hardy-Littlewood constants associated with \mathbf{c}_1 (or \mathbf{c}'_1) and \mathbf{c}_2 (or \mathbf{c}'_2) are found to be

$$h_1 = 172\,98850.522, \quad h_2 = 585\,49955.614 = \frac{44}{13} \cdot h_1,$$

respectively; therefore the pattern \mathbf{c}_2 is expected to occur 3.4 times more often than \mathbf{c}_1 . The average number of occurrences of \mathbf{c}_2 in a large interval Δ near a much larger x is about $h_2 \Delta (\log x)^{-19}$. From this we infer that the chances of the pattern \mathbf{c}_2 reoccurring at an $x < 5 \cdot 10^{25}$ are about 93%. This is also the chance of its mirror image occurring in the same range.

In the Initial Report 2001, [Projects/clprimes01.pdf](#), the idea of a search for a 19-tuple of maximum density was rejected because at that time the available hardware (the idle time of 128 nodes of Asgard) would have required an excessively long search time (up to five years).

With the installation of Hreidar the situation has changed. With the idle time of 64 nodes of Hreidar available, the speed increased about 5-fold. The quest for a 19-tuple of maximum density is on again!

We are again working in blocks of length $\Delta = 10^{24}$. In an initial attempt we were directly heading for one of the more abundant patterns, \mathbf{c}_2 . This resulted in a computation time of about 28 days per block. Unfortunately, 7 blocks done in this way produced no output. Of course, these 7 blocks have been recorded as non-carriers of Pattern \mathbf{c}_2 , but obviously a change of strategy was needed.

We decided to look for the sub-patterns

$$\mathbf{C}_2 = [41, 43, \dots, 107, 109], \quad \mathbf{C}'_2 = [-109, -107, \dots, -43, -41]$$

of 17 elements, obtained by deleting the two boundary elements from the original patterns \mathbf{c}_2 , \mathbf{c}'_2 . Deleting any other two elements from the original patterns would have resulted in a larger computational effort. From a list of subpatterns the original patterns may easily be sorted out by checking the deleted elements for primality.

One advantage of this strategy is that output is generated at a reasonable rate. With the Hardy-Littlewood constant $h = 31\,75403.027$ the estimated

number $N(x) \approx h \Delta (\log x)^{-17}$ of occurrences in a block of length $\Delta = 10^{24}$ centered at $x = X \cdot 10^{24}$ is given in the following table.

| X | +0.5 | +1.5 | +2.5 | +3.5 | +4.5 | +5.5 | +6.5 | +7.5 | +8.5 | +9.5 | sum |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 0 | 11.110 | 6.7601 | 5.7584 | 5.1955 | 4.8154 | 4.5338 | 4.3130 | 4.1330 | 3.9822 | 3.8531 | 54.45 |
| 10 | 3.7408 | 3.6417 | 3.5533 | 3.4738 | 3.4016 | 3.3357 | 3.2751 | 3.2192 | 3.1673 | 3.1190 | 88.38 |
| 20 | 3.0738 | 3.0314 | 2.9915 | 2.9538 | 2.9182 | 2.8845 | 2.8524 | 2.8219 | 2.7928 | 2.7650 | 117.47 |
| 30 | 2.7384 | 2.7130 | 2.6885 | 2.6651 | 2.6425 | 2.6208 | 2.5998 | 2.5796 | 2.5601 | 2.5412 | 143.82 |
| 40 | 2.5229 | 2.5053 | 2.4881 | 2.4715 | 2.4554 | 2.4397 | 2.4245 | 2.4097 | 2.3954 | 2.3813 | 168.31 |
| 50 | 2.3677 | 2.3544 | 2.3414 | 2.3288 | 2.3164 | 2.3044 | 2.2926 | 2.2811 | 2.2699 | 2.2589 | 191.43 |
| 60 | 2.2481 | 2.2376 | 2.2273 | 2.2172 | 2.2073 | 2.1976 | 2.1881 | 2.1788 | 2.1697 | 2.1607 | 213.46 |
| 70 | 2.1519 | 2.1433 | 2.1348 | 2.1264 | 2.1183 | 2.1102 | 2.1023 | 2.0945 | 2.0869 | 2.0794 | 234.61 |
| 80 | 2.0720 | 2.0647 | 2.0575 | 2.0505 | 2.0436 | 2.0367 | 2.0300 | 2.0234 | 2.0168 | 2.0104 | 255.01 |
| 90 | 2.0040 | 1.9978 | 1.9916 | 1.9855 | 1.9795 | 1.9736 | 1.9678 | 1.9620 | 1.9564 | 1.9508 | 274.78 |
| 100 | 1.9452 | 1.9398 | 1.9344 | 1.9291 | 1.9239 | 1.9186 | 1.9135 | 1.9084 | 1.9034 | 1.8985 | 294.00 |
| 110 | 1.8936 | 1.8887 | 1.8840 | 1.8793 | 1.8746 | 1.8700 | 1.8654 | 1.8609 | 1.8564 | 1.8520 | 312.72 |
| 120 | 1.8477 | 1.8434 | 1.8391 | 1.8349 | 1.8307 | 1.8265 | 1.8224 | 1.8184 | 1.8144 | 1.8104 | 331.01 |

The second advantage is the possibility of comparing actual counts with the HL estimates; in this way one might get an idea which pattern will occur earlier. Since the distribution of long patterns tends to have large-scale anomalies, this argument – although not well founded – may sometimes work. The price for these benefits is a somewhat larger computational effort: one block of length $\Delta = 10^{24}$ roughly takes 38 days (instead of 28 days for \mathbf{c}_2).

In the table below we summarize the current status (February 24, 2016) of the search for \mathbf{C}_2 and \mathbf{C}'_2 in the first 142 blocks. A digit in the table indicates the number of patterns \mathbf{C}_2 or \mathbf{C}'_2 located in the corresponding block. A question mark (?) means: the block has been screened for the 19-tuple \mathbf{c}_2 with a *negative* result; the number of 17-tuples \mathbf{C}_2 in the corresponding block is not known. Assuming an average turnout of 44 % (see below) we estimate the total number in the first 3 and the remaining 4 blocks of this type to be about 8 and 6, respectively. In all other cases (+) nothing has been done so far.


| | | | | | | | | | | |
|-----------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Blocks | 1-5 | 6-10 | 11-15 | 16-20 | 21-25 | 26-30 | 31-35 | 36-40 | 41-45 | 46-50 |
| \mathbf{C}_2 | 6???3 | 1++++ | ??+++ | +++++ | +++++ | +++?? | +++++ | +++++ | +++++ | +++++ |
| \mathbf{C}'_2 | 92466 | 25654 | 34422 | 73353 | 43221 | 51325 | 16034 | 22533 | 25032 | 28223 |

| | | | | | | | | | | |
|-----------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------|
| Blocks | 51-55 | 56-60 | 61-65 | 66-70 | 71-75 | 76-80 | 81-85 | 86-90 | 91-95 | 96-100 |
| \mathbf{C}'_2 | 64421 | 45062 | 20122 | 00215 | 40103 | 20113 | 02121 | 52242 | 01530 | 34342 |

| | | | | | | | | | | |
|-----------------|--------|-------|-------|-------|-------|-------|-------|-------|-------|--------|
| Blocks | 101-05 | 06-10 | 11-15 | 16-20 | 21-25 | 26-30 | 31-35 | 36-40 | 41-45 | 46-150 |
| \mathbf{C}'_2 | 14103 | 12301 | 12211 | 21001 | 32121 | 11112 | 21221 | 11242 | 33+++ | +++++ |

We infer that the first pattern is deficient in the regions considered; C_2 , C'_2 produce about 44 % or 97 %, respectively, of the expected number of patterns. Not too bad a sign for continuing the search for a 19-tuple with the pattern C'_2 !

Being unable to report a 19-tuple (among 77 integers) at this time, we will at least report the initial primes of all 357 subpatterns (17 primes among 69 integers) found so far. In the figure below the number of patterns C'_2 is plotted versus the initial prime x of the pattern, together with the Hardy-Littlewood estimate $HL(x)$. The locations of Patterns [-113..-41] with an additional prime at a distance of 4 on the lower side are marked with an asterisk; a circle marks Patterns [-109..-37].



In the Appendix all initial primes of Patterns C_2 and C'_2 found so far are listed. The “near misses” with 18 primes among 73 integers are marked

by asterisks. The position of the asterisk (left or right) indicates the side of the pattern (lower or upper) carrying the additional prime at a distance of 4. Hence the instance near $4.52 \cdot 10^{23}$ has the pattern of [41..113]; the one near $1.27 \cdot 10^{23}$ begins 4 units earlier and has the pattern of [-113..-41]. The instances near $3.24 \cdot 10^{24}$, $3.38 \cdot 10^{24}$ and $4.52 \cdot 10^{24}$ have the pattern of [-109..-37]. We are waiting for an entry framed by two asterisks.

References

- [1] C. Batut, K. Belabas, D. Bernardi, H. Cohen, M. Olivier: *The software package PARI* (freeware). <http://pari.math.u-bordeaux.fr/>
- [2] ETH Life, Die tägliche Webzeitung der ETH, 6. Dezember 2000, Archiv. *Prozessoren malochten 100 Tage*. <http://www.ethlife.ethz.ch/articles/tages/Waldvogel.html>
- [3] Tony Forbes: *Prime clusters and Cunningham chains*. Math. of Comp. **68** (1999) 1739-1747.
- [4] Tony Forbes: *Prime k-tuplets*. <http://anthony.d.forbes.googlepages.com/ktuplets.htm>
- [5] G.H. Hardy and J.E. Littlewood: *Some problems of Partitio Numerorum III*. Acta Math. **44**, 1922, 1-70.
- [6] *Number theory news*. <http://www.utm.edu/research/primes/>
- [7] Paulo Ribenboim: *The New Book of Prime Number Records*, 3rd ed. Springer 1996, 541 pp.
- [8] Hans Riesel: *Prime Numbers and Computer Methods for Factorization*, 2nd ed. Birkhäuser 1994, 464 pp.
- [9] R.L. Rivest, A. Shamir, L. Adleman: *A method for obtaining digital signatures and public-key cryptosystems*. Communications of the ACM **21** (1978) 121-126.
- [10] J. Waldvogel: *Homepage*. <http://www.math.ethz.ch/~waldvoge/Projects/>

Appendix: Subpatterns found so far

| C2=[41..109] | | C2'=[-109..-41] | |
|--------------|---------------------------|-----------------|----------------------------|
| 1 | 41 | 1 | 71542018620258822095351 |
| 2 | 112650791633055206521031 | 2 | 110251111115724441042071 |
| 3 | 272064438851059718733911 | 3 | *127372818047327460718001 |
| 4 | 452098732111641436261481* | 4 | 359103301162581793637711 |
| 5 | 636277672291047674637791 | 5 | 472773823866578374260011 |
| 6 | 649410249467730311490971 | 6 | 668134176357714775428611 |
| | | 7 | 842902600507150788234521 |
| | | 8 | 979634834795806543814351 |
| | | 9 | 980351128284287125225751 |
| | | 10 | 1448301799940226082771811 |
| | | 11 | 1791530392243173635753591 |
| | | 12 | 2287676808438268517289911 |
| | | 13 | 2638263731291524361295161 |
| | | 14 | 2861160868941024590320271 |
| | | 15 | 2965402830695807653891631 |
| | | 16 | 3239304711211315285839191* |
| | | 17 | 3381058510131934598816741* |
| | | 18 | 3491478433645050966237011 |
| | | 19 | 3663823986030922159656101 |
| | | 20 | 3774602330694614789940461 |
| | | 21 | 3878542221422020936683791 |
| 1 | 4169958545122711795353941 | 22 | 4184454523805320162938431 |
| 2 | 4262842381283978056404881 | 23 | 4225209451302652505772971 |
| 3 | 4355781833464360551222011 | 24 | 4386179912950387994201651 |
| | | 25 | 4522878605388339260634461* |
| 4 | 5332857791735927206954241 | 26 | 4716202264464890619359831 |
| | | 27 | 4869723749184137214270791 |
| | | 28 | *5188248562214464094307851 |
| | | 29 | 5413131850555132198264661 |
| | | 30 | 6423359383060950425435231 |
| | | 31 | 6462943397426544261913571 |
| | | 32 | 6515331587478752597979311 |
| | | 33 | 6657401626856646647681591 |
| | | 34 | 6732639026163593595722561 |

C2'=[-109..-41]

35 7068848143689559074537581
36 7069902747669768688311521
37 7106614823259270051322481
38 7167564348232179592490681
39 7518303244972092457804811
40 7558395658997171017779641

41 8019983382743326416141611
42 8351014354384188300523601
43 8394208478031243473535821
44 8796828522725933171551961
45 8941097330542117721700611

46 9451163321791831501251281
47 *9867358577329746770637251
48 9980642183707047492649901
49 9982676186512737822143921

50 10384473055093816251062381
51 10508599120346583663997481
52 10809519785591093308761911

53 11360432123283992291743091
54 11633244241482020341414121
55 11836184125186489787548721
56 11985312529091443124703011

57 12110376481437954916306001
58 12276728127624397775858411
59 12314036812755277924784351
60 12685651643704874036352041

61 13114239319848333922324961
62 13627173426996373227623531

63 14323963076718850133719031
64 14730444793496584954251341

65 15307029480404445010527191
66 15399775927777730924905361
67 15414617623013648097848681
68 15426394378504824654259211
69 15700071010820082671524631
70*15722057448876967802383661
71 15726957442202271450775481

C2'=[-109..-41]

72 16152301493077394927542151
73 16423284953192882928833561
74 16906332397375773026034251

75 17173948166155769230872521
76 17209884144375743712871871
77 17943521643579660773334581

78 18322687280025547085907131
79 18373102734662381624408381
80 18453788042746003529722031
81 18473759960272367077727291
82 18561322841562422318937431

83 19583322299892820386527501
84 19701331508947654781495171
85 19848748297189344094491821

86 20554940850499841671725731
87 20578602413614392636038441
88 20620171249006398317408051*
89 20698687588834882254127391

90 21111317185520011816015541
91 21156265866707710160101571*
92 21800648661381114847852421

93 22245161817448111292030171
94 22354968551813917951601561

95 23497539251772545287798361
96 23903291173457833917479591

97 24705065946193416530736881

98*25234150994366362207739471
99 25263664554515461668503801
100 25347043389795198872250341*
101 25726650749602154810197631
102 25947645160924472931625571

103 26215385108311265356059461

104 27435503504282409530404301
105 27804461307247785171181061
106 27992165933083119142384901*

C2'=[-109..-41]

107 28587021849984636782823971
108 28725150350219399690954231

109 29004431262862944827208161
110 29074293657139332704181971
111 29277232560082331577572711
112 29556637195454908075096451
113 29830012818163367509479671

114 30135761176379845423617071

115 31106642710562216131023581*
116 31510656371837402820570401
117 31608481544717546095066331
118 31822805936385714871330601
119 31833515366979568095753791
120 31952914340689883091961301

121 33829984544463298215414281
122 33913658953315256103198101
123 33976782450455103262193081

124 34334530034601738614102321
125 34684881315519357142930601
126 34693709964786919060614491*
127 34726172589975186357467321

128 35206610962106249577971951
129 35294838755236934864120021

130 36161238381434666136466721
131 36804465681616726803009191

132 37075366741403316541274951
133 37172881033460259725948201
134 37351011223126811387256971
135 37582909908839139171356921
136 37775177885317182474574301*

137 38248491989646598057728221
138 38290339015423224921781481
139 38637290403827870627172191

C2'=[-109..-41]

140 39420784963838416751763041
141 39658640177366254707662891
142 39968706603255248587728851

143 40354075779696330156755141
144 40477618815825993983000711

145 41190653587018584204355691
146 41264351829674953112072831
147 41539341349561725519019061
148 41650534847605510089523151
149 41989356607743797871598871

150*43057544631003403144463711
151*43081312682299678694677661
152 43639771287915229286719751

153 44074115153125883294198621
154 44990099513628683912367761

155 45194463980789514206295371
156 45823488455412958189729721

157 46046178308566762114318421
158 46338719607904299051165581
159 46382445063348951534677591
160 46411470092639897829243551
161 46535014899155334046114511
162 46724900004576789333778331
163 46737381948653913664277021
164 46924193138242798802093561

165 47319504598099378910445071
166 47941602150408861630295451

167 48460984964043175125051341
168 48977382368352401727753881

169 49204467559453403756919251
170 49380784584721484163799421
171 49614367573544827286513171

C2'=[-109..-41]

172 50347935949163794523932241
173 50417056266422176534833521
174 50488211153394006095463251
175 50538768446856074445690521
176 50686159063842560657028521
177 50998141230342992907120791

178 51070898064266935599875021
179 51125255743844986826912681
180 51285415832156224102724231
181 51845170207066461273918761

182 52289236400633467028713811
183 52627557253361499365946881*
184 52733694034272812983872341
185*52739555754897822763982921

186 53407151211001995334827791
187 53699155143112186922273951
188 54396744504388395445315871

189 55188915736870184544985601
190 55239999069804832367614421
191 55294887549508887603823391
192*55497352712075821192395641

193 56010289873072396015354091
194 56554948329052098487101431*
195 56626029403506045980777171
196 56642147704707425175975551
197 56878958227279168894733711

198*58152298245071252378776511
199 58294052542943324304066761
200 58315179945493641391070741
201 58369684176795608197370681
202 58452250931326222211691341
203 58746285490202984469986081

204 59254704899192057469313991
205 59707386038150921446876391

C2'=[-109..-41]

206 60304943454164475662515811
207 60345830722463589716392871

208 62211719897992766461309841

209 63272429276964882378544031
210 63903650083918050249763811*

211 64134699112056277346421671
212 64346006281871430315809501*

213 67561119397307272642795781
214 67794768223026847025213081

215 68043200340623206051953491

216 69104237948198904502931771
217 69332569607885253700760621
218 69392624402544960288451571
219 69600129710869392547182161
220 69932640879222499484105891

221 70336143673221212631084131
222 70459639345868511686577521
223 70499479425744733653419021
224 70989971724968493144557501

225 72734592371778625139601641

226 74211979717443719191660511
227 74886662719031724143222531
228 74928553149966759228885791

229 75130928599154991515842781
230 75967291917711703674164891

231 77435265650320546710691841

232 78095290682304930734579771

233 79552926097713001543119401
234 79623643523636900388931121
235 79743356926907028732517031

C2'=[-109..-41]

236 81790103060147606413849451
237*81928700810343676214635601

238 82535645126695012712880311

239 83163723992167497560211221
240 83200847503806790467069251

241 84757094727049674913002911

242 85235564164289258392843061
243 85267341076275051682057961
244 85389220242535462976082251
245 85799800160042023928322341*
246 85984419704737637198144921

247 86386503913904177739725861
248 86423046942480960718526411

249 87105045069546822910581881
250 87265775628916512266839091

251 88463224850998331042678861
252 88739768237002271090001191
253 88755597497974393054188011
254 88925664871899160394339141

255 89245134321229378500478361
256 89886550819541134968699281*

257 91450674254305715042573771

258 92037088913118518577317801
259 92099516902136090302205141
260 92101045501758290042305151
261 92196665663419186384604981
262 92480366996751441113616101

263 93056813193542798564164091
264 93606240245355109740618041
265 93819139199993179590677951*

C2'=[-109..-41]

266 95247248197635330180534041
267 95536904259961414743232631
268 95588894891587566449009921

269 96620942409228127360499921
270 96632388117810924397238111
271 *96822528705764010248629421
272 96957752342065788532028561

273 *97185033888313518629942351
274 97897240071889550869026341
275 97990071755353143269543531

276 98285596810503098975348261
277 98911690780929584204026271
278 *98939393003568465298240181
279 98956470454266970033829231

280 99288173733479552086675301
281 99767083494105820431999251

282 100920514955514684696796421

283 101012387005596252151255181
284 101190723975134428857942611*
285 101532044207218432360212551
286 101630255942059554216146531

287 102996598093096156644194891

288 104325378942756900925326221
289 104350285223660019868484051
290 104863834898014649651804501

291 105256141780180383585853151

292 106323683748837481199653271
293 106378402740750551747986541

294 107006105031999674188287491
295 107080299048686295009137351
296 107121110829656976022940861

| C2'=[-109..-41] | | C2'=[-109..-41] | |
|---------------------------------|-----------------------------|-----------------|------------------------------|
| 297 | 109213174969271098379583821 | 321 | 128258283883321490314656131 |
| 298 | 110200631471916014446877351 | 322 | 129147422114266372835458331 |
| 299 | 111004772046624579644110481 | 323 | 129991530649412842189729841 |
| 300 | 111781115172101949314370521 | 324 | 130141997309277697869235001 |
| 301 | 112011502056952356092854511 | 325 | 130617237643414983513932771 |
| 302 | 112130618794503459819398621 | 326 | 131417312794605480668475341 |
| 303 | 113700965778596253535017431 | 327 | 132217273862041931492499461 |
| 304 | 114653522511334193984286581 | 328 | 132295304994724352125654901 |
| 305 | 115372763336437245111072491 | 329 | 133168830422865624613666331 |
| 306 | 115862731750447659968270171 | 330 | 133221121010717785317291911 |
| 307*116970520931984620036816571 | | 331 | 134814184120002985070663081 |
| 308*119350280243928922603134611 | | 332 | 135465565864967513782596881 |
| 309 | 120329545398544404239146721 | 333 | 136154045050562714408062721 |
| 310 | 120529078430374537786952981 | 334 | 137854928297624635154857691 |
| 311 | 120737348813410424444589131 | 335 | 137958674217592315810642811 |
| 312 | 121066461556410591946064231 | 336 | 138201829509410126825705231 |
| 313 | 121383609931718248199865671 | 337 | 138585760086748307155542611 |
| 314 | 122435787700186403915321741 | 338 | 138601732008002676622689371 |
| 315 | 123204742436259254249128541 | 339* | 138608012676032357413775711 |
| 316 | 123337494358235211025381301 | 340 | 139305641428837702849340201 |
| 317 | 124013212267672524347831171 | 341 | 139636066510349396136649391 |
| 318 | 125459860488089856533024111 | 342 | 140207786639766762247801361 |
| 319 | 126917233130013605562352961 | 343 | 140385237967708813985355341 |
| 320 | 127720949153422449334760141 | 344 | 140440881989219134464080831 |
| | | 345 | 141135367086076411070442311 |
| | | 346 | 141430590840510059626978331* |
| | | 347 | 141589445556152778750375701 |